

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number
WO 00/78971 A2

(51) International Patent Classification⁷: C12N 15/52,
C07K 14/705, C12Q 1/68, C07K 16/18, A61K 38/00,
C12N 5/10, A61K 48/00, G01N 33/50, C12N 15/86

Avenue, Palo Alto, CA 94306 (US). ORAM, John, F.
[—/US]; 1738 N.E. Naomi Place, Seattle, WA 98195
(US). GARVIN, Michael [—/US]; 17 White Street, San
Francisco, CA 94019 (US).

(21) International Application Number: PCT/US00/16591

(22) International Filing Date: 16 June 2000 (16.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/140,264 18 June 1999 (18.06.1999) US
60/153,872 14 September 1999 (14.09.1999) US
60/166,573 19 November 1999 (19.11.1999) US

(74) Agent: HUGHES, A., Blair; McDonnell Boehnen Hulbert & Berghoff, 32nd Floor, 300 South Wacker Drive, Chicago, IL 60606 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicants (*for all designated States except US*): CV THERAPEUTICS, INC. [US/US]; 3172 Porter Drive, Palo Alto, CA 94304 (US). UNIVERSITY OF WASHINGTON [US/US]; Office of Technology Transfer, Suite 200, 1107 N.E. Street, Seattle, WA 98195 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): LAWN, Richard, M. [—/US]; 1927 8th Avenue, San Francisco, CA 94116 (US). WADE, David [—/US]; Apartment L, 383 Curtner


Published:

— Without international search report and to be republished upon receipt of that report.

[Continued on next page]

(54) Title: COMPOSITIONS AND METHODS FOR INCREASING CHOLESTEROL EFFLUX AND RAISING HDL USING ATP BINDING CASSETTE TRANSPORTER PROTEIN ABC1

CVT ABCA1 Gene Structure



(57) Abstract: The present invention relates to novel ABC1 polypeptides and nucleic acid molecules encoding the same. The invention also relates to recombinant vectors, host cells, and compositions comprising ABC1 polynucleotides, as well as to methods for producing ABC1 polypeptides. The invention also relates to antibodies that bind specifically to ABC1 polypeptides. In addition, the invention relates to methods for increasing cholesterol efflux as well as to methods for increasing ABC1 expression and activity. The present invention further relates to methods for identifying compounds that modulate the expression of ABC1 and methods for detecting the comparative level of ABC1 polypeptides and polynucleotides in a mammalian subject. The present invention also provides kits and compositions suitable for screening compounds to determine the ABC1 expression modulating activity of the compound, as well as kits and compositions suitable to determine whether a compound modulates ABC1-dependent cholesterol efflux.

WO 00/78971 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

COMPOSITIONS AND METHODS FOR INCREASING CHOLESTEROL EFFLUX AND RAISING HDL USING ATP BINDING CASSETTE TRANSPORTER PROTEIN ABC1

This application claims priority to U.S. provisional application no. 60/140,264, filed on
5 June 18, 1999; U.S. provisional application no. 60/153,872, filed on September 14, 1999; and
U.S. provisional application no. 60/166,573, filed on November 19, 1999.

TECHNICAL FIELD OF INVENTION

The present invention relates to novel ABC1 polypeptides and nucleic acid molecules
10 encoding the same. The invention also relates to recombinant vectors, host cells, and
compositions comprising ABC1 polynucleotides, as well as to methods for producing ABC1
polypeptides. The invention also relates to antibodies that bind specifically to ABC1
polypeptides. In addition, the invention relates to methods for increasing cholesterol efflux as
well as to methods for increasing ABC1 expression and activity. The present invention further
15 relates to methods for identifying compounds that modulate the expression of ABC1 and
methods for detecting the comparative level of ABC1 polypeptides and polynucleotides in a
mammalian subject. The present invention also provides kits and compositions suitable for
screening compounds to determine the ABC1 expression modulating activity of the compound,
as well as kits and compositions suitable to determine whether a compound modulates ABC1-
20 dependent cholesterol efflux.

BACKGROUND OF THE INVENTION

Circulating lipids in human plasma or lymphatic fluid consist of cholesterol, cholesteryl
esters, triglycerides and phospholipids. These lipids are transported in large molecular
25 complexes called lipoproteins, which consist of a core of cholesteryl esters and/or triglycerides,
an envelope of phospholipids and free cholesterol, and apolipoproteins (Scriber et al., Eds., *The
Metabolic and Molecular Basis of Inherited Disease*, 7th Ed., p.1841-1851 (McGraw-Hill, New
York 1995)). Apolipoproteins are involved in the assembly and secretion of the lipoprotein, as
well as the activation of lipoprotein modifying enzymes, such as lecithin cholesterol acyl
30 transferase (LCAT). In addition, apolipoproteins provide structural integrity and are ligands for
a large spectrum of receptors and membrane docking proteins. The plasma lipoproteins are
categorized into five types according to size: chylomicrons (largest in size and lowest in density),
very low density lipoproteins (VLDL), intermediate density lipoproteins (IDL), low density

lipoproteins (LDL) and high density lipoprotein (HDL).

Chylomicrons, VLDLs, IDLs, and LDLs transport exogenous and endogenous cholesterol and triacylglycerols to peripheral sites, where the lipids play a role in various metabolic pathways and serve as a major constituent of cell membranes. Chylomicrons are assembled in the intestinal mucosa as a means to transport dietary cholesterol and triacylglycerols to various tissues. VLDLs are formed in the liver to transport endogenous cholesterol and triacylglycerols synthesized by the liver to extra-hepatic tissues, such as muscle and adipose tissue. In fasting serum, VLDLs contain 10-15% of the total serum cholesterol and most of the triglyceride. In circulation, VLDLs are converted to LDLs through the action of lipoprotein lipase. LDLs are the primary plasma carriers of cholesterol for delivery to all tissues, typically containing 60-70% of the total fasting serum cholesterol.

In contrast, HDLs are involved in "reverse cholesterol transport", the pathway by which excess cholesterol is transported from peripheral sites back to the liver, where it is excreted in the form of bile salts (Glomset, J.A., *J. Lipid Res.*, 9, 155-167 (1968)). Nascent HDLs are synthesized *de novo* in the liver and small intestine, as protein-rich disc-shaped particles devoid of cholesterol and cholesterol esters. In fact, a major function of HDLs is to act as circulating stores of apolipoproteins, primarily apo C-I, apo C-II, and apoE. The nascent or protein-rich HDLs are converted into spherical lipoprotein particles through the accumulation of cholesteryl esters obtained from cellular sources. The HDL normally contain 20-30% of the total fasting serum cholesterol.

According to current theories, the reverse efflux of cellular cholesterol to HDL is mediated through two mechanisms: an aqueous diffusion pathway and an apolipoprotein-mediated pathway. The relative importance of these distinguishable mechanisms depends on the cell type and metabolic state (Oram et al., *J. Lipid Res.*, 37:2743-2491 (1996); Rothblat et al., *J. Lipid Res.*, 40:781-796 (1999); Stein et al., *Atherosclerosis*, 144:285-301 (1999)). For many cells, the aqueous diffusion pathway is the principle pathway through which cholesterol efflux occurs (Johnson et al., *Biochim. Biophys. Acta*, 1085:273-298 (1991)). This pathway involves the bidirectional exchange of cholesterol between cell membranes and a lipoprotein acceptor, such as HDL, in the extracellular space through a process of passive transport (Remaley et al., *Arterioscler. Thromb. Vasc. Biol.*, 17:1813-1821 (1997); Rothblat et al., *J. Lip. Res.*, 40:781-796 (1999)). The exchange may occur primarily at surface microdomains known as caveolae (Fielding et al., *Biochemistry*, 34:14288-14292 91995)). Net efflux can be driven by conversion of cholesterol in the extracellular compartment to cholesteryl ester by the action of LCAT.

Alternatively, in macrophage and fibroblast cells, cholesterol and phospholipid efflux is primarily mediated through apolipoproteins, such as apo A-I, apo A-II, and Apo E (Remaley, *supra* (1997); Francis, et al., *J. Clin. Invest.*, 96:78-87 (1995); Vega et al., *J. Intern. Med.*, 226:5-15 (1989); Sakar et al., *Biochim. Biophys. Acta*, 1438: 85-98 (1999); Hara et al., *J. Biol. Chem.*, 266:3080-3086 (1991); Fielding et al., *J. Lipid Res.*, 38, 1503-1521 (1997); Oram et al., *J. Lipid Res.*, 37, 2743-2491 (1996)). The process of apolipoprotein-mediated lipid efflux particularly dominates in macrophages and other scavenger cells when they are cholesterol-loaded and/or growth-arrested. Apolipoprotein-mediated efflux is an active transport process that requires the direct interaction of the apolipoprotein with the cell surface, the lipidation of the apolipoprotein, and the subsequent dissociation of the lipid-apolipoprotein particle from the cell (Oram, *supra* (1996); Mendez, A.J., *J. Lipid Res.*, 38, 1807-1821 (1997); Remaley, *supra* (1997); Mendez, A.J., *J. Lipid Res.*, 37, 2510-2524 (1996)). Once removed from the cell, the cholesterol-rich HDL particles are transported to the liver and removed from the body as described.

Abnormal lipoprotein function and/or metabolism resulting from genetic defect or as a secondary effect of another disorder can have serious biological consequences. In addition to dietary influences, disorders such as diabetes, hypothyroidism, and liver disease can result in elevated plasma levels of LDL-cholesterol and triglycerides. Elevated levels of LDL-cholesterol and triglycerides have been identified as major risk factors associated with the incidence of coronary heart disease, which is the primary cause of death in the United States and other industrialized nations (Hokanson et al., *J. Cardiovasc. Risk.*, 3:213-219 (1996); The Expert Panel, *JAMA*, 269:3015-3023 (1993)). The accumulation of excess LDL-cholesterol on arterial walls can lead to the formation of atherosclerotic plaques, which play a major role in the development of heart disease. A plaque is believed to form when free radicals released from arterial walls oxidize LDL. According to theory, the oxidized form of LDL triggers an inflammatory response, attracting circulating cells to the site which contribute to the formation of a lipid plaque. Among these are macrophages and other cells that contain scavenger receptors that accumulate cholesterol in an unregulated manner (Brown et al., *Ann. Rev. Biochem.*, 52:223-261 (1986)). Vast stores of internal cholesterol result in conversion to a foam cell phenotype, which is believed to be a major contributor to the development of vascular lesions. As the plaque builds up, the arterial walls constrict, reducing blood flow to the heart.

Interestingly, however, an estimated 60% of heart attacks occur in persons who do not have elevated blood levels of LDL-cholesterol. Of these, an estimated 45% are associated with below average blood levels of HDL-cholesterol, indicating that low HDL-cholesterol level is a

significant risk factor for coronary heart disease. In fact, recent studies have indicated that a decreased HDL-cholesterol level is the most common lipoprotein abnormality seen in patients with premature coronary artery disease (Genest J., *Circulation*, 85:2025-2033 (1992); Genest et al., *Arterioscler. Thromb.*, 13:1728-1737 (1993)). Although the basis for the inverse association
5 between HDL-cholesterol and coronary heart disease is not well understood, it has been suggested that the cardioprotective role of HDL may stem from its activity relating to the promotion of cholesterol efflux from macrophage foam cells in atherosclerotic lesions.

One example of cardiovascular disease associated with low HDL is Tangier disease (TD), a rare genetic disorder characterized by a near or complete absence of circulating HDL. In
10 addition to near zero plasma levels of HDL, patients with TD have a massive deposition and accumulation of cholesteryl esters in several tissues, including tonsils, lymph nodes, liver, spleen, thymus, intestine, and Schwann cells (Fredrickson, D.S., *J. Clin. Invest.*, 43, 228-236 (1964); Assmann et al., *The Metabolic Basis of Inherited Disease*, (McGraw-Hill, New York, 1995)). Although the cellular mechanisms have not been previously identified, recent studies
15 have shown that cells from subjects with TD are defective in the process of apolipoprotein-mediated removal of cholesterol and phospholipids (Remaley et al., *Arterioscler. Thromb. Vasc. Biol.*, 17, 1813-1821 (1997); Francis et al., *J. Clin. Invest.*, 96, 78-87 (1995); Rogler et al., *Arterioscler. Thromb. Vasc. Biol.*, 15, 683-690 (1995)). These results have led to the proposal that the severe HDL deficiency in TD patients stems from the inability of nascent apo A-I to
20 acquire lipids. Because they do not mature into lipid-rich particles, the nascent HDL in TD patients is rapidly catabolized and removed from the plasma, resulting in the near zero levels of circulating HDL (Remaley, *supra* (1997); Francis, *supra* (1995); Horowitz et al., *J. Clin. Invest.*, 91, 1743-1752 (1993); Schaefer et al., *J. Lip. Res.*, 22:217-228 (1981)).

Other disorders associated with severe premature atherosclerosis and high risk for
25 coronary heart disease resulting from diminished HDL-cholesterol levels are hypoalphalipoproteinemia and familial HDL deficiency syndrome (FHA). Persons with these disorders often have normal LDL-cholesterol and triglyceride levels. In addition, disorders such as diabetes, alcoholism, hypothyroidism, liver disease, and elevated blood pressure can result in diminished plasma levels of HDL-cholesterol, although many of these disorders are also
30 accompanied by elevated LDL-cholesterol and triglyceride levels.

Current treatments for coronary heart disease have focused primarily on diet manipulations and/or drug therapies aimed at lowering the plasma level of LDL-cholesterol by inhibiting LDL secretion or promoting LDL turnover. Derivatives of fibric acid, such as

clofibrate, gemfibrozil, and fenofibrate, promote rapid VLDL turnover by activating lipoprotein lipase. Nicotinic acid reduces plasma levels of VLDL and LDL by inhibiting hepatic VLDL secretion. In addition, HMG-CoA reductase inhibitors, such as mevinolin, mevastatin, pravastatin, simvastatin, fluvastatin, and lovastatin reduce plasma LDL levels by inhibiting the intracellular synthesis of cholesterol, which causes an increase in the cellular uptake of LDL. In addition, bile acid-binding resins, such as cholestyrene, colestipol and probucol decrease the level of LDL-cholesterol by increasing the catabolism of LDL-cholesterol in the liver.

However, many of these therapies are associated with low efficacy and/or side effects that may prevent long-term use. For example, use of HMG-CoA reductase inhibitors carry a significant risk of toxicity because they inhibit the synthesis of mevalonate, which is required for the synthesis of other important isoprenoid compounds in addition to cholesterol. Also, gemfibrozil and nicotinic acid are associated with serious adverse effects, including renal injury, myopathy, myoglobinuria and intolerable skin flushing and itching. In addition, the role of probucol in treating patients with coronary heart disease is uncertain because its administration results in lower HDL-cholesterol levels as a side effect of reducing LDL-cholesterol.

Furthermore, treating patients who have isolated low HDL-cholesterol levels provides a particularly difficult therapeutic challenge. For instance, patients with Tangier disease exhibit a 4- to 6-fold increase in cardiovascular disease even though their LDL levels are already reduced by about 50%. While there is some evidence that gemfibrozil and nicotinic acid may simultaneously elevate HDL levels, in general, therapies aimed at lowering plasma LDL-cholesterol levels are not effective for Tangier patients who suffer from coronary heart disease as a result of diminished HDL levels. Likewise, patients with hypoalphalipoproteinemia, familial HDL deficiency syndrome, or other cardiovascular disease resulting from low levels of HDL will not benefit from therapies aimed at lowering the level of plasma LDL.

The problems associated with current therapies for cardiovascular disease stem partially from the fact that the biology involved in the movement of cholesterol in and out of cells is not fully understood. Furthermore, the proteins that play a role in cholesterol movement are not fully known. Therefore, there remains a need for a better understanding of cholesterol cell biology, as well as new methods for treating humans suffering from cardiovascular disease and other disorders associated with hypercholesterolemia. Additionally, there remains a need for new methods of diagnosing cardiovascular disease and new methods of screening patients to identify those at high risk for developing cardiovascular disease.

The identification of genes and proteins involved in cholesterol transport would be useful

in the development of pharmaceutical agents for the treatment of heart disease and other disorders associated with hypercholesterolemia and atherosclerosis. In addition, the identification of such genes would be useful in the development of screening assays to screen for compounds that regulate the expression of genes associated with cholesterol transport. The
5 identification of such regulatory compounds would be useful in the development of further therapeutic agents. Furthermore, the identification of genes and proteins involved in cholesterol transport would be useful as diagnostic indicators of cardiovascular disease and other disorders associated with hypercholesterolemia.

10 SUMMARY OF THE INVENTION

The present invention provides novel polypeptides and polynucleotides involved in cholesterol efflux. Specifically, the present invention provides novel ATP-Binding Cassette (ABC1) polypeptides and novel polynucleotides that encode ABC1 polypeptides. The terms "ABC1" and "ABCA1" are alternative names for the same ATP-Binding Cassette protein and
15 gene. The invention provides ABC1 polypeptides, polypeptide fragments, and polypeptide variants. In one preferred embodiment, the present invention provides an isolated polypeptide comprising SEQ ID NO: 2. In another preferred embodiment, the present invention provides an isolated polypeptide comprising an amino acid sequence that has at least 98% identity to SEQ ID NO: 2. The present invention also provides ABC1 polypeptides from Tangier disease patients.
20 In one preferred embodiment, the present invention provides an isolated polypeptide comprising SEQ ID NO: 8. In another preferred embodiment, the present invention provides an isolated polypeptide comprising SEQ ID NO: 10.

In addition, the present invention provides ABC1 polynucleotides, polynucleotide fragments, and polynucleotide variants. In one preferred embodiment, the present invention
25 provides an isolated polynucleotide that encodes the polypeptide comprising SEQ ID NO: 2. In another preferred embodiment, the invention provides an isolated polynucleotide that encodes a polypeptide comprising an amino acid sequence that has at least 98% identity to SEQ ID NO: 2. Also, in other preferred embodiments, the invention provides an isolated polynucleotide comprising a nucleotide sequence that is complementary to the polynucleotide encoding the
30 polypeptide comprising SEQ ID NO: 2 or an isolated polynucleotide comprising a nucleotide sequence that is complementary to the polynucleotide encoding the polypeptide comprising an amino acid sequence that has at least 98% identity to SEQ ID NO: 2.

In another preferred embodiment, the present invention provides an isolated ABC1

polynucleotide comprising SEQ ID NO: 1. In a further preferred embodiment, the present invention provides an isolated polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1. In yet another preferred embodiment, the invention provides a polynucleotide comprising a nucleotide sequence that has at least 90% identity with SEQ ID NO: 1. More preferably, the polynucleotide comprises a nucleotide sequence that has at least 95%, 96%, 97%, 98%, or 99% identity to SEQ ID NO: 1. Also, in other preferred embodiments, the present invention provides an isolated polynucleotide comprising a nucleotide sequence that is complementary to the polynucleotide comprising SEQ ID NO: 1, an isolated polynucleotide comprising a nucleotide sequence that is complementary to the polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, and an isolated polynucleotide that is complementary to a polynucleotide comprising a nucleotide sequence that has at least 90% identity with SEQ ID NO: 1.

The present invention also provides ABC1 polynucleotides corresponding to the 5' flanking region of the ABC1 gene. In one preferred embodiment, the invention provides an isolated polynucleotide comprising SEQ ID NO: 3. In other preferred embodiments, the invention provides an isolated polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3. Preferably, the isolated polynucleotide comprises nucleotides 1394-1532 of SEQ ID NO: 3. In another preferred embodiment, the invention provides an isolated polynucleotide that hybridizes under stringent conditions to a polynucleotide comprising SEQ ID NO: 3. Also, in other preferred embodiments, the present invention provides an isolated polynucleotide that hybridizes under stringent conditions to a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3. In yet another preferred embodiment of the present invention, an isolated polynucleotide that has at least 80% identity to a polynucleotide comprising SEQ ID NO: 3 is provided. More preferably, the polynucleotide has at least 90% identity to a polynucleotide comprising SEQ ID NO: 3. Even more preferably, the polynucleotide has at least 95% identity to a polynucleotide comprising SEQ ID NO: 3. Also provided in preferred embodiments is an isolated polynucleotide that has at least 80% identity to a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3. More preferably, the polynucleotide has at least 90% identity, and even more preferably at least 95% identity, to a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3. In addition, the present invention provides an isolated polynucleotide comprising a nucleotide sequence that is complementary to the above described 5' flanking ABC1 polynucleotides. In one preferred embodiment, the invention provides an isolated

polynucleotide comprising a nucleotide sequence that is complementary to a polynucleotide comprising SEQ ID NO: 3. In another preferred embodiment, the present invention provides an isolated polynucleotide comprising a nucleotide sequence that is complementary to a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3.

The present invention also provides ABC1 polynucleotides corresponding to the 3' flanking region of the ABC1 gene. In preferred embodiments, the invention provides an isolated polynucleotide comprising SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6, and the complementary sequences thereof. In other preferred embodiments, the invention provides an isolated polynucleotide that hybridizes under stringent conditions to a polynucleotide comprising SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6, and the complementary sequences thereof. In still other preferred embodiments, the invention provides an isolated polynucleotide that has at least 80% identity to a polynucleotide comprising SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6, and the complementary sequence thereof. More preferably, the polynucleotide has at least 90% identity to a polynucleotide comprising SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6. Even more preferably, the polynucleotide has at least 95% identity to a polynucleotide comprising SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6.

In addition, the present invention also provides ABC1 polynucleotides from Tangier disease patients. In one preferred embodiment, the present invention provides an isolated polynucleotide encoding the polypeptide comprising SEQ ID NO: 8. In another preferred embodiment, the present invention provides an isolated polynucleotide comprising SEQ ID NO: 7. In yet another embodiment, the present invention provides an isolated polynucleotide encoding the polypeptide comprising SEQ ID NO: 10. In still another preferred embodiment, the present invention provides an isolated polynucleotide comprising SEQ ID NO: 9. The present invention further provides an isolated polynucleotide comprising a nucleotide sequence that is complementary to the described polynucleotides.

In another aspect, the present invention provides a composition comprising any of the above described polynucleotides and a suitable carrier. In one preferred embodiment, the present invention provides a composition comprising an isolated polynucleotide encoding a polypeptide comprising SEQ ID NO: 2, a polynucleotide comprising SEQ ID NO: 1, a polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, or a polynucleotide encoding the polypeptide comprising an amino acid sequence that has at least 98% identity to SEQ ID NO: 2, and a suitable carrier. In another preferred embodiment, the composition

comprises an isolated polynucleotide comprising a nucleotide sequence that has at least 90% identity with the polynucleotide comprising SEQ ID NO: 1 and a suitable carrier. In other preferred embodiments, the composition comprises an isolated polynucleotide comprising SEQ ID NO: 3 or an isolated polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3 and a suitable carrier. In still other preferred embodiments, the invention provides a composition comprising a polynucleotide that hybridizes under stringent conditions to the polynucleotide of SEQ ID NO: 3, or a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3, as well as a composition comprising a polynucleotide that has at least 80% identity to the polynucleotide comprising SEQ ID NO: 3, or a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3 and a suitable carrier. Also provided by the present invention is a composition comprising an isolated polynucleotide comprising a nucleotide sequence that is complementary to any of the described polynucleotides and a suitable carrier.

In addition, the present invention provides recombinant vectors and host cells comprising any of the described ABC1 polynucleotide sequences. In one preferred embodiment, the present invention provides a recombinant vector comprising an isolated polynucleotide encoding a polypeptide comprising SEQ ID NO: 2, an isolated polynucleotide comprising SEQ ID NO: 1, an isolated polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, or an isolated polynucleotide encoding the polypeptide comprising an amino acid sequence that has at least 98% identity to SEQ ID NO: 2. In another preferred embodiment, the recombinant vector comprises an isolated polynucleotide comprising a nucleotide sequence that has at least 90% identity, and more preferably at least 95% identity, with a polynucleotide comprising SEQ ID NO: 1. In still another preferred embodiment, the recombinant vector comprises an isolated polynucleotide comprising SEQ ID NO: 7 or SEQ ID NO: 9. The present invention further provides a recombinant vector comprising an isolated polynucleotide comprising a nucleotide sequence that is complementary to any of the described polynucleotides. In yet another preferred embodiment, the recombinant vector comprises any of the described polynucleotides and further comprises a heterologous promoter polynucleotide. One suitable heterologous promoter is a cytomegalovirus promoter. In a particularly preferred embodiment, the recombinant vector is pCEPhABC1.

The present invention also provides a recombinant vector comprising an isolated polynucleotide comprising an ABC1 5' flanking sequence. In one preferred embodiment, the

invention provides a recombinant vector comprising an isolated polynucleotide comprising SEQ ID NO: 3 or an isolated polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3. In still other preferred embodiments, the invention provides a recombinant vector comprising a polynucleotide that hybridizes under stringent
5 conditions to the polynucleotide of SEQ ID NO: 3, or a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1532 of SEQ ID NO: 3, as well as a recombinant vector comprising a polynucleotide that has at least 80% identity to these polynucleotides. The present invention further provides a recombinant vector comprising an isolated polynucleotide comprising a nucleotide sequence that is complementary to any of the
10 described polynucleotides. In yet another preferred embodiment, the recombinant vector comprises any of the described polynucleotides and further comprises at least one polynucleotide encoding a heterologous polypeptide. Suitable heterologous polypeptides include luciferase, β -galactosidase, chloramphenicol acetyl transferase, acetyl transferase, and green fluorescent proteins. Preferably, the heterologous polypeptide is a luciferase protein. In a
15 particularly preferred embodiment, the recombinant vector is pAPR1.

In addition, the present invention provides host cells comprising any of the described recombinant vectors. The present invention further provides compositions comprising any of the described recombinant vectors and a suitable carrier.

The present invention also provides methods for producing the ABC1 protein in a
20 mammalian host cell as well as methods for expressing the ABC1 protein in a mammalian subject. The method for producing an ABC1 protein in a mammalian host cell comprises the steps of: (a) transfecting the mammalian host cell with a recombinant expression vector comprising a polynucleotide encoding ABC1 in an amount sufficient to produce a detectable level of ABC1 protein, and (b) purifying the produced ABC1 protein. The method for
25 expressing ABC1 protein in a mammalian subject comprises the step of administering to a mammalian subject a recombinant expression vector comprising a polynucleotide encoding ABC1 in an amount sufficient to express ABC1 protein in the mammalian subject.

In addition, the present invention provides compositions and methods suitable for increasing cholesterol efflux from cells of a mammalian subject. In one preferred embodiment,
30 the method comprises administering to the mammalian subject a recombinant expression vector comprising a polynucleotide encoding ABC1 in an amount sufficient to increase cholesterol efflux from the cells. Suitable recombinant expression vectors include vectors comprising an isolated polynucleotide encoding a polypeptide comprising SEQ ID NO: 2, an isolated

polynucleotide comprising SEQ ID NO: 1, an isolated polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, and an isolated polynucleotide encoding the polypeptide comprising an amino acid sequence that has at least 98% identity to SEQ ID NO: 2. Preferred expression vectors include viral vectors, especially adenoviral vectors and lentiviral vectors. In other
5 embodiments, the invention provides non-viral delivery systems, including DNA-ligand complexes, adenovirus-ligand-DNA complexes, direct injection of DNA, CaPO₄ precipitation, gene gun techniques, electroporation, liposomes and lipofection.

In another preferred embodiment, the method for increasing cholesterol efflux from cells of a mammalian subject comprises administering to the mammalian subject a therapeutic amount
10 of a compound that increases the expression of ABC1 in the cells. One suitable method comprises administering to the mammalian subject a cAMP analogue. Suitable cAMP analogues include 8-bromo cAMP, N6-benzoyl cAMP, and 8-thiomethyl cAMP. Another suitable method comprises administering to the mammalian subject a compound that increases the synthesis of cAMP, e.g. forskolin. Yet another suitable method comprises administering to the mammalian
15 subject a compound that inhibits the degradation of cAMP, such as a phosphodiesterase inhibitor. Suitable phosphodiesterase inhibitors include rolipram, theophylline, 3-isobutyl-1-methylxanthine, R020-1724, vinpocetine, zaprinast, dipyridamole, milrinone, amrinone, pimobendan, cilostamide, enoximone, peroximone, and vesnarinone.

In addition, another suitable method for increasing cholesterol efflux from cells of a
20 mammalian subject comprises administering to the mammalian subject a least one ligand for a nuclear receptor in an amount sufficient to increase cholesterol efflux. Suitable ligands include LXR, RXR, FXR, SXR and PPAR ligands. In one preferred embodiment, the method comprises administering to a mammalian subject a ligand for an LXR nuclear receptor. Suitable LXR ligands include 20(S) hydroxycholesterol, 22(R) hydroxycholesterol, 24(S) hydroxycholesterol,
25 25-hydroxycholesterol, and 24(S), 25 epoxycholesterol. Preferably, the LXR ligand is 20(S) hydroxycholesterol. In another preferred embodiment, the method comprises administering to a mammalian subject a ligand for an RXR nuclear receptor. Suitable RXR ligands include 9-*cis* retinoic acid, retinol, retinal, all-trans retinoic acid, 13-*cis* retinoic acid, acitretin, fenretinide, etretinate, CD 495, CD564, TTNN, TTNNPB, TTAB, and LGD 1069. Preferably, the RXR
30 ligand is 9-*cis* retinoic acid. In another preferred embodiment, the method comprises administering to a mammalian subject a ligand for a PPAR nuclear receptor. One suitable ligand is a ligand selected from the class of thiazolidinediones. In yet another preferred embodiment,

the method comprises administering at least two ligands for a nuclear receptor. In a particularly preferred embodiment, the ligands are 20(S) hydroxycholesterol and 9-*cis* retinoic acid.

In addition, another suitable method for increasing cholesterol efflux from cells of a mammalian subject comprises administering to the mammalian subject an eicosanoid in an amount sufficient to increase cholesterol efflux. Suitable eicosanoids include prostaglandin E2, prostaglandin J2, and prostacyclin (prostaglandin I2).

In another embodiment, the present invention provides a method for increasing cholesterol efflux from cells of a mammalian subject comprising administering to the mammalian subject a compound that increases ABC1 activity in an amount sufficient to increase cholesterol efflux from the cells.

The present invention also provides methods suitable for increasing the gene expression of ABC1 in a mammalian subject. In one preferred embodiment, the method comprises administering to the mammalian subject at least one ligand for a nuclear receptor in an amount sufficient to increase the gene expression of ABC1. Suitable ligands include ligands for LXR, RXR, FXR, SXR, and PPAR nuclear receptors. In another preferred embodiment, the method comprises administering to the mammalian subject a cAMP analogue in an amount sufficient to increase the gene expression of ABC1. In yet another preferred embodiment, the method comprises administering to the mammalian subject a compound that increases the synthesis of cAMP in an amount sufficient to increase the gene expression of ABC1.

In addition, the present invention provides a method for screening a test compound for ABC1 expression modulating activity comprising the steps of: (a) operatively linking a reporter cDNA with an expression modulating portion of the mammalian ABC1 gene to produce a recombinant reporter construct; (b) transfecting the recombinant reporter construct into a population of host cells; (c) assaying the level of reporter gene expression in a sample of the host cells; (d) contacting the host cells with the test compound being screened; (e) assaying the level of reporter gene expression in a sample of the host cells after contact with the test compound; and (f) comparing the relative change in the level of reporter gene expression caused by exposure to the test compound, thereby determining the ABC1 expression modulating activity. The recombinant reporter construct comprises a reporter gene operatively linked to an expression modulating portion of the mammalian ABC1 gene, such as any of the ABC1 5' flanking region sequences provided by the present invention. In one preferred embodiment, the expression modulating portion of the ABC1 gene comprises SEQ ID NO: 3. In another preferred embodiment, the expression modulating portion of the ABC1 gene comprises nucleotides 1-

1532, 1080-1643, 1181-1643, 1292-1643, 1394-1643, or 1394-1532 of SEQ ID NO: 3. Suitable reporter cDNAs include luciferase, β -galactosidase, chloramphenicol, and green fluorescent protein cDNA. Preferably, the host cell is a mammalian cell. In a particularly preferred embodiment of the method, the recombinant reporter construct is pAPR1.

5 Also provided by the present invention is a method for screening a test compound to determine whether the test compound promotes ABC1-mediated cholesterol efflux from cells in culture comprising the steps of: (a) assaying the level of cholesterol efflux in a sample of mammalian cells maintained in culture to determine a control level of cholesterol efflux; (b) contacting the cells with the test compound being screened; (c) assaying the level of cholesterol
10 efflux in a sample of cells after contact with the test compound; (d) assaying the level of ABC1-mediated cholesterol efflux in a sample of cells after contact with the test compound, thereby determining whether the test compound promotes ABC1-mediated cholesterol efflux from cells in culture. The cells can be derived from primary cultures or a cell line. Suitable cells for screening the test compound include fibroblast, macrophage, hepatic, and intestinal cell lines.
15 Preferably, the cell line is RAW 264.7. In one preferred embodiment, the ABC1-mediated cholesterol efflux is measured using an anti-ABC1 antibody that inhibits the activity of ABC1 upon binding. In another preferred embodiment, the ABC1-mediated cholesterol efflux is measured using an antisense ABC1 polynucleotide. In a particularly preferred embodiment, the antisense polynucleotide comprises SEQ ID NO: 57.

20 In addition, the present invention provides methods for detecting the comparative level of ABC1 expression in cells of a mammalian subject. Such methods can be used to determine the susceptibility of a subject to coronary heart disease. A method for detecting the comparative level of ABC1 expression in cells of a mammalian subject is provided which comprises (a) obtaining a cell sample from the mammalian subject, (b) assaying the level of ABC1 mRNA
25 expression in the cell sample; and (c) comparing the level of ABC1 mRNA expression in the cell sample with a pre-determined standard level of ABC1 mRNA expression, thereby detecting the comparative level of ABC1 gene expression in the cells of a mammalian subject. Suitable methods for measuring the level of ABC1 mRNA expression include, for example, RT-PCR, northern blot, and RNase protection assay.

30 The present invention also provides methods for detecting the comparative level of ABC1 protein in cells of a mammalian subject. Such methods can be used to determine the susceptibility of a subject to coronary heart disease. A method for detecting the comparative amount of ABC1 protein in the cells of a mammalian subject is provided which comprises (a)

obtaining a cell sample from the mammalian subject, (b) assaying the amount of ABC1 protein in the cell sample, and (c) comparing the amount of ABC1 protein in the cell sample with a pre-determined standard amount of ABC1 protein, thereby detecting the comparative level of ABC1 protein in the cells of the mammalian subject. The amount of ABC1 protein can be determined
5 using various immunoassays available in the art. For example, the amount of ABC1 protein can be determined by (a) contacting the cell sample with a population of anti-ABC1 antibodies and (b) detecting the specific-binding ABC1 antibodies associated with the sample. Suitable methods for detecting ABC1 antibodies include western blotting, immunoprecipitation, and FACS.

10 In another aspect, the present invention provides antibodies that bind specifically to the described ABC1 polypeptides. In one preferred embodiment, the present invention provides an isolated antibody that binds specifically to an isolated polypeptide comprising SEQ ID NO: 2. In another preferred embodiment, the invention provides an isolated antibody that bind specifically to an isolated polypeptide comprising an amino acid sequence that has at least 98% identity with
15 SEQ ID NO: 2. The antibody can be a monoclonal antibody or the antibody can be a polyclonal antibody. In yet another embodiment, the antibody, upon binding to an ABC1 polypeptide, inhibits the cholesterol transport activity of the ABC1 polypeptide.

In addition, the present invention provides kits suitable for screening a compound to determine the ABC1 expression modulating activity of the compound comprising a reporter
20 cDNA operatively linked to an expression modulating portion of the mammalian ABC1 gene in an amount sufficient for at least one assay and instructions for use. In one preferred embodiment, the kit further comprises means for detecting the reporter gene. In another preferred embodiment, the expression modulating portion of the mammalian ABC1 gene comprises SEQ ID NO: 3. In yet another preferred embodiment, the expression modulating
25 portion of the mammalian ABC1 gene comprises nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, 1394-1643, or 1394-1532 of SEQ ID NO: 3. Suitable reporter cDNAs include luciferase, β -galactosidase, chloramphenicol, and green fluorescent protein cDNA. Preferably, the reporter cDNA is luciferase. In a particularly preferred embodiment of the method, the recombinant reporter construct is pAPR1.

30 The present invention also provides kits suitable for screening a compound to determine whether the compound modulates ABC1-dependent cholesterol efflux. In one preferred embodiment, the kit comprises an inactivating anti-ABC1 antibody in an amount sufficient for at least one assay and instructions for use. In another preferred embodiment, the kit comprises an

antisense ABC1 oligonucleotide in an amount sufficient for at least one assay and instructions for use. In a particularly preferred embodiment, the antisense ABC1 oligonucleotide comprises SEQ ID NO: 53.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1A-D is a graphical representation showing the results of control cholesterol efflux and cholesterol efflux in the presence of HDL and apo A-I from normal fibroblast cells (1A, C) and fibroblast cells from Tangier disease patients (1B, D). The open circles represent the cholesterol efflux from cells that were not exposed to HDL or apo A-I, the closed circles represent the cholesterol efflux from cells exposed to HDL, and the closed diamonds represent the cholesterol efflux from cells exposed to apo A-I;

Figure 2 is a graphical representation of a gene expression microarray analysis showing a comparison of the gene expression found in cells from a Tangier patient (TD1) and that found in normal cells, whereby a total of 58,800 human cDNAs were hybridized with cDNA prepared from mRNA of cAMP-treated TD1 cells cDNA (labeled with Cy3 dye) and with cDNA prepared from mRNA of cAMP-treated normal cells (labeled with Cy5 dye);

Figure 3 is a schematic diagram showing a restriction map of the recombinant expression vector pCEPhABC1, which contains the open reading frame of the human ABC1 gene;

Figure 4 is a schematic diagram of the gene structure of human ABC1, showing a comparison between the published human ABC1 amino acid sequence (GenBank, Accession # AJ012376) and the presently disclosed and claimed human ABC1 amino acid sequence ("CVT") which has sixty additional amino acids at the N-terminal end;

Figure 5 is a graphical representation showing the inhibitory effect that ABC1 transport inhibitors 4,4-diisothiocyanostilbene-2,2'-disulfonic acid (DIDS) and sulphobromophthaleine (BSP) have on apo A-I-mediated cholesterol efflux, wherein the open circles indicate the apo A-I-mediated cholesterol in the presence of BSP and the closed circles indicate the apo A-I-mediated cholesterol in the presence of DIDS;

Figure 6 is a graphical representation showing the inhibitory effect of an antisense ABC1 oligonucleotide on apo A-I-mediated cholesterol efflux, showing the apo A-I-mediated cholesterol efflux in cells incubated without antisense oligonucleotide, the apo A-I-mediated cholesterol efflux in cells exposed to 30 μ M β -globin antisense oligonucleotide, and the apo A-I-mediated cholesterol efflux in cells exposed to 30 μ M ABC1 antisense oligonucleotide;

Figure 7 is a graphical representation demonstrating the stimulation of apo A-I-mediated

cholesterol efflux caused by overexpression of the ABC1 gene using RAW 264.7 mouse macrophage cells stably transfected with an expression plasmid for human ABC1 (pCEPhABC1), showing the apo A-I-mediated cholesterol efflux in control parental cells (no pCEPhABC1) and the apo A-I-mediated cholesterol efflux in clonal cells transfected with pCEPhABC1 (L3, L5, L6);

Figure 8 is a graphical representation of reverse transcription polymerase chain reaction (RT-PCR) analyses showing the level of ABC1 gene expression in normal cells and cells from Tangier's disease patients (TD1 and TD2) that have been either exposed to albumin (closed bars), exposed to 8-Br-cAMP (open bars), cholesterol-loaded (shaded bars), or cholesterol-loaded and subsequently exposed to apo A-I (hatched bars);

Figure 9 is a graphical representation of the results of RT-PCR analyses showing the level of ABC1 gene expression in RAW 264.7 cells exposed to either ethanol (0.1% v/v), 9-*cis* retinoic acid (9-*cis* RA; 10 μ M), 20(S) hydroxycholesterol (20(S)-OH; 10 μ M), or 9-*cis* RA and 20(S)-OH (10 μ M each);

Figure 10 shows the results of immunoprecipitation analyses indicating the level of cell-surface ABC1 protein found in normal fibroblasts (NL1, 10A) and fibroblasts from a Tangier's disease patient (TD1, 10B) in the presence of either no additives (control), 8-Br-cAMP (1mM), cholesterol (30 μ g/ml), or cholesterol and 8-Br-cAMP (30 μ g/ml and 1mM, respectively);

Figure 11 is a schematic diagram showing a restriction map of the recombinant expression vector pAPR1, which contains the 5' flanking region of the ABC1 gene positioned upstream of the open reading frame of the luciferase reporter gene;

Figure 12 is a graphical representation showing the level of luciferase reporter gene expression induced in RAW 264.7 cells transfected with pAPR1 in the presence of either EtOH (control), 20(S)-OH (10 μ M), 9-*cis* RA (10 μ M), or both 20(S)-OH and 9-*cis* RA (10 μ M each);

Figure 13 is a schematic diagram of the 5' flanking region of the ABC1 gene.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In accordance with the present invention, novel polypeptides that increase the cholesterol efflux from cells are provided. In particular, the present invention provides novel ATP-Binding Cassette 1 (ABC1) polypeptides that have been shown to increase cholesterol efflux.

ABC1 is a member of the family of ATP-binding cassette proteins that reside in cell membranes and utilize ATP hydrolysis to transport a wide variety of substrates across the plasma membrane. It should be noted that the terms "ABC1" and "ABCA1" both refer to the same ATP-

binding cassette protein. The term "ABCA1" was introduced in 1999 by a nomenclature committee and has received limited acceptance in the field. To date, more than 30 members of this family have been identified in the human genome. These homologous proteins contain channel-like structures through which molecules are transported through the cell membrane and one or two domains which bind to ATP to couple energy generating ATP-hydrolysis to transport. Family members include the multidrug resistance factors (MDR/P-glycoproteins; Chen et al., *Cell*, 47: 381-389 (1986); Stride et al., *Mol. Pharmacol.*, 49:962-971 (1996)), transporters associated with antigen presentation (Neefjes et al., *Science*, 261:769-771 (1993); Shepherd et al., *Cell*, 74:577-584 (1993)), and the cystic fibrosis transmembrane conductance regulator (Chang et al., *J. Biol. Chem.*, 269:18572-18575 (1994); Rommens et al., *Science*, 245: 1059-1065 (1989)). Members of the ABC transporter family are generally composed of 4 domains found within two symmetric halves that are linked by a long charged region and a highly hydrophobic segment. Each half contains a hydrophobic domain, containing 6 transmembrane segments and a hydrophilic nucleotide binding domain containing highly conserved Walker A and B sequence motifs typical of many ATPases (Hyde et al., *Nature*, 346:362-365 (1990); Luciani et al., *Genomics*, 21: 150-159 (1994)). The transporter activity is dependent on the interaction with ATP at the nucleotide binding domains and by regulation via phosphorylation of residues in the region linking the two symmetric halves (Becq et al., *J. Biol. Chem.*, 272: 2695-2699 (1997)).

Several lines of evidence described herein identified ABC1 as a pivotal protein in the apolipoprotein-mediated mobilization of intracellular cholesterol stores. First, the studies presented herein showed that ABC1 is defective in Tangier disease, a genetic disorder characterized by abnormal HDL-cholesterol metabolism. As shown previously, and herein at Example 1, the genetic defect in Tangier disease causes a defect in the pathway of apolipoprotein mediated efflux of cholesterol from within cells, resulting in significantly decreased cholesterol efflux activity and low HDL-cholesterol levels (Oram et al., *J. Lipid Res.*, 37:2743-2491 (1996); Francis et al., *J. Clin. Invest.*, 96: 78-87 (1995)). Genetic linkage analysis of families with Tangier disease assigned the defective gene to an interval on chromosome 9q31 (Rust et al., *Nature Genetics*, 20: 96-98 (1998)). A search of public databases revealed that the ABC1 gene was localized to chromosome 9q22-9q31, which is broader than, but includes the interval revealed in Rust et al. (Luciani et al., *supra* (1994)). Based on that data, radiation hybrid mapping of the human ABC1 gene was performed, which placed the gene between two markers squarely within the 7-cM region of human chromosome 9q31 reported by Rust et al. In addition,

as shown in Example 2, microarray analysis revealed that the ABC1 gene is 2.5-fold underexpressed in Tangier patient cells as compared with normal cells. These studies identified the defective gene in Tangier disease as ABC1. In addition, further studies presented herein linked ABC1 activity to cholesterol efflux activity. First, studies showed that inhibitors of ABC1 transport activity, such as 4,4-diisothiocyanostilbene-2,2'-disulfonic acid (DIDS) and sulphobromophthaleine (BSP), also inhibited apoAI-mediated cholesterol efflux from fibroblast cells (see Example 6). Also, inhibition of ABC1 gene expression, using an antisense ABC1 oligonucleotide, was shown to inhibit apoAI-mediated cholesterol efflux from fibroblast cells (Example 7). In contrast, transfection studies, in which the ABC1 gene was transfected into mouse monocyte cells, showed that overexpression of ABC1 results in an increase in apoAI-mediated efflux (Example 8). Finally, RT-PCR performed using wildtype and Tangier patient mRNA revealed that ABC1 mRNA expression is regulated by cellular conditions related to cholesterol efflux in normal skin fibroblast cells, but not in Tangier patient fibroblasts (Example 9). Based on these findings, it was determined that ABC1 plays a major role in cholesterol efflux.

It is postulated that ABC1 plays a role in the translocation of intracellular cholesterol to the outer leaflet of the plasma membrane. Deficient transport of intracellular cholesterol due to a lack of ABC1 or defective ABC1 results in a lack of cholesterol in specific membrane domains with which apoAI and other apolipoproteins specifically interact (Stangl et al., *J. Biol. Chem.*, 273: 31002-31008 (1998); Babitt et al., *J. Biol. Chem.*, 272: 13242-13249 (1997))). The failed delivery of cholesterol to apoAI leads to the formation of cholesterol-deficient HDL particles that are rapidly removed from the plasma. (Bojanovski et al., *J. Clin. Invest.*, 80: 1742-1747 (1987)).

Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide.

As used herein, the term "polynucleotide(s)" is defined to encompass DNA and RNA of

both synthetic and natural origin. The polynucleotide may exist as single- or double-stranded DNA or RNA, or an RNA/DNA heteroduplex. Thus, the polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, or single, double-, and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or triple-stranded, or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms of polynucleotides. The term "polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

The term "polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide" refers to both short amino acid sequences, commonly referred to as peptides, as well as longer amino acid sequences, generally referred to as proteins. The polypeptide may contain amino acids other than the 20 gene encoded amino acids. Moreover, the polypeptide may be modified either by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques, which are well-known in the art. A given polypeptide may contain many types of modifications. Also, the same type of modification may be present in the same or varying degree at one or more sites in the polypeptide. Modifications may occur anywhere in the polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, formylation, gamma-carboxylation, glycosylation, hydroxylation, iodination, methylation, myristoylation, oxidation, phosphorylation, prenylation, sulfation, and selenoylation, as well as the covalent attachment of a nucleotide or nucleotide derivative, a lipid or lipid derivative, or a phosphatidylinositol. Other modifications include cross-linking, cyclization, formation of pyroglutamate, GPI anchor formation, proteolytic processing, racemization, and t-RNA-mediated addition of amino acids, such as arginylation and ubiquitination. See, for example, *Proteins- Structure and Molecular Properties*, 2nd Ed., T.E.

Creighton, W.H. Freedman and Co., New York (1993); Wold, F., Posttranslational Protein Modification: Perspectives and Prospects, in Posttranslational Covalent Modification of Proteins, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., Meth. Enzymol., 182: 626-646 (1990); and Rattan et al., Protein Synthesis: Posttranslational Modifications and Aging, Ann. N.Y. Acad. Sci., 663: 48-62 (1992)). The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

10 A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to SEQ ID NO:3 or nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, 1394-1643, or 1394-1532 of SEQ ID NO: 3, or the complements thereof. A polynucleotide of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to SEQ ID
15 NO:4, SEQ ID NO: 5, or SEQ ID NO: 6 or the complements thereof.

"Stringent hybridization conditions" refers to an overnight incubation at 42° C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about
20 65°C.

As used herein, the term "complementary" refers to the hybridization or base pairing between nucleotides, such as, for example, between the two strands of a double-stranded polynucleotide or between an oligonucleotide primer and a primer binding site on a single-stranded polynucleotide to be amplified or sequenced. Two single-stranded nucleotide
25 molecules are said to be complementary when the nucleotides of one strand, optimally aligned with appropriate nucleotide insertions, deletions or substitutions, pair with at least about 80% of the nucleotides of the other strand.

"Identity", as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences.
30 "Identity" or "similarity" also has an art-recognized meaning that refers to the degree of sequence relatedness between polypeptide or polynucleotide sequences, as determined by the match between strings of such sequences. "Identity" and "similarity" can be calculated using a number of well known methods, including those published in Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York, (1988); Biocomputing: Informatics and

Genome Projects, Smith, D.W., ed., Academic Press, New York, (1993); Computer Analysis of Sequence Data, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, (1994); Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, (1987); and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, (1991); and
5 Carillo, H., and Lipton, D., SIAM J Applied Math 48:1073 (1988)). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in "Guide to Huge Computers," Martin J. Bishop, ed., Academic Press, San Diego, (1994), and Carillo, H., and Lipton, D., SIAM J Applied Math 48:1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences
10 tested. Methods for aligning polynucleotides or polypeptides are codified in computer programs, including the GCG program package (Devereux, J., et al., Nucleic Acids Research (1984) 12(1):387 (1984)), BLASTP, BLASTN, FASTA (Atschul, S.F. et al., J. Molec. Biol. 215:403 (1990), Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711 (using the
15 local homology algorithm of Smith and Waterman, Advances in Applied Mathematics 2:482-489 (1981))).

When using any of the sequence alignment programs to determine whether a particular sequence is, for instance, 90% identical to a reference sequence, the parameters are set so that the percentage of identity is calculated over the full length of the reference polypeptide or
20 polynucleotide and that gaps in identity of up to 10% of the total number of nucleotides in the reference polynucleotide are allowed.

A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of
25 Brutlag et al. (Comp. App. Biosci. 6:237-245 (1990)). The term "sequence" includes nucleotide and amino acid sequences. In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB search of a DNA sequence to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1,
30 Joining Penalty=30, Randomization Group Length=0, and Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, and Window Size=500 or query sequence length in nucleotide bases, whichever is shorter. Preferred parameters employed to calculate percent identity and similarity of an amino acid alignment are: Matrix=PAM 150, k-tuple=2, Mismatch Penalty= 1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size

Penalty=0.05, and Window Size=500 or query sequence length in amino acid residues, whichever is shorter.

As an illustration, a polynucleotide having a nucleotide sequence of at least 90% "identity" to a sequence contained in SEQ ID NO: 1 means that the nucleotide sequence of the polynucleotide is identical to a sequence contained in SEQ ID NO: 1 except that the polynucleotide sequence may include up to ten point mutations per each 100 nucleotides of the total length of SEQ ID NO: 1. In other words, to obtain a polynucleotide comprising a nucleotide sequence that has at least 90% identity to SEQ ID NO: 1, up to 10% of the nucleotides in the sequence contained in SEQ ID NO: 1 can be deleted, inserted, or substituted with other nucleotides. These changes may occur anywhere throughout the polynucleotide, and may be interspersed either individually among nucleotides or in one or more contiguous groups within SEQ ID NO: 1.

Similarly, a polypeptide having an amino acid sequence of at least 98% "identity" to a sequence contained in SEQ ID NO: 2 means that the amino acid sequence of the polypeptide is identical to a sequence contained in SEQ ID NO: 2 except that the polypeptide sequence may include up to 2 amino acid alterations per each 100 amino acids of the total length of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 98% identical to SEQ ID NO: 2, up to 2% of the amino acid residues in the sequence contained in SEQ ID NO: 2 can be deleted, inserted, or substituted with other amino acid residues. These changes may occur anywhere throughout the polypeptide, and may be interspersed either individually among residues or in one or more contiguous groups within SEQ ID NO: 2.

"A polypeptide having biological activity" refers to a polypeptide exhibiting activity similar, but not necessarily identical, to an activity of a polypeptide of the present invention (e.g. cholesterol transport activity), as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.).

"Polypeptide variant" refers to a polypeptide differing from the ABC1 polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polypeptide comprising SEQ ID NO: 2.

Preferably, the polypeptide variant retains its biological activity, i.e., cholesterol transport activity. Variants include, but are not limited to, splice variants and allelic variants, as well as addition, deletion, and substitution variants.

Likewise, "polynucleotide variant" refers to a polynucleotide differing from the
5 polynucleotide of the present invention, but retaining essential properties thereof. The variants may contain alterations in the coding regions, non-coding regions, or both. Thus, for example, an ABC1 polynucleotide variant has a nucleotide sequence that differs from that of SEQ ID NO: 1, but encodes a polypeptide that has cholesterol transport activity. Also, for example, a polynucleotide variant has a nucleotide sequence that differs from that of SEQ ID NO: 3, but
10 retains promoter activity. Especially preferred are polynucleotide variants containing alterations that produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 10-20, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred.
15 Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (e.g. changing codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

"Allelic variants" are naturally-occurring variants that refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B.,
20 ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

The term "conservative amino acid substitution" refers to a substitution of a native amino acid residue with a nonnative residue such that there is little or no effect on the polarity or charge
25 of the amino acid residue at that position. For example, a conservative substitution results from the replacement of a non-polar residue in a polypeptide with any other non-polar residue. Another example of a conservative substitution is the replacement of an acidic residue with another acidic residue. Conservative substitutions are expected to produce ABC1 polypeptides having functional and chemical characteristics similar to those of the naturally-occurring ABC1
30 polypeptide.

The term "ortholog" refers to a polypeptide that corresponds to a polypeptide identified from a different species. For example, mouse and human ABC1 polypeptides are considered orthologs.

The term "vector" is used to refer to any molecule (e.g. nucleic acid, plasmid, or virus) used to transfer coding information to a host cell.

The term "expression vector" refers to a vector that is suitable for transformation of a host cell and contains nucleic acid sequences that direct and/or control the expression of inserted
5 heterologous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

As used herein, the term "transcriptional regulatory region" or "expression modulating portion" refers to any region of the gene, including, but not limited to, promoters, enhancers, and repressors.

10 As used herein, the term "promoter" refers to an untranscribed sequence located upstream (*i.e.*, 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that controls the transcription of the structural gene.

As used herein the term "enhancers" refers to *cis*-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively
15 orientation and position independent. They have been found 5' and 3' to the transcription unit.

"Host cell" is a cell that has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence. The term includes the progeny of the parent cell, whether or not the progeny is identical in morphology or in genetic make-up to the original parent, so long as the selected gene is present.

20 The term "operatively linked" is used herein to refer to an arrangement of flanking sequences wherein the flanking sequences so described are configured or assembled so as to perform their usual function. Thus, a flanking sequence operably linked to a coding sequence may be capable of effecting the replication, transcription and/or translation of the coding sequence. For example, a coding sequence is operably linked to a promoter when the promoter is
25 capable of directing transcription of that coding sequence. A flanking sequence need not be contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

30 The term "transfection" is used to refer to the uptake of foreign or exogenous DNA by a cell, and a cell has been "transfected" when the exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are well known in the art and are disclosed herein. See, *e.g.*, Graham *et al.*, 1973, *Virology* 52:456; Sambrook *et al.*, *Molecular Cloning*, A

Laboratory Manual (Cold Spring Harbor Laboratories, 1989); Davis *et al.*, *Basic Methods in Molecular Biology* (Elsevier, 1986); and Chu *et al.*, 1981, *Gene* 13:197. Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

ABC1 Polypeptides

5 The present invention relates to novel human ABC1 polypeptides. In one embodiment, the ABC1 polypeptide comprises the amino acid sequence shown in SEQ ID NO: 2. In contrast to the human ABC1 protein reported by others, the ABC1 polypeptide shown in SEQ ID NO: 2 has an additional 60 amino acids at the amino terminus, revealing an ABC1 protein of 2261
10 amino acids rather than 2201 amino acids (see Langmann *et al.* in *Biochem. Biophys. Res. Comm.*, 257, 29-33 (1999)). In addition, the ABC1 polypeptide shown in SEQ ID NO: 2 differs from other reported sequences at several amino acid residues. Specifically, the present ABC1 polypeptide shown in SEQ ID NO: 2 has a K for R at residue 159, I for V at 765, M for I at 823, I for T at 1495, L for P at 1588, K for R at 1914, and L for P at 2108. To remain consistent with
15 published notation, the above amino acid numbers are those of Lawn *et al.*, *J. Clin. Invest.*, 104: R25-31 (1999), rather than those of SEQ ID NO: 2. As discussed in further detail below, the sequence difference likely arises from the fact that the first ABC1 cDNA was cloned from mouse using a PCR-based strategy and the subsequently reported human ABC1 cDNA sequences were predicted from the sequence of the mouse protein. The ABC1 protein has an approximate
20 molecular weight of 240 kD as determined by SDS-PAGE.

The present invention also relates to ABC1 polypeptides comprising amino acid sequences which preferably have at least 98% identity over their entire length to the amino acid sequence of SEQ ID NO: 2. More preferably, the polypeptide has at least 99% identity over its entire length to the amino acid sequence of SEQ ID NO: 2. Most preferably, the polypeptide has
25 100% identity over its entire length to the amino acid sequence of SEQ ID NO: 2. As defined previously, the term "identity" refers to the degree of sequence relatedness between polypeptide sequences, which is further defined below.

Such related ABC1 polypeptides include substitution, deletion, and insertion variants, as well as allelic variants, splice variants, fragments, derivatives, and orthologs. Preferred
30 polypeptides and polypeptides fragments include those polypeptides and fragments that possess the biological activity of ABC1. In particular, those polypeptides and fragments that mediate reverse cholesterol transport are preferred. Also preferred are polypeptides and fragments that have improved reverse cholesterol transport activity.

Substitution, deletion, and insertion variants refer to ABC1 polypeptides comprising amino acid sequences that contain one or more amino acid sequence substitutions, deletions, and/or additions as compared to the ABC1 amino acid sequence set forth in SEQ ID NO: 2. In preferred embodiments, the variants have from about 1 to 5, or from about 1 to 10, or from about 1 to 20, or from about 1 to 40, or from about 1 to 65 amino acid substitutions, additions, and/or deletions. For example, the variants can have an addition of one or more amino acid residues anywhere in the polypeptide as well as at the carboxyl terminus and/or at the amino terminus, as long as the variant retains biological function. Also, for example, one or more amino acids can be deleted from any region of the polypeptide, including the carboxyl terminus and/or amino terminus, without substantial loss of biological function (Ron et al., *J. Biol. Chem.*, 268: 2984-2988 (1993); Dobeli et al., *J. Biotechnology*, 7: 199-216 (1988)). The amino acid substitution(s) can be conservative, non-conservative, or any combination thereof, as long as the ABC1 variant retains its biological activity. In addition, the substitution(s) can be with non-conserved amino acid residues, where the substituted residues may or may not be encoded by the genetic code, and with amino acid residues having a substituent group.

Suitable variants of ABC1 polypeptides can be determined using well-known techniques. For example, suitable ABC1 variants can be determined by identifying regions of the ABC1 molecule that may be changed without destroying biological activity. Also, as realized in the art, even regions that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure. Amino acid residues that can be changed without destroying biological activity can be determined by identifying regions of the ABC1 polypeptide that are not important for activity (Bowie et al., *Science*, 247: 1306-1310 (1990)). For example, ABC1 polypeptides from various species can be compared to determine the amino acid residues and regions of ABC1 molecules that are conserved across species. The conserved amino acid residues are likely important for biological function and/or structure. In contrast, changes in regions of the ABC1 molecule that are not conserved across species and are thus tolerated by natural selection would be less likely to adversely affect biological activity and/or structure. Accordingly, ABC1 polypeptides with additions, deletions, or substitutions in the non-conserved regions are likely suitable variants. Even in relatively conserved regions, chemically similar amino acids may be substituted for the naturally occurring residues while retaining activity.

In addition, suitable ABC1 variants can be identified using structure-function studies to determine residues in other members of the ATP-binding cassette protein family, such as ABCR and ABC-C, that are important for activity or structure. Such studies allow the prediction of

important amino acid residues in an ABC1 variant that correspond to amino acid residues that are important for activity or structure in other ATP-binding cassette proteins. For example, based on structure-function studies of other ATP-binding cassette proteins, important amino acid residues in ABC1 are likely found in regions associated with nucleotide binding and sterol transport.

- 5 Suitable variants include, for example, polypeptides having chemically similar amino acid substitutions for such predicted important amino acid residues of the ABC1 polypeptide.

Suitable ABC1 variants can also be determined using genetic engineering techniques to introduce amino acid changes at specific positions in order to identify regions critical for polypeptide function. Amino acid changes can be made using, for example, site-directed
10 mutagenesis or alanine-scanning mutagenesis (Cunningham et al., *Science*, 244: 1081-1085 (1989)). The resultant ABC1 variants can then be tested for biological activity using, for example, any one of the cholesterol efflux assays described herein. Variants having a particular amino acid residue substitution that results in destroyed cholesterol efflux activity would not be considered a suitable ABC1 variant.

- 15 Additional methods for identifying suitable variants are known in the art. Furthermore, one skilled in the art would realize amino acid changes that are likely to be permissive at certain amino acid positions in the protein (Bowie et al., *supra* (1990)). For example, it is generally known that the most buried or interior (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas fewer features of surface or exterior side chains
20 are generally conserved. Moreover, it is known that tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile, replacement of the hydroxyl residues Ser and Thr, replacement of the acidic residues Asp and Glu, replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His, replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-
25 sized amino acids Ala, Ser, Thr, Met, and Gly.

- The ABC1 variants can be naturally-occurring or artificially constructed. Examples of naturally-occurring variants are allelic variants and splice variants. Allelic variants refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism or population of organisms (Lewin, B., ed., *Genes II*, John Wiley & Sons, New York (1985)).
30 Allelic variants can vary at either the polynucleotide and/or polypeptide level. Splice variants refer to a nucleic acid molecule, usually RNA, which is generated by alternative processing of intron sequences in an RNA transcript, and the corresponding polypeptide. Alternatively, the ABC1 variants can be artificially constructed. For example, ABC1 variants can be constructed

using the technique of site-directed mutagenesis. Also, for example, ABC1 variants can be prepared from the corresponding nucleic acid molecules encoding said variants, which have a DNA sequence that varies as described from the wild-type DNA sequence as set forth in SEQ ID NO: 1.

5 Polypeptide fragments refer to polypeptides which comprise less than the full length amino acid sequence of ABC1 set forth in SEQ ID NO: 2. Preferred polypeptides fragments include those fragments that possess the biological activity of ABC1. In particular, those fragments that mediate reverse cholesterol transport or have improved reverse cholesterol transport activity are preferred. ABC1 fragments can have one or more amino acids deleted from
10 any region of the polypeptide, including the carboxyl terminus and/or amino terminus, as long as biological function is maintained. The ABC1 polypeptide fragments can occur naturally, such through alternative splicing or *in vivo* protease activity, or can be artificially constructed using well-known methods.

The invention also relates to ABC1 polypeptide derivatives, which refer to ABC1
15 polypeptides, variants, or fragments, as defined herein, that have been chemically modified. The derivatives are modified in a manner that is different from naturally-occurring ABC1 polypeptides, either in type or location of the molecules attached to the polypeptide. Derivatives may further include polypeptides formed by the deletion of one or more chemical groups which are naturally attached to the ABC1 polypeptides. In addition, the ABC1 polypeptide comprising
20 the amino acid sequence of SEQ ID NO: 2, as well as the above-described ABC1 variants and fragments, may be fused to a homologous polypeptide to form a homodimer or to a heterologous polypeptide to form a heterodimer.

Another aspect of the present invention relates to mutant ABC1 polypeptides and fragments thereof, corresponding to polypeptides isolated from Tangier patients. In one
25 preferred embodiment, the ABC1 polypeptide comprises SEQ ID NO: 8. The protein was isolated from a Tangier patient (TD1) and sequenced as described in Examples 1 and 5. The amino acid sequence set forth in SEQ ID NO: 8 is similar to the wild-type sequence with the exception of a glutamine to arginine residue substitution at position 537 (the residue number is that of Lawn et al., *J. Clin. Invest.*, 104: r25-31 (1999), corresponding to position 597 of SEQ ID
30 NO: 8). The location of this residue is within the amino-terminal hydrophilic domain, near the first predicted transmembrane domain. The substitution alters the charge of the amino acid in this region of the protein, resulting in an ABC1 protein that has significantly decreased cholesterol efflux activity, as shown in Figure 1.

In another preferred embodiment, the mutant ABC1 polypeptide comprises SEQ ID NO: 10. The protein corresponds to a polypeptide isolated from a Tangier patient (TD2) and sequenced as described in Examples 1 and 5. The amino acid sequence set forth in SEQ ID NO: 10 is similar to the wild-type sequence with the exception of an arginine to tryptophan substitution at residue 527 (the residue number is that of Lawn et al., *supra* (1999), corresponding to position 587 of SEQ ID NO: 10). Like the TD1 polypeptide, this substitution alters the charge of the amino acid residue in the amino-terminal hydrophilic domain of the ABC1 protein. The resultant mutant ABC1 protein also has significantly decreased cholesterol efflux activity, as shown in Figure 1.

ABC1 Polynucleotides

Another aspect of the present invention relates to isolated polynucleotides that encode the novel ABC1 polypeptides and variants thereof. The present invention provides, for example, isolated polynucleotides encoding the full-length ABC1 polypeptide, polynucleotides containing the full-length cDNA of wild-type ABC1, polynucleotides containing the entire length of the coding sequence of wild-type ABC1, and polynucleotides containing non-coding 5' and 3' sequences of ABC1, as well as polynucleotides of related ABC1 variants. The present invention also provides isolated polynucleotides that encode mutant ABC1 polypeptides, such as those of Tangier patients.

In one preferred embodiment, the isolated polynucleotide comprises a nucleotide sequence encoding the polypeptide comprising SEQ ID NO: 2. Importantly, in contrast to the published sequence of Langmann et al. which codes for a protein of 2201 amino acids based on a predicted start methionine found in exon 3 (Langmann et al., *Biochem. Biophys. Res. Comm.*, 257: 29-33 (1999) (GenBank Accession No. AJ012376), the presently claimed nucleotide sequence contains 50 exons and codes for a protein of 2261 amino acids (see Figure 4). The corresponding nucleotide sequence of the present invention contains a coding sequence that includes an additional 180 nucleotides at the 5' end corresponding to the following 60 amino-terminal amino acids:

MACWPQLRLLLWKNLTFRRRQTCQLLLEVAWPLFIFLILISVRLSYPPYEQHECHFPNKA

. Given that there is an in-frame stop codon 6 to 9 nucleotides upstream from this location, the newly predicted start site is the first methionine codon that could produce a continuous open reading frame. Alignment of this new ABC1 cDNA sequence with related ABC transporter sequences ABCR and ABC-C (also known as ABC3) which also contain open reading frames for

the 60 additional amino acids, indicates a high degree of similarity, implying that the homologous ABC transporter proteins begin with sequences related to the amino terminal extension sequence proposed for human ABC1. It is likely that the earlier published start site of the human ABC1 was predicted from the published mouse ABC1 cDNA sequence (Luciani et al., *Genomics*, 21:150-159 (1994); GenBank Accession no.: X75926) which contains an extra nucleotide "n" in the extension region such that the newly disclosed methionine is not in-frame. However, if the "n" nucleotide in the mouse sequence is ignored, the mouse and human sequences of the extension region are identical. In light of these results, it is likely that the full length human ABC1 protein contains 2261 amino acids rather than 2201 amino acids, as previously suggested by Langmann et al. and others. Accordingly, Langmann et al. do not present the full open reading frame of human ABC1.

In another preferred embodiment, the isolated polynucleotide comprises the full-length ABC1 cDNA, including at least a portion of either non-coding 5' and 3' sequences. Preferably, the polynucleotide comprises the nucleotide sequence shown in SEQ ID NO: 1. The 10.4 kb human ABC1 cDNA sequence shown in SEQ ID NO: 1 contains an open reading frame of 6783 nucleotides plus 5' and 3' untranslated regions. There is a start codon at position 291 and a stop codon at position 7074. The present ABC1 cDNA shown in SEQ ID NO: 1 differs from the published ABC1 cDNA (GenBank Accession No. AJ012376) in several respects. First, the present ABC1 cDNA includes an additional 350 nucleotides at the 5' end and an additional 3136 nucleotides at the 3' end (not including the poly(A) tail). The present ABC1 sequence also differs from the published ABC1 cDNA by the substitution of 10 nucleotides in the coding region. Of the ten differences, seven nucleotide differences predict amino acid changes. To remain consistent with published notation, the following nucleotide and amino acid numbers are those of Lawn et al., *J. Clin. Invest.*, 104: R25-31 (1999) and GenBank Accession No. AJ012376, rather than those of SEQ ID NO: 1. The nucleotide and amino acid changes are as follows: (1) A for G at nucleotide 414; (2) A for G at nucleotide 596 (K for R at amino acid 159); (3) T for C at nucleotide 705; (4) A for C at nucleotide 1980; (5) A for G at 2413 (I for V at amino acid 765); (6) G for A at 2589 (M for I at amino acid 823); (7) T for C at 4604 (I for T at amino acid 1495); (8) T for C at 4883 (L for P at amino acid 1588); (9) A for G at 5861 (K for R at amino acid 1914); and (10) T for C at 6443 (L for P at amino acid 2108). Five of the amino acid changes are conservative amino acid changes and may represent polymorphisms or sequence errors. In two instances, the present sequence predicts important amino acid differences from the GenBank sequence. The differences result in a leucine rather than a proline

at residue 1588 and at residue 2108. Interestingly, at both positions, the predicted leucine was also found in each of the three TD samples analyzed as well as the highly conserved mouse ABC1 protein sequence.

The present invention also relates to ABC1 polynucleotides comprising nucleotide
5 sequences that preferably have at least 80% identity over their entire length to the
polynucleotide comprising SEQ ID NO: 1. More preferably, the polynucleotide has at least
90% identity over its entire length to the polynucleotide comprising SEQ ID NO: 1. Even more
preferably, the polynucleotide has at least 95% identity over its entire length to the
polynucleotide comprising SEQ ID NO: 1. Most preferably the polynucleotide has 100%
10 identity over its entire length to the polynucleotide comprising SEQ ID NO: 1. Such related
ABC1 polynucleotides include substitution, deletion, and insertion variants, as well as allelic
variants, splice variants, fragments, derivatives, and orthologs, wherein one or more nucleotides
have been substituted, deleted, inserted, or derivatized. Preferred polynucleotides include those
polynucleotides that encode ABC1 polypeptides possessing biological activity, such as
15 cholesterol efflux activity.

In another preferred embodiment, the isolated polynucleotide comprises the entire coding
sequence of ABC1. In a particularly preferred embodiment, the polynucleotide comprises the
sequence shown as nucleotides 291-7074 of SEQ ID NO: 1. This isolated polynucleotide
contains an ABC1 open reading frame of 6783 nucleotides and encodes a polypeptide of 2261
20 amino acids, as described above.

In yet another preferred embodiment, the isolated polynucleotide comprises a nucleotide
sequence that encodes an ABC1 variant polypeptide. In particular, the isolated polynucleotide
comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence
which is at least 98% identical to the amino acid sequence of SEQ ID NO: 2. Also preferred is
25 an isolated polynucleotide comprising a nucleotide sequence that encodes a polypeptide
comprising an amino acid sequence which is at least 99% identical to the amino acid sequence of
SEQ ID NO: 2. Accordingly, the invention includes those polynucleotides that encode the
above-described ABC1 polypeptides, including the described substitution, deletion, and insertion
variants, as well as ABC1 allelic variants, splice variants, fragments, derivatives, fusion
30 polypeptides, and orthologs. Preferred polynucleotides are those polynucleotides that encode
polypeptides that possess the biological activity of ABC1. In particular, those polynucleotides
that encode polypeptides that mediate reverse cholesterol transport are preferred. Also preferred

are polynucleotides that encode polypeptides that have improved reverse cholesterol transport activity.

Yet another aspect of the invention relates to isolated polynucleotides that encode mutant ABC1 polypeptides from Tangier patients. In one preferred embodiment, the polynucleotide
5 encodes the polypeptide of SEQ ID NO: 8, which polypeptide is isolated from patient TD1 and is described above. In another preferred embodiment, the polynucleotide comprises the nucleotide sequence set forth in SEQ ID NO: 7. The nucleotide sequence set forth in SEQ ID NO: 7 contains the full open reading frame, as well as 5' and 3' flanking sequences. The open reading frame encodes a polypeptide of 2261 amino acids, containing, among other
10 substitutions, a nucleotide substitution that results in an A to G substitution at position 537 (using the numbering of Lawn et al., *supra* (1999)).

In another preferred embodiment relating to polynucleotides that encode mutant ABC1 polypeptides, the polynucleotide encodes the polypeptide of SEQ ID NO: 10, which polypeptide is isolated from Tangier patient TD2 and is also described above. In yet another preferred
15 embodiment, the polynucleotide comprises the nucleotide sequence set forth in SEQ ID NO: 9. The nucleotide sequence set forth in SEQ ID NO: 9 contains the full open reading frame, as well as 5' and 3' flanking sequences. The open reading frame encodes a polypeptide of 2261 amino acids, containing, among other substitutions, a polynucleotide substitution that results in an Arg to Trp substitution at residue 527 (using the numbering of Lawn et al., *supra* (1999)).

Another aspect of the present invention relates to isolated polynucleotides that comprise the non-coding 5' flanking and 3' flanking regions of ABC1. In one embodiment, the isolated polynucleotide comprises the non-coding 5' flanking region of ABC1. Preferably, the 5' flanking region contains, but is not limited to, the ABC1 promoter region. Thus, in a preferred
20 embodiment, the polynucleotide comprises the sequence shown in SEQ ID NO: 3. As demonstrated by heterologous reporter assays, discussed in Example 15, the polynucleotide set forth in SEQ ID NO: 3 contains the transcriptional regulatory region of the ABC1 gene. As shown in Figure 13, the polynucleotide set forth in SEQ ID NO: 3 is a 1643 b.p. non-coding sequence that contains several transcription regulatory elements, including a TATA box at positions 1522, 1435, and 1383, as well as transcription factor binding sites, including several
25 putative SP1 sites, and several nuclear receptor half sites. In addition, an identified sterol response element is found at position 1483-1500. Further heterologous reporter assays described in Example 17 revealed that several discrete portions of SEQ ID NO: 3 retained promoter activity. Accordingly, in another preferred embodiment, the polynucleotide comprises
30

nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3. In an especially preferred embodiment, the polynucleotide comprises nucleotides 1394-1532 of SEQ ID NO: 3, which sequence has been shown to have ABC1 promoter activity (see Example 17). In yet another preferred embodiment, the polynucleotide comprises nucleotides 1480-1510 of
5 SEQ ID NO: 3, which is shown to regulate the ABC1 transcriptional response to LXR ligands.

The 5' flanking polynucleotide also comprises a nucleotide sequence that hybridizes, under stringent conditions, to the nucleotide sequence set forth in SEQ ID NO: 3, wherein the nucleotide sequence has ABC1 promoter activity. In yet another embodiment, the polynucleotide comprises a nucleotide sequence that hybridizes, under stringent conditions, to
10 the nucleotide sequence comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3, wherein the nucleotide sequence has ABC1 promoter activity.

In another embodiment, the isolated polynucleotide comprises the 3' flanking region of ABC1. Several 3' untranslated regions have been identified which may represent alternate sites of polyadenylation of the ABC1 transcript. Preferably, the 3' flanking region contains
15 regulatory sequences. For example, the full length 3' UTR (SEQ ID NO: 6) contains 46 sequences (AA)nCU/UC(AA)n which have been shown to be necessary for binding of Vigilin. Vigilin, a ubiquitous protein with 14K homology domains, is the estrogen-inducible vitellogenin mRNA 3'-untranslated region binding protein (*J. Biol. Chem.*, 272: 12249-12252 (1997)). In addition to binding HDL, Vigilin has been shown to bind to the 3' flanking region of
20 mRNAs and to increase the half-life of the mRNA transcript (*Mol. Cell. Biol.*, 18:3991-4003 (1998)). Thus, the 3' flanking region could be altered, for example, to increase the binding of Vigilin, thereby increasing the half-life of the ABC1 mRNA. Preferably, the isolated polynucleotide comprises the sequence shown in SEQ ID NO: 4. Also preferably, the isolated polynucleotide comprises the sequence shown in SEQ ID NO: 5. In another preferred
25 embodiment, the isolated polynucleotide comprises the sequence shown in SEQ ID NO: 6. In other preferred embodiments, the polynucleotide comprises a sequence that hybridizes, under stringent conditions, to the nucleotide sequence set forth in SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6.

The present invention also includes related ABC1 5' and 3' flanking polynucleotides.
30 Accordingly, the invention relates to polynucleotides comprising nucleotide sequences that have at least 80% identity over their entire length to the polynucleotide comprising SEQ ID NO: 3, the polynucleotide comprising SEQ ID NO: 4, the polynucleotide comprising SEQ ID NO: 5, the polynucleotide comprising SEQ ID NO: 6, or the polynucleotide comprising nucleotides 1-

1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3. Preferably, the polynucleotide has at least 80%, more preferably at least 90%, even more preferably at least 95%, and most preferably 100% identity over its entire length to any one of the aforementioned flanking polynucleotides. Preferred polynucleotides include those polynucleotides that possess
5 biological activity, such as transcriptional regulatory activity.

It is understood that the present invention further relates to isolated polynucleotides that are complementary to any one of the above-described polynucleotide sequences. As used herein, the term "complementary" refers to the hybridization or base pairing between nucleotides, such as, for example, between the two strands of a double-stranded polynucleotide.
10 Two single-stranded nucleotide molecules are said to be complementary when the nucleotides of one strand, optimally aligned with appropriate nucleotide insertions, deletions or substitutions, pair with at least about 80% of the nucleotides of the other strand.

Another aspect of the present invention relates to compositions comprising the novel ABC1 polynucleotides described above and a suitable carrier. In one embodiment, the composition comprises a polynucleotide encoding the polypeptide comprising SEQ ID NO: 2, a
15 polynucleotide comprising SEQ ID NO: 1, a polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, or a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 98% identical to the amino acid sequence of SEQ ID NO: 2 and a suitable carrier. In another embodiment, the composition comprises a polynucleotide having at
20 least 80%, preferably 90%, or more preferably 95% identity over its entire length to the polynucleotide comprising SEQ ID NO: 1 and a suitable carrier.

In another embodiment, the composition comprises a polynucleotide comprising an ABC1 5' flanking sequence and a suitable carrier. Preferably, the composition comprises a polynucleotide comprising SEQ ID NO: 3 or a polynucleotide comprising nucleotide fragments
25 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3 and a suitable carrier. Also preferably, the composition comprises a polynucleotide having at least 80%, 90%, or 95% identity over its entire length to the polynucleotide comprising any one of the described 5' flanking sequences and a suitable carrier. In yet another embodiment, the composition comprises a polynucleotide comprising an ABC1 3' flanking sequence and a suitable carrier.
30 Preferred compositions comprise a polynucleotide comprising SEQ ID NO: 4, a polynucleotide comprising SEQ ID NO: 5, or a polynucleotide comprising SEQ ID NO: 6, as well as a polynucleotide having at least 80%, preferably 90%, or more preferably 95% identity over its entire length to any of these polynucleotides, and a suitable carrier. Still other compositions of

the invention comprise mutant ABC1 polynucleotides. Preferably, the composition comprises a polynucleotide comprising SEQ ID NO: 7 or a polynucleotide comprising SEQ ID NO: 9 and a suitable carrier.

In addition, a composition of the present invention may comprise, in any combination, two or more of the above-described ABC1 polynucleotides and a suitable carrier. Any suitable aqueous carrier can be used in the composition. Preferably, the carrier renders the composition stable at a desired temperature, such as room temperature or storage temperature (i.e. 4°C to -20°C), and is of approximately neutral pH. Examples of suitable carriers are known to those of skill in the art and include Tris-EDTA buffer and DEPC-H₂O.

ABC1 Vectors and Host Cells

The present invention also relates to recombinant vectors that comprise one or more of the above-described ABC1 polynucleotides, host cells that are genetically engineered with the vectors comprising ABC1 polynucleotides, and the production of ABC1 polypeptides by recombinant techniques. As mentioned, the invention provides recombinant vectors that comprise one or more of the above-described wild-type ABC1 polynucleotides. In preferred embodiments, the recombinant vector comprises the polynucleotide encoding the polypeptide comprising SEQ ID NO: 2, the polynucleotide comprising SEQ ID NO: 1, and the polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1. In another preferred embodiment, the recombinant vector comprises the variant polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 98% identical to the amino acid sequence of SEQ ID NO: 2. Also, in another preferred embodiment, the recombinant vector comprises a variant polynucleotide that is at least 80% identical to a polynucleotide comprising SEQ ID NO: 1. In still another preferred embodiment, the recombinant vector comprises a polynucleotide that is complementary to any of these polynucleotides.

In another embodiment, the recombinant vector comprises a polynucleotide comprising a mutant ABC1 polynucleotide isolated from a Tangier disease patient. In a preferred embodiment, the recombinant vector comprises the polynucleotide comprising SEQ ID NO: 8. In another preferred embodiment, the recombinant vector comprises the polynucleotide comprising SEQ ID NO: 10. The recombinant vectors may also comprise a polynucleotide sequence that is complementary to these sequences.

It is also understood that the recombinant vector may also comprise, in any combination, two or more of the above-described wild-type, variant, or mutant ABC1 polynucleotides.

An isolated ABC1 polynucleotide, such as any of the above-described wild-type, variant, or mutant polynucleotides, is inserted into a vector using well-known ligation and cloning techniques. Cloning techniques have been described in several standard laboratory manuals, including Davis et al., *Basic Methods in Molecular Biology* (1986); Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., (Cold Spring Harbor Press, Cold Spring Harbor, N.Y. 5 (1989)); Ausubel et al. eds., *Current Protocols in Molecular Biology*, (Wiley and Sons (1994)); Goeddel, ed., *Gene Expression Technology* (Methods in Enzymology (1991)); Murray, ed., *Gene Transfer and Expression Protocols* (Human Press, Clifton, NJ).

Any vector suitable for ABC1 polynucleotide insertion can be used. The vector is 10 typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery such that amplification and/or expression of the gene can occur). Preferably, the vector is compatible with bacterial, insect, or mammalian host cells. Also preferably, the vector is an expression vector (for a review of expression vectors, see Goeddel, D.V. ed., *Methods Enzymol.*, Academic Press Inc., San Diego, CA (1990)). The 15 vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such 20 sequences, collectively referred to as "flanking sequences" should preferably include one or more of the following nucleotide sequences: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region 25 for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element.

The flanking sequences may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of flanking sequences from more than one source), or synthetic. Also, the 30 flanking sequences may be native sequences which normally function to regulate ABC1 polypeptide expression. The source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

The vector should also preferably include at least one selectable marker for propagation in a host. A selectable marker is a gene element that encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Suitable selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, tetracycline, or kanamycin for prokaryotic host cells; (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers include zeocin, G418, hygromycin, or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin, or ampicillin resistance for culturing in *E. coli* and other bacteria.

Other suitable selection genes include those that are used to amplify the expressed gene. Amplification is the process wherein genes that are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell transformants are placed under selection pressure wherein only the transformants are uniquely adapted to survive by virtue of the selection gene present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to the amplification of both the selection gene and the ABC1 gene. As a result, increased quantities of ABC1 polypeptide are synthesized from the amplified DNA.

The vector should also preferably contain a transcription termination sequence, which is typically located 3' of the end of a polypeptide coding region and serves to terminate transcription. Usually, the transcription termination sequence in prokaryote cells is a G-C rich fragment followed by a poly T sequence. The sequence can be purchased as part of a commercial vector or synthesized using well-known methods for nucleic acid synthesis.

The vector should also preferably contain a ribosome binding site, which is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of the ABC1 polypeptide to be expressed. The Shine-Dalgarno sequence is varied but is typically a polypurine (*i.e.*, having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using well-known methods.

The vector should also preferably contain a promoter that is recognized by the host organism and operably linked to the encoded polynucleotide. The promoter can be an inducible

promoter or a constitutive promoter. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. In contrast, constitutive promoters initiate continual gene product production; consequently there is little or no control over gene expression. A suitable promoter is operably linked to a polynucleotide, by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. As mentioned, a native promoter can be used to direct amplification and/or expression of the polynucleotide. Thus, the recombinant vector can comprise any one of the above-described wild-type, variant, or mutant ABC1 polynucleotides and an ABC1 promoter, such as that found in SEQ ID NO: 3. The recombinant vector can also comprise, in any combination, two or more of the above-described wild-type, variant, or mutant ABC1 polynucleotides and an ABC1 promoter.

Preferably, a heterologous promoter is used if it permits greater transcription and higher yields of ABC1 protein as compared to the native ABC1 promoter, and if it is compatible with the host cell system that has been selected for use. The heterologous promoter can be used alone or in conjunction with the native ABC1 promoter. Thus, in one preferred embodiment, the recombinant vector comprises any one of the above-described wild-type ABC1 polynucleotides and a heterologous promoter. In another preferred embodiment, the recombinant vector comprises any one of the above-described variant ABC1 polynucleotides and a heterologous promoter. In yet another preferred embodiment, the recombinant vector comprises any one of the above-described mutant ABC1 polynucleotides and a heterologous promoter. Preferred embodiments also include recombinant vectors that contain any combination of two or more of the above-described wild-type, variant, and mutant ABC1 polynucleotides and a heterologous promoter.

Heterologous promoters suitable for use with prokaryotic hosts include, but are not limited to, the beta-lactamase and lactose promoter systems (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-31), alkaline phosphatase, a tryptophan (trp) promoter system, and hybrid promoters such as the tac promoter (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-31). Their sequences have been published, thereby enabling one skilled in the art to ligate them to the desired DNA sequence, using linkers or adapters as needed to supply any useful restriction sites. Other suitable heterologous promoters will be known to those skilled in the art.

Suitable heterologous promoters for use with mammalian host cells are also well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, retroviruses, hepatitis-B virus and Simian Virus 40 (SV40). Other
5 suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter. Additional suitable promoters include, but are not limited to: the SV40 early promoter and late promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-10); the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, *et al.*, 1980, *Cell* 22:787-97); the herpes thymidine kinase promoter (Wagner *et al.*,
10 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-45); and the regulatory sequences of the metallothioneine gene (Brinster *et al.*, 1982, *Nature* 296:39-42). Preferably, the promoter is a cytomegalovirus or SV40 promoter. Thus, in especially preferred embodiments, the recombinant vector comprises one of the above-described wild-type ABC1 polynucleotides, one of the above-described variant ABC1 polynucleotides, or one of the above-described mutant ABC1
15 polynucleotides and a cytomegalovirus promoter. In another especially preferred embodiment, the recombinant vector comprises, in any combination, two or more of the above-described wild-type, variant, or mutant ABC1 polynucleotides and a cytomegalovirus promoter.

The vector also preferably contains an enhancer sequence to increase the transcription of a polynucleotide, such as ABC1. Suitable enhancers for the activation of eukaryotic promoters
20 include viral enhancers, such as the SV40, cytomegalovirus early promoter, polyoma, and adenovirus enhancers.

Expression vectors of the invention may be constructed from a starting vector, such as a commercially available vector, which may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already
25 present in the vector, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

Preferred vectors are those which are compatible with bacterial, insect, and mammalian host cells. Vectors preferred for use in bacteria include, for example, pQE70, pQE60, and pQE-9 (Quiagen, Inc.), pBluescript vectors, Phagescript vectors, pNH16A, pNH18A, pNH46A
30 (Stratagene Cloning Systems, Inc.), ptrc99a, pKK223-3, pDR540, pRIT5 (Pharmacia Biotech, Inc.), and pCEP4 (Invitrogen Corp., Carlsbad, CA). Preferred eukaryotic vectors include, but are not limited to, pWLNEO, pSV2CAT, pOG44, pXTI and pSG (Stratagene), pSVK3, pBPV, pMSG, and pSVL (Pharmacia), and pGL3 (Promega, Madison, WI). Other suitable vectors will

be readily apparent to the skilled artisan.

In an especially preferred embodiment, the recombinant vector comprises pCEPhABC1, which is described in Example 4 and shown in Figure 3. The recombinant vector pCEPhABC1 comprises the plasmid pCEP4 (Invitrogen), an expression vector containing the cytomegalovirus promoter and enhancer. The vector pCEPhABC1 further comprises an ABC1 polynucleotide operatively linked to the heterologous cytomegalovirus promoter. The ABC1 polynucleotide comprises SEQ ID NO: 1, which contains the full-length ABC1 cDNA, including non-coding 5' flanking (i.e., native ABC1 promoter) and 3' flanking sequences.

In addition, the present invention provides recombinant vectors comprising ABC1 flanking sequence polynucleotides. In one embodiment, the recombinant vector comprises a polynucleotide comprising a ABC1 5' flanking sequence that preferably contains promoter activity. Thus, in a preferred embodiment, the recombinant vector comprises the polynucleotide comprising SEQ ID NO: 3. In another preferred embodiment, the recombinant vector comprises the polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3. In an especially preferred embodiment, the polynucleotide comprises nucleotides 1394-1532 of SEQ ID NO: 3. Also, in another embodiment, the recombinant vector comprises a polynucleotide that hybridizes, under stringent conditions, to the polynucleotide set forth in SEQ ID NO: 3 or the polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3. In yet another embodiment, the polynucleotide comprises a nucleotide sequence that has at least 80%, more preferably at least 90%, and even more preferably at least 95% identity over its entire length to the polynucleotide comprising SEQ ID NO: 3 or the polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3. The recombinant vector also comprises a polynucleotide that is complementary to any of the above-described 5' flanking sequences.

In another embodiment, the recombinant vector comprises a polynucleotide comprising a 3' flanking sequence of ABC1. In a preferred embodiment, the recombinant vector comprises the polynucleotide comprising SEQ ID NO: 4. In another preferred embodiment, the recombinant vector comprises the polynucleotide comprising SEQ ID NO: 5. In an equally preferred embodiment, the recombinant vector comprises the polynucleotide comprising SEQ ID NO: 6. Also, in another embodiment, the recombinant vector comprises a polynucleotide that hybridizes, under stringent conditions, to the polynucleotide set forth in SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6. In yet another embodiment, the polynucleotide comprises a

nucleotide sequence that has at least 80%, more preferably at least 90%, and even more preferably at least 95% identity over its entire length to the polynucleotide comprising SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6. The recombinant vector also comprises a polynucleotide that is complementary to any of the above-described 3' flanking sequences.

5 An isolated ABC1 flanking sequence polynucleotide, such as any of the above-described 5' or 3' flanking sequence polynucleotides, is inserted into a vector using well-known ligation and cloning techniques. Any of the previously described vectors can be used. Preferably, the vector is compatible with bacterial, insect, or mammalian host cells. Also preferably, the vector is an expression vector. The vector may be, for example, a phage, plasmid, viral, or retroviral
10 vector.

 In addition to the ABC1 flanking sequence, the vector may contain one or more of the following flanking nucleotide sequences: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion,
15 a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Any of the previously described flanking nucleotide sequences are suitable. The flanking sequences may be homologous, heterologous, hybrid, or synthetic. Also, the flanking sequences may be native sequences which normally function to regulate ABC1 polypeptide expression. The source of a
20 flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

 Preferred vectors are those which are compatible with bacterial, insect, and mammalian host cells. Suitable vectors have been previously described and include pQE70, pQE60, pQE9,
25 pBluescript vectors, Phagescript vectors, pNH16A, pNH18A, pNH46A, ptrc99a, pKK223-3, pDR540, pRIT5, and pCEP4 for use in bacteria and pWLNEO, pSV2CAT, pOG44, pXTI, pSG, pSVK3, pBPV, pMSG, pSVL, and pGL3 for use in eukaryotic cells.

 In one particularly preferred embodiment, the recombinant vector comprises a polynucleotide comprising the 5' flanking region of the ABC1 gene and further comprises at least
30 one polynucleotide encoding a heterologous polypeptide. The heterologous polynucleotide is operatively linked to the ABC1 5' flanking sequence. The ABC1 5' flanking sequence preferably contains the ABC1 promoter. Thus, preferably, the 5' flanking sequence comprises the sequence set forth in SEQ ID NO: 3. Equally preferably, the 5' flanking sequence comprises nucleotides

1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3. The heterologous polynucleotide preferably encodes a polypeptide that is a complete protein or a biologically active fragment of a protein. The vector may also contain more than one heterologous polynucleotide. More preferably, the heterologous polynucleotide encodes a reporter protein. In
5 such case, the recombinant vector preferably does not contain any additional promoter sequences. Examples of suitable reporter proteins include luciferase, β -galactosidase, chloramphenicol acetyl transferase, acetyl transferase, and green fluorescent protein. Preferably, the reporter polypeptide is luciferase. Thus, in one especially preferred embodiment, the recombinant vector comprises the 5' flanking sequence set forth in SEQ ID NO: 3 and a
10 luciferase reporter polynucleotide. In other equally preferred embodiments, the recombinant vector comprises a polynucleotide comprising nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3 and a luciferase reporter polynucleotide.

Expression vectors comprising the ABC1 5' flanking sequence can be constructed from a starting vector, such as a commercially available vector, which contains a reporter
15 polynucleotide. Examples of suitable expression vectors include pGL3-Basic, which contains a luciferase reporter gene (Promega, Madison, WI) and p β Gal-Basic (Clontech, Palo Alto, CA). A preferred vector is the pGL3-Basic luciferase reporter vector, which is promoterless. A 5' flanking sequence containing the ABC1 promoter, for example SEQ ID NO: 3, can be ligated into one of the above expression vectors using well-known methods, including the methods
20 described herein (see Example 15). Thus, in an especially preferred embodiment, the recombinant vector is pAPR1, a reporter gene construct comprising SEQ ID NO: 3 and a luciferase reporter gene in a pGL3 vector (see Figure 11).

The present invention also relates to host cells comprising any one of the above-described recombinant vectors. After the vector has been constructed, the completed vector can be inserted
25 into a suitable host cell for amplification and/or polypeptide expression. Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast cell, an insect cell, or a vertebrate cell). The host cell, when cultured under appropriate conditions, synthesizes an ABC1 polypeptide or, alternatively, a reporter polypeptide, which can be subsequently measured. The host cell can be a mammalian host cell, such as a primate cell line or a rodent cell line,
30 including transformed cell lines. Normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Candidate host cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene.

A number of suitable host cells are known in the art and many are available from the American Type Culture Collection (ATCC), Manassas, VA. Suitable mammalian host cells include, but are not limited to, chinese hamster ovary cells (CHO), CO DHFR- cells (Urlaub *et al.*, 1980, *Proc. Natl. Acad. Sci.*, 97:4216-20), human embryonic kidney (HEK) 293 or 293T
5 cells, or 3T3 cells, monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa cells, mouse I-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HAK hamster cell lines, Thp-1, HepG2, and mouse RAW cell lines. Each of these cell lines is known by and available to those skilled in the art of protein expression. A preferred host cell is the
10 mouse monocytic cell line RAW 264.7, which use is described in Example 8.

Suitable bacterial host cells include various strains of *E. coli* (e.g., HB101, DH50, DH10, and MC1061), which are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas spp.*, other *Bacillus spp.*, *Streptomyces spp.*, and the like may also be employed in this method. Also, many strains of yeast cells are also available as host cells for the
15 expression of ABC1 polypeptides. Preferred yeast cells include, for example, *Saccharomyces cerevisiae* and *Pichia pastoris*. Additionally, insect cell systems may be suitable host cells. Insect cell systems are described, for example, in Kitts *et al.*, 1993, *Biotechniques*, 14:810-17; Lucklow, 1993, *Curr. Opin. Biotechnol.* 4:564-72; and Lucklow *et al.*, 1993, *J. Virol.*, 67:4566-79. Preferred insect cells are Sf-9 and Hi5 (Invitrogen).

20 The present invention also provides a method for expressing an ABC1 protein in a mammalian host cell comprising the steps of: (a) transfecting the mammalian host cell with a recombinant expression vector comprising a polynucleotide encoding ABC1 in an amount sufficient to produce a detectable level of ABC1 protein; and (b) purifying the produced ABC1 protein. In one preferred embodiment, the recombinant expression vector comprises a
25 polynucleotide encoding the polypeptide comprising SEQ ID NO: 2. In another preferred embodiment, the polynucleotide comprising SEQ ID NO: 1. In yet another preferred embodiment, the polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1. In still another preferred embodiment, the polynucleotide encoding a polypeptide that is at least 98% identical to the polypeptide comprising SEQ ID NO: 2.

30 Introduction of the recombinant ABC1 vector into a mammalian host cell can be effected by methods well-known in the art and described in standard laboratory manuals, such as Sambrook, *supra*. Preferably, the recombinant vector is introduced into a host cell in a precipitate or in a complex with a charged lipid. Suitable methods for introduction of the ABC1

vector include calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods known in the art. These methods are described, for example, in Davis et al., *Basic Methods in Molecular Biology* (1986). If the recombinant ABC1 vector is a viral vector, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

The ABC1 polypeptide can be recovered and purified from recombinant cell cultures using well-known methods, including ammonium sulfate or ethanol precipitation, acid, extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography (see, e.g., Smith and Johnson, *Gene* 67:31-40 (1988)). Preferably, affinity chromatography using anti-ABC1 antibodies is employed for purification.

In addition, the present invention provides a method for expressing ABC1 protein in a mammalian subject comprising the step of administering to a mammalian subject a recombinant expression vector comprising a polynucleotide encoding ABC1 in an amount sufficient to express ABC1 protein in said mammalian subject. Preferably, the recombinant expression vector comprises a polynucleotide encoding the polypeptide comprising SEQ ID NO: 2, the polynucleotide comprising SEQ ID NO: 1, the polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, or the polynucleotide encoding a polypeptide that is at least 98% identical to the polypeptide comprising SEQ ID NO: 2. Introduction of the recombinant ABC1 vector into a mammalian subject can be effected by methods well-known in the art and are described in detail herein below. Expression of ABC1 can be measured by obtaining a blood sample from the subject to whom the recombinant ABC1 vector was administered, separating the monocyte population, and measuring the ABC1 gene expression in macrophage cells. The ABC1 gene expression can be measured using methods well-known in the art, such as RT-PCR, and methods described herein (see Examples 9 and 10). The level of ABC1 protein can be measured by obtaining a blood sample from the subject, separating the monocyte population and measuring the ABC1 protein in macrophage cells using well-known methods, such as immunoprecipitation, described herein at Example 11.

Methods and Compounds for Increasing Cholesterol Efflux

In another aspect of the present invention, a method suitable for increasing cholesterol efflux from cells in a mammalian subject is provided. Such method comprises administering to

the mammalian subject a recombinant expression vector comprising an ABC1 polynucleotide in an amount sufficient to increase cholesterol efflux from said cells. The recombinant vector can be any of the above-described vectors containing any of the previously described wild-type or variant ABC1 polynucleotides, as long as the encoded ABC1 polypeptide has biological activity
5 (i.e., cholesterol transport activity). Preferably, the recombinant vector comprises the polynucleotide encoding the polypeptide comprising SEQ ID NO: 2, the polynucleotide comprising SEQ ID NO: 1, the polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, or the polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 98% identical to the amino acid sequence of SEQ ID NO: 2. Also, preferably, the
10 recombinant vector comprises a variant polynucleotide that is at least 80%, 90%, or 95% identical to the polynucleotide comprising SEQ ID NO: 1, as long as the encoded ABC1 polypeptide has cholesterol transport activity.

The administration of a recombinant ABC1 expression vector to a mammalian subject can be used to express the ABC1 gene in said subject for the treatment of cardiovascular disease.
15 Specifically, this method would achieve its therapeutic effect by the introduction of the ABC1 gene into macrophage cells and other cholesterol-accumulating cells found in the arterial lesions of mammals with cardiovascular disease. Expression of the ABC1 polynucleotide in target cells would effect greater production of the ABC1 protein. The subsequently produced ABC1 protein would ameliorate the disease by increasing the efflux of cholesterol from macrophage and other
20 cholesterol-laden cells found in arterial lesions onto nascent HDL particles. The cellular efflux would lead to the overall removal of cholesterol from peripheral sites, such as the cholesterol-rich core of arterial plaques. A concurrent reduction in the size of these pathological lesions reduces the risk of arterial blockage that leads to heart attacks and angina. This method could also be used prophylactically to prevent the accumulation of cholesterol in arterial walls.

25 A sufficient amount of a ABC1 expression vector is the amount of ABC1 vector that increases cholesterol efflux from the cells of a mammalian subject. Such amount can be determined by measuring the cholesterol efflux in the cells of a subject before (control level) and after administration of the recombinant ABC1 expression vector at various dosages and determining the dose that effects an increase in cholesterol efflux compared to control level. The
30 cholesterol efflux can be measured by obtaining a blood sample from the subject, separating the monocyte population, and measuring the amount of cholesterol efflux in a subject's macrophage cells. Any of the assays described herein can be used to measure cholesterol efflux.

In addition, cholesterol efflux can be measured by determining the level of plasma HDL-cholesterol in a subject before (control) and after administration of a recombinant ABC1 expression vector. An observed increase in HDL-cholesterol in the serum of a subject following administration of a recombinant ABC1 expression vector indicates an increase in cholesterol efflux. HDL-cholesterol levels in serum can be determined using methods well-known in the art.

In addition, cholesterol efflux can be measured by measuring the size of atherosclerotic lesions found in the arterial wall of a subject before (control level) and after administration of the recombinant ABC1 expression vector. A reduction in the size of the arterial lesion indicates an increase in cholesterol efflux. Assays for measuring arterial lesions are well known in the art. For example, increased cholesterol efflux from arterial lesions can be measured using the mouse model of atherosclerosis described in Lawn et al., *Nature*, 360: 670-672 (1992)), as well as any of the other known models. Using the LDL receptor knockout mouse described in Lawn et al., cholesterol efflux can be measured before and after administration of the ABC1 vector. Fatty streak lesion size in groups of animals fed an atherogenic diet can be measured by oil-red O staining of aortic sections as described in Lawn et al. A reduction in size of fatty streak lesions in the group receiving the ABC1 expression vector compared to a group receiving a control vector indicates an increase in cholesterol efflux from the lesions. In humans, the size of atherosclerotic lesions found in arterial walls can be measured using, for example, angiography and non-invasive ultrasound methods.

Alternatively, cholesterol efflux can be measured by obtaining a blood sample and measuring the level of ABC1 mRNA or ABC1 protein in the macrophage cells of a subject before and after administration of a recombinant ABC1 expression vector. Routine assays can be performed to determine the correlation between increasing ABC1 mRNA concentrations and cholesterol efflux. Likewise, assays can be performed to correlate the amount of ABC1 protein with the amount of cholesterol efflux. Using such correlation data, an observed increase in ABC1 mRNA or ABC1 protein in the cells of a subject following administration of a recombinant ABC1 expression vector can be used to indicate an increase in cholesterol efflux. ABC1 mRNA and ABC1 protein levels can be measured using the assays described herein and other well-known techniques for mRNA and protein quantitation. Therapeutic dosages and formulations are discussed in further detail below.

There are available to one skilled in the art multiple viral and non-viral methods suitable for introduction of a nucleic acid molecule into a target cell. For example, viral delivery vectors suitable for gene therapy include, but are not limited to, adenovirus, herpes simplex virus, pox

virus (i.e., vaccinia), hepatitis virus, parvovirus, papovavirus, alphavirus, coronavirus, rhabdovirus, papilloma virus, adeno-associated virus (AAV), polio virus, and RNA viruses, such as a retroviruses and Sindbis virus. The ABC1 polynucleotide can also be delivered using a non-viral delivery system, such as naked DNA delivery (direct injection), receptor-mediated transfer
5 (DNA-ligand complexes), electroporation, adenovirus-ligand-DNA complexes, calcium phosphate (CaPO_4) precipitation, microparticle bombardment (gene gun techniques), liposome-mediated transfer, and lipofection.

Genetic modification of a cell may be accomplished using one or more techniques well known in the gene therapy field (Mulligan, R., 1993, *Science*, 260 (5110): 926-32). Gene
10 therapy materials and methods can include inducible promoters, tissue-specific enhancer-promoters. DNA sequences designed for site-specific integration, DNA sequences capable of providing a selective advantage over the parent cell, labels to identify transformed cells, negative selection systems and expression control systems (safety measures), cell-specific binding agents (for cell targeting), cell-specific internalization factors, and transcription factors to enhance
15 expression by a vector as well as methods of vector manufacture. Examples of methods and materials for the practice of gene therapy techniques are described in U.S. Patent Nos. 4,970,154 (involving electroporation techniques), 5,679,559 (describing a lipoprotein-containing system for gene delivery), 5,676,954 (involving liposome carriers), 5,593,875 (describing methods for calcium phosphate transfection), and 4,945,050 (describing a process wherein biologically active
20 particles are propelled at cells at a speed whereby the particles penetrate the surface of the cells and become incorporated into the interior of the cells), and PCT Pub. No. WO 96/40958 (involving nuclear ligands).

Adenoviral vectors have proven especially useful for gene transfer into eukaryotic cells (Rosenfeld, M., *et al.*, *Science*, 252: 431-4 (1991); U.S. Patent No. 5,631,236). The first trial of
25 Ad-mediated gene therapy in human was the transfer of the cystic fibrosis transmembrane conductance regulator (CFTR) gene to lung (Crystal, R., *et al.*, 1994, *Nat. Genet.*, 8 (1): 42-51). Experimental routes for administering recombinant Ad to different tissues *in vivo* have included intratracheal instillation (Rosenfeld, M., *et al.*, 1992, *Cell*, 68 (1): 143-55) injection into muscle (Quantin, B., *et al.*, 1992, *Proc. Natl. Acad. Sci. U.S.A.*, 89 (7): 2581-4), peripheral intravenous
30 injection (Herz, J., and Gerard, R., 1993, *Proc. Natl. Acad. Sci. U.S.A.*, 90 (7): 2812-6) and stereotactic inoculation to brain (Le Gal La Salle, G., *et al.*, 1993, *Science*, 259 (5097): 988-90). The adenoviral vector, then, is widely available to one skilled in the art and is suitable for use in the present invention.

Adeno-associated virus (AAV) has recently been introduced as a gene transfer system with potential applications in gene therapy. Wild-type AAV demonstrates high-level infectivity, broad host range and specificity in integrating into the host cell genome (Hermonat, P., and Muzyczka, N., 1984, *Proc. Natl. Acad. Sci. U.S.A.*, 81 (20): 6466-70). Herpes simplex virus type-1 (HSV-1) is a preferred vector system (Geller, A., *et al.*, 1991, *Trends Neurosci.*, 14 (10): 428-32; Glorioso, J., *et al.*, 1995, *Mol. Biotechnol.*, 4 (1): 87-99; Glorioso, J., *et al.*, 1995, *Annu. Rev. Microbiol.*, 49: 675-710). Vaccinia virus, of the poxvirus family, has also been developed as an expression vector (Smith, G., and Moss, B., 1983, *Gene*, 25 (1): 21-8; Moss, B., 1992, *Biotechnology*, 20: 345-62; Moss, B., 1992, *Curr. Top. Microbiol. Immunol.*, 158: 25-38). Each of the above-described vectors is widely available to one skilled in the art and would be suitable for use in the present invention.

Preferably, the viral delivery system utilizes a retroviral vector. Retroviral vectors are capable of infecting a large percentage of the target cells and integrating into the cell genome (Miller, A., and Rosman, G., *Biotechniques*, 7(9): 980-2, 984-6, 989-90 (1989); U.S. Patent No. 5,672,510). Retroviruses were developed as gene transfer vectors relatively earlier than other viruses, and were first used successfully for gene marking and transducing the cDNA of adenosine deaminase (ADA) into human lymphocytes. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus, or is a lentiviral vector. An especially preferred retroviral vector is a lentiviral vector. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), SIV, BIV, HIV and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated.

Since recombinant retroviruses are defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. The helper plasmids are missing a nucleotide sequence that enables the packaging mechanism to recognize an RNA transcript for encapsitation. Thus, helper cell lines produce empty virions, since no genome is packaged. Suitable helper cell lines include, but are not limited to Ψ 2, PA317 and PA12. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced. The vector virions

produced by this method can then be used to infect a tissue cell line, such as NIH 3T3 cells, to produce large quantities of chimeric retroviral virions.

The vectors of the present invention may be constructed using standard recombinant techniques widely available to one skilled in the art. Such techniques may be found in common molecular biology references such as *Molecular Cloning: A Laboratory Manual* (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press), *Gene Expression Technology* (Methods in Enzymology, Vol. 185, edited by D. Goeddel, 1991, Academic Press, San Diego, CA), and *PCR Protocols: A Guide to Methods and Applications* (Innis, et al. 1990, Academic Press, San Diego, CA).

In order to obtain transcription of the ABC1 polynucleotide within a target cell, a transcriptional regulatory region capable of driving gene expression in the target cell must be utilized. The transcriptional regulatory region preferably comprises a promoter and/or enhancer, which is operatively linked to the ABC1 polynucleotide. The promoter can be homologous or heterologous to the ABC1 gene, provided that it is active in the cell or tissue-type into which the construct will be inserted. The transcriptional regulatory region chosen should drive high level gene expression in the target cell. Preferably, a macrophage-specific promoter, such as a scavenger receptor type A, matrix metalloproteinase promoter (MMP-12), or macrophage-tropic lentivirus promoter (Fabunmi et al., *Atherosclerosis*, 148: 375-386 (1999)), is used. A particularly preferred promoter is the 5' region of the scavenger receptor type A gene, which contains a strong macrophage promoter that can be used to drive the transcription of the ABC1 gene. In addition, a means to increase endogenous ABC1 polypeptide expression in a cell is to insert one or more enhancer elements into the promoter region, where the enhancer elements can serve to increase transcriptional activity of the ABC1 gene. Similarly, the enhancer element(s) used is selected based on the tissue in which one desires to activate the gene. Thus, for example, enhancer elements known to confer promoter activation in cells found in arterial tissue, especially macrophage cells, will be selected. Other transcriptional regulatory regions suitable for use in the present invention include but are not limited to the human cytomegalovirus (CMV) immediate-early enhancer/promoter, the SV40 early enhancer/promoter, the JC polyomavirus promoter, the albumin promoter, PGK and the α -actin promoter coupled to the CMV enhancer (Doll, R., et al., 1996, *Gene Ther.*, 3 (5): 437-47).

Other components of the vector construct may optionally include DNA molecules designed for site-specific integration (e.g., endogenous sequences useful for homologous recombination), tissue-specific promoters, DNA molecules capable of providing a selective

advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (*e.g.*, for cell targeting), cell-specific internalization factors, transcription factors enhancing expression from a vector, and factors enabling vector production.

5 In one embodiment, the vector can include targeting DNA for site-specific integration. The targeting DNA is a nucleotide sequence that is complementary (homologous) to a region of the gene of interest, for example, the ABC1 gene. Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to the ABC1 gene by attaching it to the targeting DNA. DNA sequences for insertion include, for example, regions of DNA that
10 may interact with or control the expression of an ABC1 polypeptide, *e.g.*, flanking sequences. Thus, the expression of the desired ABC1 polypeptide is achieved not by transfection of DNA that encodes the ABC1 gene itself, but rather by the use of targeting DNA coupled with DNA regulatory segments that provide the endogenous gene sequence with recognizable signals for transcription of an ABC1 polypeptide (Sauer, *Curr. Opin. Biotechnol.*, 5:521-27 (1994); Sauer,
15 *Methods Enzymol.*, 225:890-900 (1993)).

 In yet other embodiments, regulatory elements can be included for the controlled expression of the ABC1 gene in the target cell. Such elements are turned on in response to an appropriate effector. In this way, a therapeutic ABC1 polypeptide can be expressed when desired. One conventional control means involves the use of small molecule dimerizers or
20 rapalogs to dimerize chimeric proteins which contain a small molecule-binding domain and a domain capable of initiating a biological process, such as a DNA-binding protein or transcriptional activation protein (*see* PCT Pub. Nos. WO 96/41865, WO 97/31898, and WO 97/31899). The dimerization of the proteins can be used to initiate transcription of the transgene. Another suitable control means or gene switches includes the use of mifepristone (RU486),
25 which is a progesterone antagonist. The binding of a modified progesterone receptor ligand-binding domain to the progesterone antagonist activates transcription by forming a dimer of two transcription factors that then pass into the nucleus to bind DNA. The ligand-binding domain is modified to eliminate the ability of the receptor to bind to the natural ligand. The modified steroid hormone receptor system is further described in U.S. Patent No. 5,364,791 and PCT Pub.
30 Nos. WO 96/40911 and WO 97/10337. Yet another control means uses a positive tetracycline-controllable transactivator. This system involves a mutated tet repressor protein DNA-binding domain (mutated tet R-4 amino acid changes which resulted in a reverse tetracycline-regulated transactivator protein, *i.e.*, it binds to a tet operator in the presence of tetracycline) linked to a

polypeptide which activates transcription. Such systems are described in U.S. Patent Nos. 5,464,758, 5,650,298, and 5,654,168. Additional expression control systems and nucleic acid constructs are described in U.S. Patent Nos. 5,741,679 and 5,834,186.

Viral delivery vectors containing an ABC1 polynucleotide can be made target specific by
5 altering the viral coat such that it contains a ligand that is specific for another molecule found on the target cell. This will allow the vector to bind specifically to the desired cell-type. The ligand can be any compound of interest that will bind specifically to a molecule found on the target cell, such as a cell-surface receptor. Preferably, the receptor is found exclusively on target cells and not on other cells. For example, ligands for scavenger receptor A can be used to direct viral
10 delivery vectors to macrophage cells. Alternatively, the viral coat can be altered such that it contains an antibody or antibody fragment, such as Fab, or F(ab')₂, that recognizes and binds to an antigenic epitope on the target cells. The viral coat can be altered by inserting an additional polynucleotide that encodes the ligand into the viral genome. Those of skill in the art will know of other specific polynucleotide sequences which can be inserted into the viral genome to allow
15 target specific delivery of the vector containing the ABC1 polynucleotide.

In addition, a naked ABC1 polynucleotide can be administered. ABC1 polynucleotides and recombinant ABC1 expression vectors, such as those described above, can be administered as a pharmaceutical composition. Such a composition comprises an effective amount of an ABC1 polynucleotide or recombinant ABC1 expression vector, as previously defined herein, and
20 a pharmaceutically acceptable formulation agent selected for suitability with the mode of administration. Suitable formulation materials preferably are non-toxic to recipients at the concentrations employed and are described herein below.

The pharmaceutical composition comprising an ABC1 polynucleotide or an ABC1 recombinant expression vector may contain formulation materials for modifying, maintaining, or
25 preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption, or penetration of the composition. Suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine, or lysine), antimicrobials, antioxidants (such as ascorbic acid, sodium sulfite, or sodium hydrogen-sulfite), buffers (such as borate, bicarbonate, Tris-HCl, citrates,
30 phosphates, or other organic acids), bulking agents (such as mannitol or glycine), chelating agents (such as ethylenediamine tetraacetic acid (EDTA)), complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin, or hydroxypropyl-beta-cyclodextrin), fillers, monosaccharides, disaccharides, and other carbohydrates (such as glucose, mannose, or

dextrins), proteins (such as serum albumin, gelatin, or immunoglobulins), coloring, flavoring and diluting agents, emulsifying agents, hydrophilic polymers (such as polyvinylpyrrolidone), low molecular weight polypeptides, salt-forming counterions (such as sodium), preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid, or hydrogen peroxide), solvents (such as glycerin, propylene glycol, or polyethylene glycol), sugar alcohols (such as mannitol or sorbitol), suspending agents, surfactants or wetting agents (such as pluronics; PEG; sorbitan esters; polysorbates such as polysorbate 20 or polysorbate 80; triton; tromethamine; lecithin; cholesterol or tyloxapal), stability enhancing agents (such as sucrose or sorbitol), tonicity enhancing agents (such as alkali metal halides – preferably sodium or potassium chloride – or mannitol sorbitol), delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. See *Remington's Pharmaceutical Sciences* (18th Ed., A.R. Gennaro, ed., Mack Publishing Company 1990).

The pharmaceutically active compounds (i.e. ABC1 polynucleotide or ABC1 vector) can be processed in accordance with conventional methods of pharmacy to produce medicinal agents for administration to patients, including humans and other mammals. Thus, the pharmaceutical composition comprising an ABC1 polynucleotide or an ABC1 recombinant expression vector may be made up in a solid form (including granules, powders or suppositories) or in a liquid form (e.g., solutions, suspensions, or emulsions). Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral or parenteral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting sweetening, flavoring, and perfuming agents. For example, a suitable carrier for injection may be water, physiological saline solution, or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary carriers. Other

exemplary pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute.

The dosage regimen for treating a cardiovascular disease with a composition comprising an ABC1 polynucleotide or ABC1 expression vector is based on a variety of factors, including the type of cardiovascular disease, the age, weight, sex, medical condition of the patient, the severity of the condition, the route of administration, and the particular compound employed. Thus, the dosage regimen may vary widely, but can be determined routinely using standard methods. For example, the amount of ABC1 polynucleotide or ABC1 expression vector to be administered is an amount sufficient to increase cholesterol efflux from the cells of a mammalian subject. Such amount can be determined, for example, by measuring the plasma HDL-cholesterol level of a subject before and after administration of the ABC1 polynucleotide or ABC1 expression vector. A sufficient amount of ABC1 polynucleotide or ABC1 expression vector is an amount that increases the plasma HDL-cholesterol level of a subject. Accordingly, the clinician can titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. A typical dosage may range from about 0.1 μ g/kg to about 100 mg/kg or more, depending on the factors mentioned above.

The frequency of dosing will depend upon the pharmacokinetic parameters of the ABC1 polynucleotide or vector in the formulation being used. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them.

Appropriate dosages may be ascertained through use of appropriate dose-response data.

The cells of a mammalian subject may be transfected *in vivo*, *ex vivo*, or *in vitro*.

Administration of an ABC1 polynucleotide or a recombinant vector containing an ABC1 polynucleotide to a target cell *in vivo* may be accomplished using any of a variety of techniques well known to those skilled in the art. For example, U.S. Patent No. 5,672,344 describes an *in vivo* viral-mediated gene transfer system involving a recombinant neurotrophic HSV-1 vector.

The above-described compositions of ABC1 polynucleotides and recombinant ABC1 vectors can be transfected *in vivo* by oral, buccal, parenteral, rectal, or topical administration as well as by inhalation spray. The term "parenteral" as used herein includes, subcutaneous, intravenous, intramuscular, intrasternal, infusion techniques or intraperitoneally.

For oral administration, the pharmaceutical composition containing the ABC1 polynucleotide or recombinant ABC1 vector may be in the form of, for example, a capsule, a tablet, a suspension, or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a given amount of DNA or viral vector particles. For example, these
5 may contain an amount from about 10^3 - 10^{15} viral particles, preferably from about 10^6 - 10^{12} viral particles. A suitable daily dose for a human or other mammal may vary widely depending on the condition of the patient and other factors, but, once again, can be determined using routine methods.

The pharmaceutical composition containing the ABC1 DNA or recombinant ABC1
10 vector can also be administered rectally. Suitable suppositories for rectal administration of the vector can be prepared by mixing the vector with a suitable non-irritating excipient such as cocoa butter and polyethylene glycols that are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the vector.

A pharmaceutical composition also can be formulated for inhalation. For example, an
15 ABC1 polynucleotide or vector may be formulated as a dry powder for inhalation. Also, ABC1 polynucleotide or vector inhalation solutions can be formulated with a propellant for aerosol delivery. In yet another embodiment, solutions may be nebulized.

The pharmaceutical composition containing the ABC1 DNA or recombinant ABC1 vector can also be injected. Injectable preparations, such as sterile injectable aqueous or
20 oleaginous suspensions, may be formulated according to the known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. A particularly suitable carrier for parenteral injection is sterile distilled water in which an ABC1 polynucleotide or vector is formulated as a
25 sterile, isotonic solution, properly preserved. Among the other acceptable carriers and solvents that may be employed are Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed, including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables. Yet another
30 preparation can involve the formulation of the desired ABC1 molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds, beads or liposomes, that provides for the controlled or sustained release of the ABC1 product (*see, e.g.* PCT/US93/00829; Epstein et al., *Proc. Natl. Acad. Sci.*, 82: 3688-3692 (1985)).

Also, the pharmaceutical composition containing the ABC1 DNA or recombinant ABC1 vector can administered topically. For topical administration, the vector may comprise from 0.001% to 10% w/w, *e.g.*, from 1% to 2% by weight of the formulation, although it may comprise as much as 10% w/w, but preferably not more than 5% w/w, and more preferably from 0.1% to 1% of the formulation. Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin (*e.g.*, liniments, lotions, ointments, creams, or pastes) and drops suitable for administration to the eye, ear, or nose. A suitable topical dose of active ingredient of a vector of the present invention is administered one to four, preferably two or three times daily.

While the nucleic acids and /or vectors of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more vectors of the invention or other agents. When administered as a combination, the therapeutic agents can be formulated as separate compositions that are given at the same time or different times, or the therapeutic agents can be given as a single composition.

In another embodiment of the present invention, a target cell is transfected *in vivo* by implantation of a "producer cell line" in proximity to the target cell population (Culver, K., *et al.*, 1994, *Hum. Gene Ther.*, 5 (3): 343-79; Culver, K., *et al.*, *Cold Spring Harb. Symp. Quant. Biol.*, 59: 685-90; Oldfield, E., 1993, *Hum. Gene Ther.*, 4 (1): 39-69). The producer cell line is engineered to produce a viral vector containing the ABC1 polynucleotide and to release its viral particles in the vicinity of the target cells, *i.e.* preferably macrophage cells found in atherosclerotic lesions. A portion of the released viral particles contact the target macrophage cells and infect those cells, thus delivering an ABC1 polynucleotide to the target macrophage cell. Following infection of the target cell, expression of the ABC1 polynucleotide occurs, providing the macrophage cell with functional ABC1 protein.

In another embodiment, the invention provides a method of treating a cardiovascular disease by the *ex vivo* introduction of an ABC1 polynucleotide or recombinant ABC1 expression vector. In such instances, cells, tissues, or organs that have been removed from the patient are exposed to ABC1 compositions after which the cells, tissues, or organs are subsequently implanted back into the patient. For example, one method includes the removal of a blood sample from a subject with cardiovascular disease, enriching the sample for monocytes, and contacting the isolated monocytes with a recombinant expression vector containing the ABC1 polynucleotide and, optionally, a target specific gene. Optionally, the monocyte cells can be treated with a growth factor, such as GM-CSF, to stimulate cell growth, before reintroducing the

cells into the subject. When reintroduced, the cells will specifically target the cell population from which they were originally isolated. In this way, the transport activity of the ABC1 polypeptide may be used to promote cholesterol efflux in a subject.

Another method of *ex vivo* administration involves introducing the ABC1 polynucleotide or recombinant ABC1 vector into the mammalian subject by means of skin transplants of cells containing the virus. Preferably, a retrovirus used for this method of administration. Long term expression of foreign genes in implants, using cells of fibroblast origin, may be achieved if a strong housekeeping gene promoter is used to drive transcription. For example, the dihydrofolate reductase (DHFR) gene promoter may be used. Cells such as fibroblasts, can be infected with virions containing a retroviral construct containing the ABC1 polynucleotide together with a gene which allows for specific targeting, such as scavenger receptor A, and a strong promoter. The infected cells can be embedded in a collagen matrix that can be grafted into the connective tissue of the dermis in the recipient subject. As the retrovirus proliferates and escapes the matrix it will specifically infect the target cell population. In this way the transplantation results in increased amounts of cholesterol efflux activity in cells manifesting the transport disorder.

In another embodiment, the recombinant expression vector comprising the ABC1 polynucleotide can be administered using *in vitro* techniques, such as described in U.S. Patent No. 5,399,346. For example, an ABC1 polypeptide can be delivered by implanting certain cells that have been genetically engineered, using methods such as those described herein, to express the ABC1 polypeptide. In order to minimize a potential immunological reaction in patients being administered an ABC1 polypeptide, as may occur with the administration of a polypeptide of a foreign species, it is preferred that the natural cells producing ABC1 polypeptide be of human origin and produce human ABC1 polypeptide. Thus, it is preferred that the recombinant cells producing ABC1 polypeptide be transformed with an expression vector containing a gene encoding a human ABC1 polypeptide. The cells can be autologous or heterologous. Optionally, the cells can be immortalized. In order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues. The transfected cells can be administered to a patient using the above-described methods.

Techniques for the encapsulation of living cells are known in the art, and the preparation of the encapsulated cells and their implantation in patients may be routinely accomplished. For example, Baetge *et al.* (PCT Pub. No. WO 95/05452 and PCT/US94/09299) describe membrane capsules containing genetically engineered cells for the effective delivery of biologically active molecules. The capsules are biocompatible and are easily retrievable. The capsules encapsulate cells transfected with recombinant DNA molecules comprising DNA sequences coding for biologically active molecules operatively linked to promoters that are not subject to down regulation *in vivo* upon implantation into a mammalian host. The devices provide for the delivery of the molecules from living cells to specific sites within a recipient. In addition, *see* U.S. Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating living cells is described in PCT Pub. No. WO 91/10425 (Aebischer *et al.*). *See also*, PCT Pub. No. WO 91/10470 (Aebischer *et al.*); Winn *et al.*, 1991, *Exper. Neurol.* 113:322-29; Aebischer *et al.*, 1991, *Exper. Neurol.* 111:269-75; and Tresco *et al.*, 1992, *ASAIO* 38:17-23.

Another delivery system for polynucleotides encoding ABC1 is a "non-viral" delivery system. Techniques that have been used or proposed for gene therapy include DNA-ligand complexes, adenovirus-ligand-DNA complexes, direct injection of DNA, CaPO₄ precipitation, gene gun techniques, electroporation, lipofection, and colloidal dispersion (Mulligan, R., 1993, *Science*, 260 (5110): 926-32). Any of these methods are widely available to one skilled in the art and would be suitable for use in the present invention. Other suitable methods are available to one skilled in the art, and it is to be understood that the present invention may be accomplished using any of the available methods of transfection. Several such methodologies have been utilized by those skilled in the art with varying success (Mulligan, R., 1993, *Science*, 260 (5110): 926-32).

Preferably, the non-viral delivery system is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome, which is an artificial membrane vesicle useful as delivery vehicles *in vitro* and *in vivo*. Liposomes are self-assembling, colloidal particles in which a lipid bilayer, composed of amphiphilic molecules such as phosphatidyl serine or phosphatidyl choline, encapsulates a portion of the surrounding media such that the lipid bilayer surrounds a hydrophilic interior. Unilamellar or multilamellar liposomes can be constructed such that the interior contains a desired chemical, drug, or, as in the instant invention, an isolated DNA molecule. For example, it has been shown that large unilamellar vesicles (LUV), which

range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules, such as RNA, DNA and intact virions. Once encapsulated within the aqueous interior, these macromolecules can be delivered to mammalian cells in a biologically active form (Fraley, R. and Papahadjopoulos, D. 1981, *Trends Biochem. Sci.*, 6: 77-80). In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, R., *et al.*, 1988, *Biotechniques*, 6 (7): 682-90).

The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations. Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebroside, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.

The targeting of liposomes has been classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries. Active targeting, on the other hand, involves alteration of the liposome by coupling the liposome to a specific ligand, such as a polyclonal or monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the liposome in order to achieve targeting to organs and cell types other than the naturally occurring sites of localization. Preferably, the ligand is a polyclonal or monoclonal antibody which can be used to target liposomes to specific cell-surface ligands. The ligand can also be an antibody fragment, such as Fab, or F(ab')_2 , as long as it binds efficiently to an the antigenic epitope on the target cells. Preferably, the antibody or antibody fragment recognizes an antigen that is found exclusively on the target cell. For

example, certain antigens expressed specifically on macrophage cells, such as scavenger receptor A, may be exploited for the purpose of targeting antibody-ABC1 liposomes directly to a macrophage cell.

5 A number of procedures can be used to covalently attach either polyclonal or monoclonal antibodies to a liposome bilayer. Also, lipid groups can be incorporated into the lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. In addition, various linking groups can be used for joining the lipid chains to the targeting ligand.

Studies presented herein showed that ligands for nuclear receptors were able to effect an increase in cholesterol efflux. Accordingly, in another embodiment, the present invention provides a method suitable for increasing cholesterol efflux from cells in a mammalian subject comprising the step of administering to the mammalian subject at least one ligand for a nuclear receptor in an amount sufficient to increase cholesterol efflux from said cells. A pharmaceutical composition comprising a ligand for a nuclear receptor can be prepared and administered using the above-described methods for formulating and administering pharmaceutical compositions. A sufficient amount of a nuclear receptor ligand is the amount of ligand that increases cholesterol efflux. Such amount can be determined by measuring the cholesterol efflux before and after administration of the ligand at various dosages and determining the dose that effects an increase in cholesterol efflux. The cholesterol efflux can be measured using assays previously described.

20 Nuclear receptors are ligand-activated transcription factors that play a critical role in vertebrate development and adult physiology by transducing the effects of small, lipophilic hormones into transcriptional responses. Several families of nuclear receptors exist, including peroxisome proliferator-activated receptors (PPARs), liver X receptors (LXR), retinoid X receptor (RXR), the farnesoid X receptor (FXR), and the steroid and xenobiotic receptor (SXR). The PPAR family comprises the three closely related gene products PPAR α , PPAR γ , and PPAR β/δ . PPAR α has been implicated as a key regulator of intra- and extracellular lipid metabolism. When bound to fatty acids, PPAR α stimulates the proliferation of peroxisomes and induces the synthesis of several enzymes involved in the β -oxidation of fatty acids. The PPAR α receptor is also the molecular target for the fibrates, drugs that are prescribed for the reduction of high triglyceride levels (Isseman et al., *Nature*, 347, 645 (1990)). Fibrates act as PPAR ligands to regulate the transcription of a large number of genes that affect lipoprotein and fatty acid metabolism. In addition, PPAR γ ligands, such as compounds belonging to the class of

thiazolidinediones compounds, have been shown to increase HDL and reduce triglyceride levels in humans.

LXR is an oxysterol receptor that regulates the catabolism of excess cholesterol. LXR α has been shown to bind as a heterodimer with RXR to a DNA response element in the CYP7a gene, which encodes the enzyme responsible for the rate-limiting step in the conversion of cholesterol to bile acids. Studies have shown that mice lacking LXR α accumulate enormous amounts of cholesterol esters in their livers when fed a cholesterol-rich diet, due to their inability to increase CYP7a gene transcription in response to dietary cholesterol (Peet et al., *Cell*, 93: 693 (1998)). Further studies with LXR α null mice show that LXR α is also involved in the regulation of several other genes that participate in cholesterol and fatty acid homeostasis. The biological role of a closely related nuclear receptor, LXR β , which is expressed in several tissues and is activated by the same oxysterols as LXR α , remains unclear (Song et al., *Proc. Natl. Acad. Sci.*, 91: 10809 (1994); Seol, *Mol. Endocrinol.*, 9: 72 (1995)).

FXR, which is evolutionarily related to LXR α , is also involved in cholesterol homeostasis. Like LXR α , FXR functions as a heterodimer with the RXR receptor (Schwartz et al., *Curr. Opin. Lipidol.*, 9: 113 (1998); Vlahcevic et al., *Gastroenterology*, 113: 1949 (1997)). FXR is activated by the synthetic retinoid TTNPB and superphysiological concentrations of all-trans retinoic acid (Zavacki et al., *Proc. Natl. Acad. Sci.*, 94: 7909 (1997)). Recent studies indicate that FXR is a nuclear bile acid receptor. First, FXR is abundantly expressed in tissues through which bile acids circulate, including the liver, intestine, and kidney (Seol et al., *Mol. Endocrinol.*, 9: 72 (1995); Forman et al., *Cell*, 81: 687 (1995)). Also, FXR has recently been shown to serve as a receptor for physiological concentrations of several bile salts, among which chenodeoxycholic acid (CDCA) is the most potent (Kliwer et al., *Science*, 284: 757-284 (1999); Makishima et al., *Science*, 284: 1362-1363 (1999)). CDCA is known to regulate the expression of several genes that participate in bile salt homeostasis, including those encoding CYP7a and the intestinal bile acid-binding protein.

As described in detail in Example 13, ligands for LXR, RXR, and PPAR nuclear receptors were shown to increase apoAI-induced cholesterol efflux in cholesterol-loaded mouse macrophage cells. For example, administration of 9 *cis*-retinoic acid (30 ng/ml) produced approximately a 3-fold increase in apoAI-induced cholesterol efflux in these cells. Similarly, administration of 22(R)hydroxycholesterol (5 μ g/ml) produced approximately a 3-fold increase in apoAI-induced cholesterol efflux. Cells that received fenfibrate (3 μ g/ml) produced an approximate 2-fold increase in cholesterol efflux. These results indicate that nuclear receptors

may be modulated to increase the rate of apolipoprotein-mediated cholesterol efflux from macrophages. Furthermore, as described in Example 13, 9-*cis*-RA mediated cholesterol efflux from macrophage cells in a dose-dependent manner. Other nuclear receptor activators, such as bezafibrate, were shown to increase cholesterol efflux (data not shown).

5 Accordingly, in the method for increasing cholesterol efflux by administering a nuclear receptor ligand, the ligand is preferably selected from the group consisting of LXR, RXR, PPAR, FXR, and SXR nuclear receptor ligands. In the preferred embodiment wherein a LXR ligand is used to increase cholesterol efflux, the ligand is more preferably selected from the group consisting of 20(S) hydroxycholesterol, 22(R) hydroxycholesterol, 24-hydroxycholesterol, 25-
10 hydroxycholesterol, and 24(S), 25 epoxcholesterol LXR ligands. Most preferably, the LXR ligand is 20(S) hydroxycholesterol. In the preferred embodiment wherein a RXR ligand is used to increase cholesterol efflux, the ligand is more preferably selected from the group consisting of 9-*cis* retinoic acid, retinol, retinal, all-trans retinoic acid, 13-*cis* retinoic acid, acitretin, fenretinide, etretinate, CD 495, CD564, TTNN, TTNNPB, TTAB, LGD 1069. Most preferably,
15 the RXR ligand is 9-*cis* retinoic acid. In another preferred embodiment wherein a PPAR ligand is used to increase cholesterol efflux, the ligand is preferably selected from the class of thiazolidinedione compounds.

 In another preferred embodiment, more than one nuclear receptor ligand is administered to the mammalian subject to increase cholesterol efflux. Preferably, when two or more nuclear
20 receptor ligands are administered to a subject, the ligands are an LXR and an RXR ligand. More preferably, the nuclear receptor ligands are 20(S) hydroxycholesterol and 9-*cis* retinoic acid.

 In still another embodiment, the present invention provides a method suitable for increasing cholesterol efflux from cells in a mammalian subject comprising the step of administering to the mammalian subject an eicosanoid in an amount sufficient to increase
25 cholesterol efflux. A pharmaceutical composition comprising an eicosanoid can be prepared and administered using the above-described methods for formulating and administering pharmaceutical compositions. A sufficient amount of eicosanoid is the amount that increases cholesterol efflux. Such amount can be determined by measuring the cholesterol efflux before and after administration of the eicosanoid at various dosages and determining the dose that
30 effects an increase in cholesterol efflux. The cholesterol efflux can be measured using assays and methods previously described.

 As described in Example 14, eicosanoids were shown to increase apoAI-induced cholesterol efflux in cholesterol-loaded mouse macrophage cells. For example, administration of

PGI₂ (25 nm) produced approximately a 2-fold increase in apoAI-induced cholesterol efflux in these cells. Likewise, administration of PGE₁ (25 nM) produced approximately a 3-fold increase in apoAI-induced cholesterol efflux. These results demonstrate that eicosanoids can increase the rate of apolipoprotein-mediated cholesterol efflux from macrophages. Accordingly, in a preferred embodiment, the eicosanoid is selected from the group consisting of prostaglandin E₂, prostacyclin (prostaglandin I₂), and prostaglandin J₂ eicosanoids.

Methods and Compounds for Increasing ABC1 Expression/Activity

Given that ABC1 functions to promote cholesterol efflux, one way to increase cholesterol efflux is to increase the cellular expression of ABC1. Accordingly, the present invention also provides methods suitable for increasing cholesterol efflux from cells in a mammalian subject by administering to the mammalian subject a therapeutic amount of a compound that increases the expression of ABC1 in said cells. A therapeutic amount of compound is the amount of compound that increases ABC1 expression. Such amount can be determined by measuring the gene expression of ABC1 before and after administration of the compound at various dosages and determining the dose that effects an increase in ABC1 gene expression. The ABC1 gene expression can be measured by obtaining a blood sample from the subject, separating the monocyte population, and determining the concentration of ABC1 mRNA using methods known in the art and described herein, such as RT-PCR.

In one preferred embodiment, the method comprises administering a cAMP analogue to increase the gene expression of ABC1. As shown in Figure 8, cAMP increases the expression of ABC1 mRNA in normal fibroblast cells approximately 10-fold. Preferably, the cAMP analogue is selected from the group consisting of 8-bromo cAMP, N⁶-benzoyl cAMP, and 8-thiomethyl cAMP. In another preferred embodiment, the method comprises administering a compound that increases the synthesis of cAMP to increase the gene expression of ABC1. Preferably, the compound is forskolin. In yet another preferred embodiment, the method comprises administering a compound that inhibits the breakdown of cAMP to increase the gene expression of ABC1. An example of such a compound is a phosphodiesterase inhibitor. Preferably, the phosphodiesterase inhibitor is selected from the group consisting of rolipram, theophylline, 3-isobutyl-1-methylxanthine, R020-1724, vinpocetine, zaprinast, dipyridamole, milrinone, amrinone, pimobendan, cilostamide, enoximone, peroximone, and vesnarinone phosphodiesterase inhibitors.

In another preferred embodiment, the method comprises administering to the mammalian subject a ligand for a nuclear receptor in an amount sufficient to increase the gene expression of ABC1. As described in Example 17 and shown in Figure 12, ligands for nuclear receptors can up-regulate the gene expression of ABC1. Transfection studies using pAPR1, which contains a luciferase reporter gene under the control of the ABC1 promoter, showed that the ABC1 promoter was activated in the presence of ligands for LXR and RXR nuclear receptors. Specifically, macrophage cells transfected with pAPR1 produced a 19-fold increase in luciferase reporter activity in the presence of 20 OH-chol, a 16-fold increase in luciferase activity in the presence of 9-*cis* RA, and a 280-fold increase in luciferase activity in the presence of both ligands compared with EtOH control. The results indicate that both sterols and retinoids elicit a strong transcription response from the ABC1 promoter. Further, there is an apparent synergistic effect between the two classes of compounds, as can be seen by the dramatic increase in luciferase activity found in cells treated with both ligands. In accordance with the inventive method, preferably, the ligand is selected from the group consisting of LXR, RXR, PPAR, FXR, and SXR ligands.

In addition to increasing cellular levels of ABC1 protein, reverse cholesterol transport can be promoted by enhancing the activity of ABC1 protein. Thus, in another embodiment, the present invention provides a method suitable for increasing cholesterol efflux from cells in a mammalian subject comprising the step of administering to the mammalian subject a therapeutic amount of a compound that increases ABC1 activity in an amount sufficient to increase cholesterol efflux. A pharmaceutical composition comprising such a compound can be prepared and administered using the above-described methods for formulating and administering pharmaceutical compositions. A therapeutic amount of compound is the amount of compound that increases cholesterol efflux. Such amount can be determined by measuring the cholesterol efflux before and after administration of the compound at various dosages and determining the dose that effects an increase in cholesterol efflux using methods previously described. To determine whether an increase in cholesterol efflux is due to an increase in ABC1 activity, the amount of ABC1 protein present in a cell sample before and after administration of the compound is determined, using methods described herein (see Example 11). For both measurements (i.e. pre- and post-administration of the compound), the amount of cholesterol efflux activity is divided by the concentration of ABC1 protein to determine the amount of cholesterol activity found in a standard concentration of ABC1 protein. An observed increase in

cholesterol activity standardized to protein concentration indicates that the increase is due to an increase in ABC1 activity.

Methods for Identifying Therapeutic Compounds

5 Another aspect of the present invention relates to methods for screening a compound to determine whether the compound modulates (i.e., up-regulates or down-regulates) the gene expression of ABC1. Such compounds may be useful in the development of therapeutic compounds that increase ABC1 expression and thereby promote cholesterol efflux and raise blood levels of HDL-cholesterol. Accordingly, methods for identifying compounds that may be
10 useful in the treatment of cardiovascular disease are provided. In one embodiment, the present invention provides a method for screening a test compound for ABC1 expression modulating activity comprising the steps of: (a) operatively linking a reporter cDNA with an expression modulating portion of the mammalian ABC1 gene to produce a recombinant reporter construct; (b) transfecting the recombinant reporter construct into a population of host cells; (c) assaying
15 the level of reporter gene expression in a sample of the transfected host cells; (d) contacting the transfected host cells with the test compound being screened; (e) assaying the level of reporter gene expression in a sample of the transfected host cells after contact with the test compound; and (f) comparing the relative change in the level of reporter gene expression caused by exposure to the test compound, thereby determining the ABC1 expression modulating activity.

20 First, a recombinant reporter construct comprising a heterologous reporter gene operatively linked to an expression modulating portion of the ABC1 gene is constructed. The ABC1 expression modulating polynucleotide and reporter gene can be inserted into a vector using well-known ligation and cloning techniques, such as those described herein and in standard laboratory manuals, including Davis et al., *Basic Methods in Molecular Biology*
25 (1986); Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., (Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1989)); and Ausubel et al. eds., *Current Protocols in Molecular Biology*, (Wiley and Sons (1994)). Alternatively, the ABC1 expression modulating polynucleotide can be inserted into a commercially available reporter construct, such as those previously described. Any vector suitable for ABC1 polynucleotide and reporter gene insertion
30 can be used. The chosen vector should be functional in the particular host cell employed. Preferably, the vector is compatible with mammalian host cells.

Preferably, the expression modulating portion of the ABC1 gene is the 5' flanking region of ABC1, containing ABC1 promoter activity. In one preferred embodiment, the expression

modulating portion of the ABC1 gene comprises SEQ ID NO: 3. In another preferred embodiment, the expression modulating portion of the ABC1 gene comprises nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, 1394-1643, or 1394-1532 of SEQ ID NO: 3. Also, preferably the heterologous reporter is selected from the group consisting of polynucleotides that encode the luciferase, β -galactosidase, chloramphenicol, and green fluorescent proteins. More preferably, the heterologous reporter is a polynucleotide that encodes the luciferase protein. In a particularly preferred embodiment, the recombinant reporter construct is pAPR1, shown in Figure 11.

Next, the recombinant reporter construct is transfected into a population of host cells.

The recombinant reporter construct can be introduced into the host cells using any of the previously described transfection methods, as well as the methods described in Examples 8 and 15. For example, the reporter construct can be transfected using calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or any of the other known and described methods (see, e.g., Davis et al., *Basic Methods in Molecular Biology* (1986)). The host cell can be any cell that, when cultured under appropriate conditions, synthesizes a reporter polypeptide, which can be subsequently measured. Preferably, the host cell is a mammalian host cell. More preferably, the mammalian host cell is a macrophage, fibroblast, hepatic, or intestinal cell. Most preferably, the host cell is selected from the group consisting of RAW 264.7 cells, Thp-1 cells, and HepG2 cells. The concentration of reporter construct and duration of the transfection can vary, depending on the transfection method and the type and concentration of host cell used. Determination of the appropriate concentration of reporter construct and transfection time is well within the skill of the ordinary artisan.

Following transfection, a sample of transfected host cells that was not exposed to the test compound is assayed to determine the level of reporter gene expression. The level of reporter gene expression found in the sample of unexposed transfected host cells provides a control measurement. The transfected host cells are lysed and the level of reporter gene expression is assayed using any of the methods well-known in the art. The assays used to measure the level of reporter gene expression differ, depending on the reporter construct used in the transfections. For example, if a luciferase reporter construct is used, the luciferase activity of the cell lysate is measured as light units using a luminometer, as described in Example 15.

A different sample of the transfected host cells is contacted with the test compound being screened and the level of reporter gene expression found in these cells is subsequently measured.

Preferably, the transfected host cells are contacted with the test compound for about 4-48 hours. More preferably, the transfected host cells are contacted with the test compound for about 8-36 hours. Even more preferably, the transfected host cells are contacted with the test compound for about 24 hours. The same assay used to measure the level of reporter gene expression in
5 unexposed (control) cells should be used to measure the level of reporter gene expression in the cells exposed to the test compound.

Finally, the level of reporter gene expression found in unexposed control cells is compared with the level of reporter gene expression found in cells exposed to the test compound to determine whether the test compound has ABC1 expression modulating activity. If the level
10 of gene expression in both cell samples are the same or about the same, the test compound does not modulate ABC1 gene expression. A higher level of reporter gene expression in cells exposed to the test compound relative to the level of reporter gene expression found in control cells, indicates that the test compound up-regulates the gene expression of ABC1. A lower level of
reporter gene expression in cells exposed to the test compound relative to the level of reporter
15 gene expression found in control cells, indicates that the test compound down-regulates the gene expression of ABC1.

Another aspect of the present invention relates to methods for screening a test compound to determine whether the compound promotes ABC1-mediated cholesterol efflux. Such method comprises: (a) assaying the level of cholesterol efflux in a sample of mammalian cells
20 maintained in culture to determine a control level of cholesterol efflux, (b) contacting the mammalian cells with the test compound being screened; (c) assaying the level of cholesterol efflux in a sample of cells after contact with the test compound; and (d) assaying the level of ABC1-dependent cholesterol efflux in a sample of cells after contact with the test compound, thereby determining whether the test compound promotes ABC1-mediated cholesterol efflux
25 from cells in culture.

The level of cholesterol efflux in a sample of cultured cells can be determined using methods known in the art and described herein (see Example 1). Any mammalian cells that can be maintained in culture can be used to measure cholesterol efflux. The cells can be derived from primary cultures or from immortalized cell lines. For convenience, cells can be
30 immortalized by transfecting them with amphotropic retroviruses containing vectors with inserts of human papillomavirus 16, oncogenes E6 and E7, and a selectable marker gene, as described in Example 1. Preferably, the cultured cells are fibroblast, macrophage, hepatic, or intestinal cells. More preferably, the cultured cells are RAW 264.7 cells.

The level of cholesterol efflux in a sample of cells that has not been contacted with the test compound is measured to obtain a control level of cholesterol efflux. In addition, the level of cholesterol efflux in a sample of mammalian cells that has been contacted with the test compound is measured to determine the amount of cholesterol efflux affected by the test compound. Also, the level of ABC1-mediated cholesterol efflux in a sample of mammalian cells that has been contacted with the test compound is measured to determine the amount of ABC1-mediated cholesterol efflux affected by the test compound. Preferably, the cells are contacted with the test compound for about 8-24 hours before cholesterol efflux or ABC1-mediated cholesterol efflux is assayed. The level of ABC1-mediated cholesterol efflux can be assayed using an anti-ABC1 antibody that, upon binding, inhibits the activity of ABC1. Alternatively, the level of ABC1-mediated cholesterol efflux can be assayed using an anti-sense ABC1 polynucleotide that inhibits the expression of ABC1. For example, the level of ABC1-mediated cholesterol efflux can be assayed using the anti-sense ABC1 polynucleotide comprising SEQ ID NO: 57 (see Example 7). The cells should be contacted with the anti-ABC1 antibody or anti-sense ABC1 polynucleotide at the same time and for the same duration that it is contacted with the test compound.

If the level of control cholesterol efflux is the same or about the same as the level of cholesterol efflux found in cells contacted with the test compound, the compound does not promote cholesterol efflux. An increase in the level of cholesterol efflux found in cells contacted with test compound over the control level of cholesterol efflux indicates the amount of cholesterol efflux promoted by the test compound. The difference between the cholesterol efflux found in cells contacted with test compound alone and the cholesterol efflux found in cells contacted with test compound and anti-ABC1 antibody or anti-sense ABC1 polynucleotide indicates the amount of cholesterol efflux mediated through ABC1. For example, if control level of cholesterol efflux is 1.0 and the level of cholesterol efflux found in cells contacted with test compound is 1.1, the test compound promotes cholesterol efflux by 10%. If the cholesterol efflux found in cells contacted with test compound and anti-ABC1 antibody or anti-sense ABC1 polynucleotide is 1.0, the increase in cholesterol efflux caused by the test compound is entirely ABC1-mediated.

Methods for Detecting Susceptibility to Coronary Heart Disease

The present invention also relates to methods for detecting the comparative level of ABC1 gene or protein expression in a mammalian subject, including a human subject. Given the

role of ABC1 in cholesterol efflux, the determination of a decreased level of ABC1 gene or protein expression in a mammalian subject relative to a pre-determined standard level of ABC1 gene or protein expression can be used to indicate a susceptibility to coronary heart disease in the subject. Accordingly, the present invention provides a method for detecting the comparative
5 level of ABC1 gene expression in a mammalian subject comprising the steps of: (a) obtaining a test cell sample from the mammalian subject; (b) assaying the level of ABC1 mRNA expression in the test cell sample; and (c) comparing the level of ABC1 mRNA expression in the test cell sample with a pre-determined standard level of ABC1 mRNA expression, thereby detecting the comparative level of ABC1 gene expression in the mammalian subject.

10 A test cell sample is first obtained from a mammalian subject, including a human subject. The test cell sample can be a blood sample, wherein the monocyte population has been enriched. Monocytes can be enriched using well-known cell separation procedures based on, for example, cell size, cell density or cell affinity. Next, the level of ABC1 mRNA expression in the test cell sample is assayed. The level of ABC1 mRNA expression can be assayed using any of the well-
15 known methods for mRNA preparation and detection, including the methods described herein at Examples 2 and 9. For example, the concentration of ABC1 mRNA can be determined by reverse transcription polymerase chain reaction, northern blot analysis, or RNase protection assay. The ABC1 mRNA concentration should be standardized to the concentration of total mRNA found in the test cell sample. Finally, the ABC1 expression in the test cell sample is
20 compared with a pre-determined standard level of ABC1 mRNA expression. A pre-determined standard level of ABC1 mRNA expression can be obtained by determining the average concentrations and distribution of ABC1 mRNA found in cell samples taken from a representative population of mammalian subjects, wherein the mammalian subjects are the same species as the subject from which the test cell sample was obtained, and wherein the mammalian
25 subjects do not have coronary heart disease, Tangier disease, or other disease associated with low HDL-cholesterol and are considered to have cholesterol efflux activity within a normal range (i.e., as indicated by an HDL-cholesterol level within a normal range). The determination of a decreased level of ABC1 mRNA expression in the test cell sample of a mammalian subject relative to the pre-determined standard level of ABC1 mRNA expression can be used to indicate
30 a susceptibility to coronary heart disease in the mammalian subject.

Likewise, the detection of a decreased level of ABC1 protein can be used to indicate decreased capacity for cholesterol efflux and a susceptibility to coronary heart disease. Accordingly, another embodiment of the present invention provides a method for detecting the

comparative level of ABC1 protein in a mammalian subject. Such method comprises the steps of: (a) obtaining a test cell sample from the mammalian subject; (b) assaying the amount of ABC1 protein in the test cell sample; and (c) comparing the amount of ABC1 protein in the test cell sample with a pre-determined standard amount of ABC1 protein, thereby detecting the

5 comparative level of ABC1 protein in the mammalian subject.

The amount of ABC1 protein can be assayed using any of the well-known methods of measuring protein. Preferably, the amount of ABC1 protein is measured using an immunoassay.

In one embodiment, the amount of ABC1 protein is determined by (a) contacting the cell sample with a population of anti-ABC1 antibodies and (b) detecting the anti-ABC1 antibodies associated
10 with the cell sample. For example, the ABC1 protein can be contacted with an antiserum raised against a synthetic peptide corresponding to KNQTVVDAVLTSFLQDEKVKES located at the C-terminus, as described in Example 11. The anti-ABC1 antibodies can be detected using several methods known in the art, including, for example, western blotting, immunoprecipitation, and FACS, wherein the detection can be accomplished using radioactive, colorimetric, or
15 fluorescent labeling. One preferred method for measuring the amount of ABC1 protein in a cell sample is immunoprecipitation, wherein biotinylated ABC1 proteins are contacted with anti-ABC1 antibody and the bound anti-ABC1 antibody is detected using streptavidin horse radish peroxidase.

The amount ABC1 protein in the test cell sample is compared with a pre-determined
20 standard amount of ABC1 protein. A pre-determined standard amount of ABC1 protein can be obtained by determining the average concentration of ABC1 protein found in cell samples taken from a population of mammalian subjects, wherein the mammalian subjects are the same species as the subject from which the test cell sample was obtained, and wherein the mammalian subjects do not have coronary heart disease, Tangier disease, or other disease associated with low HDL-
25 cholesterol and are considered to have cholesterol efflux activity within a normal range (i.e., as indicated by an HDL-cholesterol level within a normal range).

ABC1 Antibodies

As used herein, the term "antibody" (Ab) or "monoclonal antibody" (Mab) is meant to
30 include intact molecules as well as antibody fragments (such as, for example, Fab and F(ab')₂ fragments) which are capable of specifically binding to protein. Fab and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody (Wahl et al., *J. Nucl. Med.*, 24:316-325

(1983)). Thus, these fragments are preferred, as well as the products of a FAB or other immunoglobulin expression library. Moreover, antibodies of the present invention include chimeric, single chain, and humanized antibodies.

Additional embodiments include chimeric antibodies, e.g., humanized versions of murine
5 monoclonal antibodies. Such humanized antibodies may be prepared by known techniques, and offer the advantage of reduced immunogenicity when the antibodies are administered to humans.

In one embodiment, a humanized monoclonal antibody comprises the variable region of a murine antibody (or just the antigen binding site thereof) and a constant region derived from a human antibody. Alternatively, a humanized antibody fragment may comprise the antigen
10 binding site of a murine monoclonal antibody and a variable region fragment (lacking the antigen-binding site) derived from a human antibody. Procedures for the production of chimeric and further engineered monoclonal antibodies include those described in Riechmann et al. (*Nature* 332:323, 1988), Liu et al. (*PNAS* 84:3439, 1987), Larrick et al. (*Bio/Technology* 7:934, 1989), and Winter and Harris (*TIPS* 14:139, May, 1993).

One method for producing an antibody comprises immunizing a non-human animal, such as a transgenic mouse, with a polypeptide translated from a polynucleotide comprising SEQ ID NO: 1, a polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, or a polynucleotide comprising a nucleotide sequence that has at least 90% identity with a polynucleotide comprising SEQ ID NO: 1, whereby antibodies directed against the polypeptide translated from the
20 described polynucleotides are generated in said animal. Procedures have been developed for generating human antibodies in non-human animals. The antibodies may be partially human, or preferably completely human. Non-human animals (such as transgenic mice) into which genetic material encoding one or more human immunoglobulin chains has been introduced may be employed. Such transgenic mice may be genetically altered in a variety of ways. The genetic
25 manipulation may result in human immunoglobulin polypeptide chains replacing endogenous immunoglobulin chains in at least some (preferably virtually all) antibodies produced by the animal upon immunization. Antibodies produced by immunizing transgenic animals with a polypeptide translated from any of the described polynucleotides are provided herein.

Mice in which one or more endogenous immunoglobulin genes are inactivated by
30 various means have been prepared. Human immunoglobulin genes have been introduced into the mice to replace the inactivated mouse genes. Antibodies produced in the animals incorporate human immunoglobulin polypeptide chains encoded by the human genetic material introduced into the animal. Examples of techniques for production and use of such transgenic

animals are described in U.S. Patents 5,814,318, 5,569,825, and 5,545,806, which are incorporated by reference herein.

Monoclonal antibodies may be produced by conventional procedures, e.g., by immortalizing spleen cells harvested from the transgenic animal after completion of the immunization schedule. The spleen cells may be fused with myeloma cells to produce hybridomas, by conventional procedures.

A method for producing a hybridoma cell line comprises immunizing such a transgenic animal with an immunogen comprising at least seven contiguous amino acid residues of a polypeptide translated from one of the described polynucleotides; harvesting spleen cells from the immunized animal; fusing the harvested spleen cells to a myeloma cell line, thereby generating hybridoma cells; and identifying a hybridoma cell line that produces a monoclonal antibody that binds a polypeptide translated from one of the described polynucleotides. Such hybridoma cell lines, and monoclonal antibodies produced therefrom, are encompassed by the present invention. Monoclonal antibodies secreted by the hybridoma cell line are purified by conventional techniques.

The antibodies, upon specific binding to an ABC1 polypeptide, may inhibit the activity of the ABC1 polypeptide. Preferably, the antibody, upon binding, inhibits the cholesterol transport activity of the ABC1 polypeptide. Such antibodies can be made by immunizing a non-human animal with a polypeptide corresponding to the region essential for cholesterol transport. The antibody can be tested to determine whether it inhibits cholesterol efflux using any of the described cholesterol efflux assays. Such inactivating antibodies can be employed in an *in vitro* assay, such as any of the cholesterol efflux assays described herein, to determine whether a test compound promotes ABC1-mediated cholesterol efflux. The inactivating antibodies can also be used in *in vitro* assays to detect the comparative level of ABC1 protein in the cells of a mammalian subject. The inactivating antibodies are also useful in kits suitable for screening a compound to determine whether the compound modulates ABC1-dependent cholesterol efflux.

Kits for Identifying Therapeutic Compounds

The present invention also includes a kit suitable for screening a compound to determine the ABC1 expression modulating activity of a compound. The kit includes, in an amount sufficient to perform at least one assay, a recombinant reporter construct comprising a reporter cDNA operatively linked to an expression modulating portion of the mammalian ABC1 gene, as a separately packaged reagent. Instructions for use of the packaged reagent(s) are also typically

included. The expression modulating portion of the mammalian ABC1 gene comprises the 5' flanking sequence. In one preferred embodiment, the expression modulating portion of the mammalian ABC1 gene comprises SEQ ID NO: 3. In other preferred embodiments, the expression modulating portion of the mammalian ABC1 gene comprises nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, 1394-1643, or 1394-1532 of SEQ ID NO: 3. The reporter cDNA can be any suitable reporter gene, including In a particularly preferred embodiment, the recombinant reporter construct is pAPR1. In another embodiment, the kit further comprises means for detecting the reporter protein. Thus, the kit comprises reagents, such as buffers and substrates, used for reporter protein detection.

As used herein, the term "package" refers to a solid matrix or material such as glass, plastic (e.g., polyethylene, polypropylene or polycarbonate), paper, foil and the like capable of holding within fixed limits a recombinant vector of the present invention. Thus, for example, a package can be a glass vial used to contain milligram quantities of a contemplated vector.

"Instructions for use" typically include a tangible expression describing the reagent concentration or at least one assay method parameter such as the relative amounts of reagent and sample to be admixed, maintenance time periods for reagent/ sample admixtures, temperature, buffer conditions and the like.

In addition, the present invention also includes a kit suitable for screening a compound to determine whether the compound modulates ABC1-dependent cholesterol efflux. In one embodiment, the kit includes, in an amount sufficient to perform at least one assay, an inactivating anti-ABC1 antibody, as a separately packaged reagent. Instructions for use of the packaged reagent(s) are also typically included.

In another embodiment, the kit includes an antisense ABC1 oligonucleotide in an amount sufficient for at least one assay and instructions for use. Preferably, the antisense ABC1 oligonucleotide comprises SEQ ID NO: 53.

The following examples further illustrate the present invention but should not be construed to limit the present invention in any way.

30

EXAMPLE 1

This example demonstrates that patients with Tangier Disease (TD) have an absence of apo A-I-mediated lipid efflux.

Cell Cultures: Human fibroblasts were obtained from skin explants from two normal

subjects (NL1) and three unrelated patients with Tangier disease (TD). TD1 cells were obtained from a 53 year-old female with extremely low plasma HDL cholesterol and apo A-I levels and clinical symptoms typical of Tangier disease. TD2 cells were obtained from a 56 year-old male with clinical, morphological, and biochemical features of Tangier disease, including very low
5 levels of plasma HDL cholesterol and apoA-I (Francis, et al., *J. Clin. Invest.*, 96, 78-87 (1995)). TD3 cells were obtained from an 18 year-old male with Tangier disease who presented with orange tonsil remnants, asymmetrical motor neuropathy, plasma HDL cholesterol of 5 mg/dL, and LDL cholesterol of 16mg/dL (Lawn et al., *J. Clin. Invest.*, 104, R25-R31 (1999)). The normal cells and TD subject cells were immortalized as described in Oram et al., *J. Lipid Res.*,
10 40: 1769-1781 (1999). Briefly, the cells were transfected with amphotropic retroviruses containing vectors with inserts of human papillomavirus 16 oncogenes E6 and E7 and a neomycin resistance selectable marker. Control cells were infected with vector alone (mock-infected). Pooled cell populations were selected in the presence of G418 for two passages, after which G418 was excluded from the medium. Fibroblasts were used between the fifth and
15 sixteenth passage (primary) or sixth and fourteenth passage (immortalized). The immortalized normal and TD cells were seeded into 16-mm wells or 35-mm dishes and grown to confluence in Dulbecco's modified Eagle's medium (DMEM) plus 10% fetal bovine serum (FBS) before experimental use. RAW 264.7 mouse monocytic cells (American Type Culture Collection, Rockville, MD) were also maintained in DMEM containing 10% FBS.

20 Assay to Measure Lipid Efflux: Apo AI-mediated efflux of cholesterol and phospholipid was assayed according to the method described in Francis, et al., *J. Clin. Invest.*, 96: 78-87 (1995). The cultured skin fibroblasts from normal and TD subjects were labeled by growth to confluence in the presence of 0.2-0.5 $\mu\text{Ci/ml}$ [^3H]cholesterol (40-60 Ci/mmol, Amersham Corp., Arlington Heights, IL). The radioactive cholesterol was added to serum-containing growth
25 medium when the cells were 60-80% confluent. After 3 days, the cells were washed twice with PBS/BSA and simultaneously growth-arrested and cholesterol-loaded to maximize apolipoprotein-mediated lipid efflux. This was achieved by incubating the cells for 48 hours in serum-free DMEM plus 2 mg/ml fatty acid-free bovine serum albumin (DMEM/BSA) (Sigma Chemical Co., St. Louis, MO) and 30 $\mu\text{g/ml}$ non-radioactive cholesterol. RAW 264.7 cells were
30 cholesterol-loaded through the scavenger receptor by 24-hour incubation with acetylated LDL as described in Smith et al., *J. Biol. Chem.*, 271:30647-30655 (1996). Briefly, RAW 264.7 cells in 24-well dishes were cholesterol-loaded and labeled overnight in 0.5 ml of DMEM supplemented with 50 $\mu\text{l/ml}$ 1M glucose, 10 $\mu\text{l/ml}$ 200 mM glutamine and 2% BSA and with 50 $\mu\text{g/ml}$

acetylated low density lipoprotein (AcLDL) and [³H]-cholesterol which had been pre-incubated for 30 minutes at 37°C with AcLDL to yield a final concentration of 0.33 µCi/ml [³H]-cholesterol. Cells were subsequently washed five times with PBS containing 1% BSA and incubated overnight (16-18 hours) in DMEM/BSA to allow for equilibrium of cholesterol pools.

5 After equilibrium of cholesterol pools, cells were rinsed four times with PBS/BSA and incubated for one hour at 37°C with DMEM/BSA before the efflux incubations. Efflux medium (DMEM/BSA) containing either albumin alone (control), albumin plus HDL (40µg protein/ml), or albumin plus apo A-I (10 µg/ml, Biodesign International, Kennebunk, ME) was added and the cells were incubated for 4, 24, or 48 hours. Phospholipids were labeled by including 10 µCi/ml
10 [³H]choline (75-85 Ci/mmol, Amersham Corp.) in the DMEM/BSA overnight equilibrium medium. The radioactivity found in the culture medium was measured by scintillation counting after a 15-minute centrifugation at 12,000g. The radioactivity in the cells was measured by scintillation counting after solubilization in 0.5 ml of 0.2M NaOH (Smith et al., *J. Biol. Chem.*, 271:30647-30655 (1996)) or extraction in hexane:isopropanol (3:2 v/v) as described in Francis,
15 et al., *J. Clin. Invest.*, 96, 78-87 (1995). Cells containing labeled phospholipids were extracted with 1 ml of isopropanol for 1 hour and then with hexane:isopropanol as described above. The efflux of cholesterol or phospholipid was expressed as the percentage of tritiated lipid counts in the medium over the total tritiated lipid counts recovered from the cells and medium (cpm medium / cpm (medium + lysate) x 100).

20 As shown in Figure 1A and C, the addition of HDL or apo A-I results in the removal of cholesterol from cholesterol-laden fibroblasts obtained from normal subjects. However, in TD cells, the ability of HDL to remove cholesterol is slightly diminished and the ability of apo A-I to remove cholesterol is completely absent. Figure 1 shows that normal and TD fibroblast cells release about 3-4% of the cellular [³H]-cholesterol into the medium during 48-hour incubation
25 with albumin. Addition of HDL to the albumin medium increased the efflux of [³H]-cholesterol from both normal and TD fibroblasts, although to a lesser extent with TD cells (Figure 1B, D). Addition of apo A-I promoted the efflux of [³H]-cholesterol from normal fibroblasts (Figure 1A, C), but had little or no effect on [³H]-cholesterol efflux from TD fibroblasts (Figure 1B, D).

30

EXAMPLE 2

This example demonstrates that 175 genes show at least 2.5-fold decreased expression and 375 genes show at least 2.5-fold increased expression in TD cells compared with normal cells. The differential gene expression was determined using gene-expression microarray (GEM)

analysis of cDNAs from normal individuals (non-TD) and from patients with TD.

Cell Cultures: The immortalized cell cultures obtained from normal individuals and TD patients described in Example 1 were used. Confluent cultures were maintained in DMEM/BSA and supplemented for 24 hours with 1mM 8-Bromo cyclic adenosine monophosphate (8-Br-cAMP, Sigma Chemical Co., St. Louis, MO).

mRNA Extraction and cDNA Synthesis: mRNA from both normal and TD fibroblast cells was prepared from total RNA extracted from cells with Trizol (Life Technologies Inc., Bethesda, MD, Cat. # 15596-026). The mRNA was isolated using the Oligotex mRNA kit (Qiagen Inc., Valencia, CA, Cat. # 70022) according to vendor's protocols. The mRNA was reverse transcribed using Cy3 or Cy5 fluorescent dye to create fluorescently labeled cDNA according to the method described in DeRisi et al., *Science*, 24:680-686 (1997). The resultant cDNA from TD cells was labeled with Cy3 fluorescent dye and cDNA from normal cells was labeled with Cy5 fluorescent dye (Incyte Genomics, Palo Alto, CA).

Microarray Analysis: To analyze differential gene expression in cells from individuals with TD and normal individuals, Cy3 and Cy5 fluorescent labeled cDNA samples prepared as described above were hybridized to a set of Gene Album microarrays (GEMs) on microscope slides (Incyte Genomics, Palo Alto, CA). Each of six slides contained about 9,800 human cDNA samples plus 200 control samples, resulting in a microarray of 58,800 partial cDNAs. Therefore, allowing for estimates of redundancy, approximately 30-50% of expressed human genes were represented. The hybridization of Cy3-labeled cDNA prepared from TD1 cells and Cy5-labeled cDNA from normal cells allowed comparison of the relative RNA content of TD cells versus normal cells for the expressed genes. In addition, Cy3-labeled cDNA prepared from TD2 cells and Cy5-labeled cDNA from normal cells were hybridized to the same set of microarrays to examine the variation of gene expression between different TD patients.

Results: Data were analyzed using GemTools software (Incyte Genomics, Palo Alto, CA) and expressed as ratios of TD cell to normal cell mRNA. The results indicated that the majority of genes are comparably expressed in TD1 and normal cells. As shown in Figure 2 (in the section above and to the left of the diagonal) only 175 genes were more than 2.5-fold underexpressed in TD1 cells compared with normal cells, whereas 375 genes were more than 2.5-fold overexpressed in TD1 cells compared with normal cells (below and to the right of the diagonal). Genes more highly expressed in the TD cells could include those that are differentially regulated as a consequence of the Tangier mutation, either as a compensatory response or as a contributor to the disease pathology. Among the genes that could contribute to

the observed phenotype of TD and are more highly expressed in TD cells include interferon- β (IFN- β), macrophage inflammatory protein-2 α , granulocyte chemotactic protein-2, IL-11, prostaglandin endoperoxide synthase-2 (COX-2), thrombospondin, and monocyte chemotactic proteins 1, 3, and 4 (Lawn et al., *J. Clin. Invest.*, 104:R25-R31 (1999)).

5 No single RNA that was expressed in the normal fibroblasts was found completely absent in either the TD1 or TD2 cells. Also, comparison of the differentially expressed genes in TD1 and TD2 revealed very little variation between the individual TD patients. For instance, of the most highly down-regulated genes in TD2 cells, 92% were also under-expressed in TD1 cells compared with normal cells. One of the genes more than 2.5-fold under-expressed in TD1 or
10 TD2 versus normal cells was the gene for ABC1 protein. The ABC1 gene was pursued due to the ascribed functions of some of its homologues and also because the gene was localized to the approximate chromosome region reported as the TD gene region.

EXAMPLE 3

15 This example demonstrates that the ABC1 gene is localized to the human chromosome 9q31.

Previous genetic linkage analysis mapped the TD gene to the 7-cM region of human chromosome 9q31 (Rust et al., *Nat. Genet.*, 20, 96-98 (1998)). In addition, *in situ* hybridization analyses revealed that the ABC1 gene was localized to the broader chromosomal interval 9q22 -
20 9q31 (Luciani et al., *Genomics*, 21, 150-159 (1994)). Using PCR methods with the GeneBridge 4 panel of human/hampster radiation hybrids (Research Genetics, Inc., Huntsville, AL), human ABC1 was determined to be located between the markers WI-14706 and WI-4062, corresponding to the 7-cM region of human chromosome 9q31. DNA from 93 human/hampster hybrid cell lines was amplified by PCR using human ABC1-specific primers LF:
25 CCTCTCATTACACAAAACCAGAC (SEQ ID NO: 11) and LR:
GCTTTCTTTCACTTCTCATCCTG (SEQ ID NO: 12). Each line was scored as positive or negative for the human ABC1 amplification product and the mapping of ABC1 derived from analysis of this data was accomplished using the Whitehead Institute/MIT Center for Genome Research software, accessed via the internet ([http://carbon.wi.mit.edu:8000/cgi-](http://carbon.wi.mit.edu:8000/cgi-bin/contig/rhmapper.pl)
30 [bin/contig/rhmapper.pl](http://carbon.wi.mit.edu:8000/cgi-bin/contig/rhmapper.pl)). These results were further confirmed by southern blot hybridization to human genomic/yeast artificial chromosome clones (Research Genetics, Inc.) from the equivalent interval. In addition, public database searching (GeneMap '98; National Center for Biotechnology Information) and radiation hybrid mapping eliminated the other significantly

underexpressed genes in the microarray data from the location in the reported genetic interval. These complementary data demonstrate that the ABC1 gene is located on human chromosome 9q31 and further indicate that the ABC1 gene is associated with Tangier disease.

EXAMPLE 4

This example shows the determination of the nucleotide sequence of the wildtype ABC1 gene, including the flanking regions and the entire coding region.

DNA sequencing was performed using an ABI Prism 310 Genetic Analyzer or by Davis Sequencing (Davis, CA). Both strands were sequenced throughout. The sequence of the open reading frame of the ABC1 gene from a normal subject was determined from a full-length cDNA clone obtained from an expression plasmid library constructed from normal fibroblast RNA. To construct the plasmid library, cDNA was synthesized according to the Stratagene kit protocol (Stratagene, La Jolla, CA). Briefly, first strand cDNA was synthesized from mRNA using an oligo-dT primer with an XhoI site and MMLV reverse transcriptase in the presence of 5-methyl dCTP. The second strand was synthesized using RNase H and DNA polymerase I in the presence of unmodified dNTPs. After the cDNA was blunt-ended with pfu DNA polymerase, an EcoRI linker was ligated to the cDNA. The cDNA was then digested with XhoI, creating XhoI ends at the 3' end of the cDNA. The internal XhoI sites were protected from this digestion due to the semi-methylation during the first strand synthesis. The synthesized cDNA was cloned into the HindIII and XhoI sites of the plasmid pCEP4 (Invitrogen Corp., Carlsbad, CA #VO44-50), an expression vector containing the cytomegalovirus promoter/enhancer. A 585 bp ABC1 probe was generated by reverse transcriptase polymerase chain reaction (RT-PCR) using primers based on known ABC1 sequence, which were 5'-TCCTTGGGTTTCAGGGGATTATC (SEQ ID NO: 13) and 5'-CAATGTTTTTGTGGCTTCGGC (SEQ ID NO: 14). Using this ABC1 probe, a clone containing a 10.5 kb insert of human ABC1 cDNA was recovered from the library using the CloneCapture selection kit according to the manufacturer's protocol (CLONTECH Laboratories, Inc., Palo Alto, CA). This clone is shown in Figure 3 as pCEPhABC1. The 10.5 kb ABC1 cDNA insert sequence is shown in SEQ ID NO: 1. Sequence determination confirmed that pCEPhABC1 contains the human ABC1 open reading frame of 6783 nucleotides plus 5' and 3' untranslated regions, having a larger open reading frame than the cDNA sequence reported by Langmann et al. in *Biochem. Biophys. Res. Comm.*, 257, 29-33 (1999) (GenBank Accession No. AJ012376).

EXAMPLE 5

This example demonstrates the sequence differences between the wildtype ABC1 gene and the TD1, TD2, and TD3 gene sequences.

cDNA Synthesis of TD1, TD2, and TD3: cDNA was prepared from TD1, TD2, and TD3 cells by reverse transcription polymerase chain reaction (RT-PCR) using the Superscript Choice cDNA system and the Advantage cDNA polymerase mix following the manufacturer's protocol (CLONTECH, Palo Alto, CA; Cat. #8417-1) using two sets of primer pairs designed from the normal human ABC1 gene sequence, designated: (1) sacIhabcf, 5'-AGTCGAGCTCCAAACATGTCAGCTGTTACTGGAAGTGGCC (SEQ ID NO: 15); habcr3851, 5'-TCTCTGGATTCTGGGTCTATGTCAG (SEQ ID NO: 16) and (2) habcf3585, 5'-GGGAGCCTTTGTGGAAGTCTTTC (SEQ ID NO: 17); habcrsall, 5'-ACTGGTCGACCATTGAATTGCATTGCATTGAATAGTATCAG (SEQ ID NO: 18). Amplification of 0.2-0.5 µg polyA+ RNA with these primers at a final concentration of 0.4µM generated two overlapping templates of approximately 3.5 kb. The templates were gel-purified using the QIAEX II system (QIAGEN, Inc., Valencia, CA; Cat. #20021) and adjusted to a concentration of 100ng/µl.

Sequencing of TD1, TD2, and TD3 cDNA: Eight µl of each template generated as described above was sequenced in a reaction with individual sequencing primers designed on the basis of wildtype ABC1 sequence at a final concentration of 0.5µM. The primers were as follows: 1F: 5'-TTTCCTGGTGGACAATGAA (SEQ ID NO: 19), 2F: 5'-AGTGACATGCGACAGGAG (SEQ ID NO: 20); 3F: 5'-GATCTGGAAGGCATGTGG (SEQ ID NO: 21); 4F: 5'-CCAGGCAGCATTGAGCTG (SEQ ID NO: 22); 5F: 5'-GGCCTGGACAACAGCATA (SEQ ID NO: 23); 6F: 5'-GGACAACCTGTTTGAGAGT (SEQ ID NO: 24); 7F: 5'-AAGACGACCACCATGTCA (SEQ ID NO: 25); 8F: 5'-ATATGGGAGCTGCTGCTG (SEQ ID NO: 26); 9F: 5'-GGGCATGAGCTGACCTATGTGCTG (SEQ ID NO: 27); 10F: 5'-AAGAGACTGCTAATTGCC (SEQ ID NO: 28); 11F: 5'-AGCGACAAAATCAAGAAG (SEQ ID NO: 29); 12F: 5'-TGGCATGCAATCAGCTCT (SEQ ID NO: 30); 13F: 5'-TCCTCCACCAATCTGCCT (SEQ ID NO: 31); 14F: 5'-TTCTTCCTCATTACTGTT (SEQ ID NO: 32); 15F: 5'-GATGCCATCACAGAGCTG (SEQ ID NO: 33); 16F: 5'-AGTGTCCAGCATCTAAA (SEQ ID NO: 34); 1R: 5'-CAAAGTTCACAAATACTT (SEQ ID NO: 35); 2R: 5'-CTTAGGGCACAATTCCACA (SEQ ID NO: 36); 3R: 5'-TGAAAGTTGATGATTTTC (SEQ ID NO: 37); 4R: 5'-TTTTTCACCATGTGCGATGA (SEQ ID NO: 38)

NO: 38); 5R: 5'- CTCCACTGATGAACTGC (SEQ ID NO: 39); 6R: 5'-
 GTTCTTCATTTGTTTGA (SEQ ID NO: 40); 7R: 5'- AGGGCGTGTCTGGGATTG (SEQ ID
 NO: 41); 8R: 5'- CAGAATCATTTGGATCAG (SEQ ID NO: 42); 9R: 5'-
 CATCAGAACTGCTCTGAG (SEQ ID NO: 43); 10R: 5'- AGCTGGCTTGTTTTGCTTT SEQ
 5 ID NO: 44), 11R: 5'- TGGACACGCCCAGCTTCA (SEQ ID NO: 45), 12R: 5'-
 CCTGCCATGCCACACACA (SEQ ID NO: 46), 13R: 5'- CTCATCACCCGCAGAAAG (SEQ
 ID NO: 47), 14R: 5'- CAACTCCATGAAGCGAG (SEQ ID NO: 48), 15R: 5'-
 TCCAGATAATGCGGGAAA (SEQ ID NO: 49), 16R: 5'- TCAGGATTGGCTTCAGGA (SEQ
 ID NO: 50), UTR1R: 5'- AAGTTTGAGCTGGATTCTTG (SEQ ID NO: 51).

10 Results: The nucleotide numbering follows the numbering found in Lawn et al. (1999).
 Patient TD1 retained the full open reading frame, with 2 substantial differences from the wild-
 type sequence (SEQ ID NO: 8). One of these is an A to G substitution, resulting in a change
 from a glutamine to arginine residue at position 537 of the 2201 amino acid sequence, as
 published by Lawn et al. (1999). The location of this residue is within the NH₂-terminal
 15 hydrophilic domain, near the first predicted transmembrane domain. Patient TD2 also retained
 the open reading frame with an arginine to tryptophan substitution at residue 527 (SEQ ID NO:
 10). Thus, both TD1 and TD2 contain a substitution altering the charge of an amino acid in the
 same region of the protein. TD3 DNA contains a 14 nucleotide insertion in its ABC1 cDNA
 following nucleotide 5697 in one allele and a 138 bp insertion after nucleotide 5062 in the other
 20 allele.

Genomic sequencing of the TD1, TD2, and TD3 DNAs confirmed the changes found in
 the respective cDNAs. The genomic sequence was generated by PCR amplification of a 156 bp
 region of genomic DNA isolated from fibroblasts that contained the mutations found in the
 cDNA from TD1 and TD2. The genomic sequencing also indicated that patient TD1 was
 25 homozygous for the glutamine to arginine substitution. Genomic DNA analysis showed that
 TD2 was a compound heterozygote with one allele containing the detected substitution and the
 second allele (which failed to produce detectable mRNA) containing an undetermined defect.
 Neither of the substitution mutations was found in more than 80 alleles of genomic DNA of
 non-TD individuals. TD3 insertions were identified by sequence analysis and confirmed by RT-
 30 PCR using primers surrounding the insertion points. The 14-bp insertion following nucleotide
 5697 causes a frameshift, resulting in the replacement of the wild-type amino acid sequence from
 a location before the second ATP binding domain, up to the point of a premature protein
 termination. The 138 bp insertion following nucleotide 5062 in the other allele contains an

inframe stop codon.

EXAMPLE 6

This example demonstrates that inhibitors of ABC1 transport activity also inhibit apo A-I-mediated cholesterol efflux from fibroblast cells.

5 To test whether inhibition of ABC1 could affect the process of apolipoprotein-mediated cholesterol efflux, two compounds reported to be ABC1 inhibitors were tested in assays which monitor apolipoprotein mediated cholesterol efflux. The compounds 4,4-diisothiocyanostilbene-2,2'-disulfonic acid (DIDS) and sulphobromophthaleine (BSP) were reported to inhibit anion transport activities of ABC1 in a dose-dependent fashion (Becq et al., *J. Biol. Chem.*, 272:2695-10 2699 (1997); Hamon et al., *Blood*, 90:2911-2915 (1997)). The apolipoprotein-mediated cholesterol efflux assays were performed as described in Example 1 with the noted changes. Cholesterol-loaded and [³H]cholesterol-labeled normal fibroblasts (n=3) were incubated for 6 hours with or without 5µg/ml apo A-I and either 0, 0.2mM, or 0.4mM DIDS. In addition, cholesterol-loaded and [³H]cholesterol-labeled normal fibroblasts (n=3) were incubated for 15 hours with or without 5µg/ml apo A-I and either 0, 0.2mM, or 0.4mM BSP. [³H]cholesterol efflux was measured by scintillation counting as described in Example 1 and calculated as the percentage of total radiolabeled cholesterol appearing in the medium. The results are shown in Figure 5 as the mean ± SD (n=3) of efflux in the presence of apo A-I after subtraction of values for apo A-I-free medium. Figure 5 shows that both DIDS and BSP inhibit the 6-hour efflux of 20 tritiated cholesterol mediated by apolipoprotein A-I. In addition, similar inhibition was observed with the efflux of tritiated phosphatidyl choline using DIDS and BSP (data not shown). The results of these tests mimic the efflux defect in fibroblasts derived from patients with TD, described in Example 1.

EXAMPLE 7

25 This example demonstrates that antisense inhibition of ABC1 mRNA expression inhibits apo A-I-mediated cholesterol efflux from fibroblast cells.

Normal skin fibroblasts were labeled with [³H]cholesterol as described in Example 1. The cells were then loaded with oligonucleotide by scraping in the presence of either 30µM control Morpholino oligonucleotide (5'-CCTCTTACCTCAGTTACAATTTATA-3' 30 corresponding to the antisense complement of a β-globin thalassemic mRNA; SEQ ID NO: 52) or 30µM ABC1 antisense Morpholino oligonucleotide (5'-CATGTTGTTTCATAGGGTGGGTAGCTC-3'; SEQ ID NO: 53) and reseeding on new dishes. Control cells were mock-loaded after [³H]cholesterol-labeling by scraping and reseeding in the

absence of oligonucleotide. Apo A-I-mediated efflux was measured after 12 hours by scintillation counting as the percentage of total radiolabeled cholesterol appearing in the medium. The results are shown in Figure 6 as the mean \pm SEM of three separate experiments, normalized to the value for apo A-I-specific efflux in the absence of oligonucleotide in each experiment. As shown in Figure 6, antisense oligonucleotides directed against ABC1 mRNA caused a 50% reduction in cholesterol efflux from normal fibroblasts compared with control antisense oligonucleotide (β -globin antisense oligonucleotide).

EXAMPLE 8

This example demonstrates that over expression of the human ABC1 gene results in an increase in apo A-I-mediated cholesterol efflux from monocyte cells.

Stable Transfection of RAW 264.7 Cells: Mouse monocytic RAW 264.7 cells were stably transfected with the pCEPhABC1 expression plasmid for human ABC1. Construction of the pCEPhABC1 plasmid containing the open reading frame of human ABC1 is described in Example 4. Approximately 1×10^6 RAW 264.7 cells were transfected for 5 hours with 2 μ g of pCEPhABC1 DNA and 12 μ l Geneporter transfection reagent (Gene Therapy Systems, Inc., San Diego, CA; Cat. #T201007) in 0.8ml serum-free DMEM. Two days later, cells were split at ratios ranging from 1:2 - 1:50 and selection applied by adding 150 μ g/ml hygromycin to the culture medium. After two weeks, hygromycin-resistant colonies were picked and expanded.

Apo A-I-mediated Cholesterol Efflux Assay: Parental RAW 264.7 cells and three clonal lines (L3, L5, and L6) stably expressing human ABC1 were grown to confluence. The cells were cholesterol-loaded and labeled by incubation for 24 hours with 0.5 μ Ci/ml [3 H]cholesterol and 50 μ g/ml acetylated LDL as described in Example 1. After equilibrium of cholesterol pools by an overnight incubation in DMEM/BSA, cells were washed and the efflux medium was added as described in Example 1. Apo A-I-mediated cholesterol efflux was measured as previously described by scintillation counting of the tritiated cholesterol in the cell medium, expressed as a percentage of the total counts recovered from the cells and medium. The results are presented as the mean \pm SEM of three separate experiments normalized to the value for apo A-I-specific efflux from parental RAW 264.7 cells within each experiment. Figure 7 shows the apo A-I-mediated cholesterol efflux from parental RAW 264.7 cells and L3, L5, and L6 transfected cell lines. As can be seen, transfection with the ABC1 expression vector results in a 4-fold (L6) to 8-fold (L3 and L5) increase in apo A-I-mediated cholesterol efflux. These results indicate that overexpression of the ABC1 gene can substantially increase the amount of cholesterol efflux

from macrophage cells.

EXAMPLE 9

This example demonstrates that ABC1 mRNA expression is regulated by cellular
5 conditions related to cholesterol efflux in normal skin fibroblasts, but not in TD fibroblasts.

To determine whether ABC1 plays a rate-limiting role in cellular sterol efflux, the
synthesis of ABC1 was measured under various cellular conditions related to cholesterol efflux
processes. Specifically, normal fibroblasts and TD fibroblasts were individually exposed to
conditions of excess cAMP, cholesterol, or Apo A-I. Cell cultures of normal skin fibroblasts and
10 TD1 and TD2 fibroblasts were prepared as described in Example 1. The level of ABC1 mRNA
was measured by RT-PCR.

Cell cultures: Immortalized cell cultures of normal skin fibroblasts and TD1 and TD2
fibroblasts were prepared as described in Example 1. Cells were grown to subconfluence in
DMEM/10% FBS before replacement with DMEM/BSA and the indicated additive for 24 or 48
15 hours. RNA was prepared as described in Example 2.

RT-PCR: Quantitative PCR was carried out using the GeneAmp 5700 Sequence
Detection System (Perkin-Elmer Applied Biosystems, Foster City, CA). Briefly, 500ng of
DNase-treated mRNA was reverse transcribed using random hexamer primers at 2.5 μ M.
Approximately 5% of this reaction was amplified by PCR using the SYBR green core kit (PE
20 Applied Biosystems, Foster City, CA; Cat. #4304886) and human ABC1 primers LF: 5'-
CCTCTCATTACACAAAAACCAGAC (SEQ ID NO: 11) and LR: 5'-
GCTTTCTTTCACTTCTCATCCTG (SEQ ID NO: 12) to yield an 82 bp fragment
corresponding to nucleotides 7018-7099 of human ABC1. PCR cycle conditions were as
follows: 10 minutes 95°C; followed by 40 cycles of 95°C, 15 seconds; and 60°C, 60 seconds.
25 The mRNA in each sample was quantitated by detecting the increase in fluorescence caused by
SYBR green binding to the double-stranded amplification product generated during each PCR
cycle. All samples were run in triplicate and normalized against β -actin mRNA, amplified in
parallel reactions with primers actinF: 5'-TCACCCACACTGTGCCATCTACGA (SEQ ID NO:
54) and actinB: 5'-CAGCGGAACCGCTCATTGCCAATGG (SEQ ID NO: 55). Standard
30 curves were run for both ABC1 and β -actin on the same PCR plate.

8-Br-cAMP Assay: Normal, TD1, and TD2 fibroblast cells were grown to subconfluence
in DMEM/10% FBS and then treated with 1 mM 8-Br-cAMP in DMEM/BSA for 24 hours.

Cholesterol Assay: Normal, TD1, and TD2 fibroblast cells were grown to subconfluence in DMEM/10% FBS and then treated with 30µg/ml free cholesterol in DMEM/BSA for 48 hours followed by 18-24 hours of equilibrium in DMEM/BSA.

5 Apo A-I Assay: Normal, TD1, and TD2 fibroblast cells were grown to subconfluence in DMEM/10% FBS and then treated with 30µg/ml free cholesterol in DMEM/BSA for 48 hours followed by 18-24 hours of equilibrium in DMEM/BSA plus 10µg/ml apo A-I.

Results: Figure 8 shows that in normal fibroblasts ABC1 mRNA is increased approximately 10-fold by exposure to 8-Br-cAMP and increased approximately 17-fold by exposure to cholesterol in serum-free medium. Subsequent exposure of cholesterol-loaded cells to Apo A-I results in a marked decrease in ABC1 mRNA expression. Although the mechanism has not been demonstrated, previous studies have shown that cholesterol efflux is promoted in the presence of such compounds as cAMP and cholesterol (Hokland et al., *J. Biol. Chem.*, 268:25343-25349 (1993)). The present results indicate that in normal fibroblasts, ABC1 mRNA is induced by these known effectors of the cholesterol efflux pathway and repressed by exposure to an apolipoprotein cholesterol acceptor, demonstrating that the expression of ABC1 is regulated by cellular conditions related to apolipoprotein-mediated cholesterol efflux. In contrast, fibroblast cells from TD patients are not regulated by effectors of cholesterol efflux. First, the cAMP-inducible level of ABC1 mRNA in both TD1 and TD2 cells is only approximately 40% of that in normal cells. Further, exposure of cholesterol-loaded cells to Apo A-I either did not alter ABC1 expression (TD1 cells) or slightly increased ABC1 expression (TD2 cells). These results reflect the defect in Apo-A-I-mediated cholesterol efflux described for TD cells. Interestingly, growth of cells in serum-containing medium suppressed ABC1 message to near the limit of detection (data not shown). This may reflect the fact that functioning of the lipid efflux pathway requires cell quiescence or other cellular states of reduced cholesterol need. In conclusion, conditions that are associated with increased efflux of cellular cholesterol (i.e., cholesterol loading, cAMP treatment, serum deprivation) also result in increased expression of ABC1 mRNA in normal fibroblast cells. Conversely, exposure of cholesterol-loaded normal fibroblast cells to apo A-I reduces ABC1 expression.

30

EXAMPLE 10

This example demonstrates that ligands for the LXR nuclear receptor, such as 20-hydroxycholesterol, and ligands for the RXR receptor, such as 9-*cis* retinoic acid, can increase ABC1 gene expression in mouse RAW 264.7 cells.

LXR nuclear receptors are transcription factors that form obligate heterodimers with the nuclear receptor RXR, and are activated to enhance transcription of their target genes by binding a class of oxysterols including 22-hydroxycholesterol and 20-hydroxycholesterol (Janowski et al., *Nature*, 383:728-731 (1996)). As such, they are candidates for the mediation of cholesterol-induced gene transcription. Further, in light of studies which showed that ABC1 mRNA and protein increase in fibroblasts and macrophages in response to cholesterol loading, and other studies which showed that LXR and RXR expression increase in cholesterol-loaded macrophage cells by exposure to oxidized LDL, the LXR and RXR nuclear receptors are highly plausible candidates for transcriptional activators of the ABC1 gene. To determine whether LXR and RXR receptors play a role in ABC1 gene expression, the level of ABC1 mRNA was measured in response to 20-hydroxycholesterol and 9-*cis* retinoic acid.

Mouse RAW 264.7 cells were grown to subconfluence in DMEM/10% FBS and then treated for 24 hours in serum-free DMEM/BSA with either 9-*cis* retinoic acid (10 μ M), 20-hydroxycholesterol (10 μ M), or both ligands together (20 μ M total). Control cells received ethanol vehicle only (0.1% v/v). RNA was extracted, treated with DNase, and ABC1 mRNA measured by RT-PCR using PE Biosystems SYBR Green Technology as described in Example 9. Figure 9 shows that treatment with either 20-hydroxycholesterol or 9-*cis* retinoic acid results in an increase in ABC1 mRNA expression. In addition, Figure 9 shows that treatment with both ligands together results in a markedly synergistic effect, with an approximate 6-fold increase over the ABC1 expression observed with either ligand alone. These results demonstrate that ligands for the nuclear receptors LXR and RXR can increase the expression of the ABC1 gene.

EXAMPLE 11

This example demonstrates that enhanced expression of ABC1 protein in the plasma membrane is associated with lipid efflux.

Cell-surface labeling and immunoprecipitation was used to determine whether increased expression of ABC1 protein in the plasma membrane is correlated with an increase in cholesterol efflux (Figure 10). The relative amount of ABC1 on the cell surface was determined by cross-linking surface proteins on intact cells with the membrane-impermeable agent sulfo-NHS-biotin, followed by the steps of membrane solubilization, immunoprecipitation with ABC1 antibody, SDS-PAGE, and detection with streptavidin.

Cell Culture: Normal and TD1 fibroblast cells were immortalized as described in Example 1. Both normal and TD1 cells were cultured under control conditions and conditions

known to increase apolipoprotein-mediated cholesterol efflux (Oram, et al., *J. Lip. Res.*, 40: 1769-1781 (1999)). Control cells were grown to confluence in DMEM/10% FBS and then incubated in DMEM/BSA for 18 hours with no additives (control). cAMP-treated cells were grown to confluence in DMEM/10% FBS and then incubated in DMEM/BSA for 18 hours with 1mM 8-Br-cAMP(cAMP). Cholesterol-loaded cells were grown to confluence in DMEM/10% FBS and then incubated in DMEM/BSA for 48 hours with 30µg/ml cholesterol plus 18 hours with no additives (cholesterol). Cholesterol-loaded cells treated with cAMP were grown to confluence in DMEM/10% FBS and then incubated in DMEM/BSA for 48 hours with 30µg/ml cholesterol plus 18 hours with 1mM 8-Br-cAMP (cholesterol + cAMP).

10 Cell-Surface Labeling: For selective labeling of plasma membrane ABC1, the cells were incubated for 30 minutes at 0°C with PBS containing 1 mg/ml sulfo-NHS-biotin (Pierce, Rockford, IL; Cat. #21217) to biotinylate cell-surface proteins (see Walker et al., *Biochemistry*, 50:14009-14014 (1993)).

15 Immunoprecipitation: Rabbit antiserum for ABC1 was raised against a synthetic peptide corresponding to the deduced peptide KNQTVVDAVLTSFLQDEKVKES located at the C-terminus of human ABC1. Immunoprecipitation was performed by solubilizing the cells in PBS containing 1% Triton X-100 (Sigma, St. Louis, MO) and protease inhibitors leupeptin (1mM), pepstatin (1mM), and aprotinin (1mM). The cell extract was incubated overnight at 4°C with anti-ABC1 antiserum at 1:200 dilution followed by an additional 1 hour incubation with 5µl proteinA-coated magnetic beads (Dynal, Lake Success, NY; Cat. #1001.01). The antibody-antigen complex was sedimented with a magnet, the beads were washed twice with 1% Triton-X/PBS, and the proteins were eluted with 1% acetic acid.

20 Detection of ABC1 Protein: The eluted biotinylated proteins were subjected to SDS-PAGE (6% gel; 150V, 5 hours) and transferred to nitrocellulose membrane (200mA, 18 hours). The nitrocellulose was probed with streptavidin-horse radish peroxidase (Amersham Pharmacia, Piscataway, NJ; Cat. #RPN 1231) diluted 300-fold and detected by enhanced chemiluminescence labeling (ECL) according to vendor's protocol (Amersham Pharmacia, Piscataway, NJ). To test for possible biotinylation of intracellular proteins, the post-immunoprecipitation supernatant was treated with a mouse monoclonal antibody to the intracellular protein β-COP and immunoprecipitated biotinylated β-COP was assayed by streptavidin blotting. None was detected.

30 Results: As shown in Figure 10, the 240 kDa ABC1 protein appears as a doublet. The ABC1 protein is partially localized to the plasma membrane in both normal (10A) and TD1

(10B) fibroblast cells. Similar results were seen with a second normal fibroblast cell line and with TD2 fibroblasts (data not shown). Cell-surface expression of ABC1 was increased slightly when cells grown in serum (normal and TD1 cells) were treated with 8-Br-cAMP. Serum deprivation and cholesterol-loading of both normal and TD1 cells markedly increased cell-surface expression of ABC1, which was further enhanced by cAMP treatment. These results indicate that expression of ABC1 at the cell surface is regulated by conditions that enhance apolipoprotein-mediated lipid efflux, consistent with the idea that its localization to the plasma membrane plays a key role in its lipid transport function. The mutations in TD1 and TD2 cells do not appear to severely impair expression or processing of ABC1, implying that secondary effects on lipid transport or interactions with accessory proteins depend on its NH₂-terminal domain, where the mutations occur.

EXAMPLE 12

This example shows that agents that inhibit the degradation of 3'5' cyclic AMP, such as phosphodiesterase inhibitors, increase apolipoprotein A-I-mediated efflux from macrophage cells.

As shown in Figure 10, cAMP increases the activity of ABC1. The present studies were performed to determine the cholesterol efflux from macrophage cells in the presence of elevated cAMP. Elevated levels of cAMP can be attained in the presence of agents that either stimulate cAMP synthesis or inhibit the degradation of cAMP. For example, rolipram is a compound that regulates cAMP levels by inhibiting phosphodiesterases, a group of enzymes that degrade cAMP. The effect of elevated cAMP on cholesterol efflux was determined using the apolipoprotein-mediated cholesterol efflux assays described in Example 1. Briefly, RAW 264.7 cells suspended at a density of 1.25×10^5 cells/ml were grown in DMEM/10% FBS supplemented with pyruvate. After 24 hours, the medium was removed and replaced with DMEM/BSA plus radiolabeled cholesterol (1 μ Ci/ml ³[H]-cholesterol) and 50 μ g/ml of acetylated LDL for 24 hours. The cells were then maintained for 24 hours in equilibrium medium consisting of DMEM/BSA plus either apo A-I alone (20 μ g/ml), apo A-I and 8-bromo 3', 5' cAMP (1mM) or apo A-I and rolipram (50 μ M). After 12-24 hours, [³H]cholesterol efflux was measured by scintillation counting as described in Example 1 and calculated as the percentage of total radiolabeled cholesterol appearing in the medium. The results indicated that cholesterol-loaded control cells that received no apo A-I showed a 3% cholesterol efflux, while cells that received apo A-I only showed a 5% efflux. Cholesterol-loaded cells that received apo

A-I and cAMP showed a 32% cholesterol efflux, demonstrating that elevated cAMP promotes cholesterol efflux. Similarly, cells that received apo A-I and a phosphodiesterase inhibitor (rolipram) showed a 17% cholesterol efflux.

5

EXAMPLE 13

This example shows that agents that are ligands for nuclear receptors, such as LXR, RXR, and PPAR nuclear receptors, increase apolipoprotein A-I-mediated efflux from macrophage cells.

To determine whether ligands for nuclear receptors affect the process of apolipoprotein-mediated cholesterol efflux, various ligands were tested using the apo A-I-mediated cholesterol efflux assay described in Example 12. The nuclear receptor superfamily includes several members, such as the liver receptor LXR, the retinoid receptor RXR, and the peroxisome proliferator-activated receptor PPAR, which have been implicated in lipid metabolism (Russell, D.W., *Cell*, 97:539-542 (1999); Spiegelman, B.W., *Cell*, 93:153-155 (1998); Janowski et al., *Nature*, 383:728-731 (1996)). Further, ligands for some of these receptors have been observed to increase plasma HDL and gene expression profiling (microarray) data have shown that hormone receptors respond to cholesterol loading via oxidized LDL. Using the above-described assay, 9-*cis*-retinoic acid (RXR ligand), oxysterol (LXR ligand), and fenofibrate (PPAR ligand) was tested to determine the effect on cholesterol efflux. Cholesterol-loaded control cells that received no apo A-I showed a 3% cholesterol efflux, while cells that receive apo A-I only showed a 5% efflux. In contrast, cholesterol-loaded cells that receive apo A-I and 9-*cis*-retinoic acid (30 ng/ml) showed a 16% cholesterol efflux. Cells that receive apo A-I and oxysterol (5 µg/ml) showed a 14% cholesterol efflux. Cells that receive apo A-I and fenofibrate (3 µg/ml) showed a 10% cholesterol efflux. These results indicate that hormone receptors may be modulated to increase the rate of apolipoprotein-mediated cholesterol efflux from macrophages.

Further, when the efflux assay was performed using various concentrations of 9-*cis*-RA (0.3ng/ml, 3.0ng/ml, or 30ng/ml), the results showed that 9-*cis*-RA mediated cholesterol efflux from macrophage cells in a dose-dependent manner. Specifically, control cells (apo A-I only) showed 1890 c.p.m., 0.3 ng/ml 9-*cis*-RA showed 1522 c.p.m., 3.0 ng/ml 9-*cis*-RA showed 3568 c.p.m., and 30 ng/ml 9-*cis*-RA showed 8597 c.p.m. In addition, using a similar assay where RAW 264.7 cells were cholesterol-loaded for 48 hours, other nuclear receptor activators, such as 22-hydroxycholesterol (LXR ligand) and benzofibrate, were shown to increase cholesterol efflux (data not shown).

EXAMPLE 14

This example shows that eicosanoids, such as prostaglandin E1 and prostacyclin PG12, increase apolipoprotein A-I-mediated efflux from macrophage cells.

5 Eicosanoids, such as prostaglandins and prostacyclins, have been shown to be effective in the treatment of hypercholesterolemia. To determine whether eicosanoids affect the process of apolipoprotein-mediated cholesterol efflux, PGE1 and PG12 were tested using the apo A-I-mediated cholesterol efflux assay described in Example 12. This assay showed that cholesterol-loaded control cells that receive no apo A-I have a 3% cholesterol efflux, while cells that receive
10 apo A-I only have a 5% efflux. Cholesterol-loaded cells that receive apo A-I and PG12 (25 nm) showed a 10% cholesterol efflux. Cells that receive apo A-I and PGE1 (25 nM) showed a 15% cholesterol efflux. These results demonstrate that eicoasnoids can increase the rate of apolipoprotein-mediated cholesterol efflux from macrophages.

EXAMPLE 15

15 This example demonstrates that a reporter gene under the control of an ABC1 promoter can be used to test compounds for the ability to regulate ABC1 gene expression.

The pGL3 luciferase reporter vector system (Promega, Madison, WI) was used to create a recombinant plasmid to measure reporter gene expression under control of the ABC1 promoter.

20 Construction of Reporter Plasmids: Plasmid pGL3-Basic (Promega, Madison, WI; Cat. #E1751) was used as a control plasmid containing the promoterless luciferase gene. The reporter construct containing the ABC1 promoter and luciferase gene was made by cloning a genomic fragment from the 5' flanking region of the ABC1 gene (hAPR1 5' promoter, corresponding to nucleotides 1080-1643 of SEQ ID NO: 3) into the SacI site of the GL3-Basic plasmid to generate
25 plasmid GL-6a. Next, plasmid GL-6a was digested with SpeI and Acc65I. A BsiWI-SpeI fragment excised from a lambda subclone, representing the ABC1 genomic sequence corresponding to nucleotides 1-1534 of SEQ ID NO: 3 was ligated into the remaining vector/ABC1 promoter fragment produced by this digestion. The resultant plasmid, pAPR1, encodes the luciferase reporter gene under transcriptional control of 1.75 kb of the human ABC1
30 promoter sequence.

Transfection of Reporter Constructs: The above-described control or pAPR1 plasmid was transfected into confluent cultures of RAW 264.7 cells maintained in DMEM containing 10% fetal bovine serum. Each well of a 12 well dish was transfected for 5 hours with either

pGL3-Basic, pGL3-Promoter or pAPR1 DNA (1 µg), luciferase plasmid DNA (1 µg), and 12 µl of Geneporter reagent (Gene Therapy Systems, San Diego, CA; Cat. #T201007). In addition, 0.1 µg of pCMVβ plasmid DNA (Clontech, Palo Alto, CA, Cat. #6177-1) was added as a control for transfection efficiency. After 5 hours, the culture medium was replaced with serum-free
5 DMEM/BSA in the presence of or absence of acetylated LDL (100 µg/ml) and incubated for 24 hours.

For added convenience in high throughput screening, cultured cells can be stably transfected with reporter plasmids using the following procedure. First, 5×10^6 RAW 264.7 cells are transfected for 5 hours in a 60mm dish with 9 µg of the pAPR1 plasmid and pCMVscript
10 (Stratagene, LaJolla, CA) in 10 ml of serum-free DMEM with 50 µl Geneporter transfection reagent (Gene Therapy Systems, San Diego, CA). Subsequently, the transfection medium is replaced with complete medium and the cells incubated overnight at 37°C. Subsequently, the cells are transferred to separate dishes at dilutions ranging from 1:5 to 1:1000 and incubated in selection medium containing 800 µg/ml G418 (Life Technologies, Bethesda, MD) for 20 days.
15 Visible colonies are picked, expanded, and assayed for luciferase activity as described below. Using this method, five clonal cell lines positive for luciferase activity were identified for use in high throughput assays.

Luciferase Assay: Following transfection, the cells in each well were lysed in 70 µl of 1X cell lysis reagent (Promega, Madison, WI, Cat. #E3971), subjected to one freeze-thaw cycle,
20 and the lysate cleared by centrifugation for 5 minutes at 12,000g. After centrifugation, 100 µl of luciferase assay reagent (Promega, Madison, WI; Cat. #E1501) was added to 10 µl of lysate. The luciferase activity of each lysate was measured as light units using a luminometer. Additionally, the β-galactosidase activity of each lysate was measured using the chemiluminescent assay reagents supplied in the Galacto-light kit according to the
25 manufacturer's instructions (Tropix Inc., Bedford, MA; Cat. #BL100G). The normalized luciferase activity for each lysate was determined by dividing the luciferase activity value by the determined β-galactosidase value and reported as relative light units.

Results: The luciferase activity detected in cells transfected with pAPR1 was 3.3-fold higher than the activity detected in control cells transfected with pGL3-Basic plasmid containing
30 luciferase cDNA only. These results indicated that the transcriptional regulatory regions of ABC1 were in place. When the pAPR1 transfected cells were incubated with 100 µg/ml acetyl LDL for 24 hours, the luciferase activity was 3.25-fold higher than in cells that had not been treated with acetyl LDL. These results suggest that the genomic ABC1 sequence contains a

"cholesterol responsive" element found in the 5'flanking region which mediates the cholesterol loading response of the native ABC1 gene. This reporter system can also be used to test other compounds to determine whether the compound modulates ABC1 expression.

5

EXAMPLE 16

This example demonstrates an additional assay that can be used to test compounds for the ability to regulate ABC1 gene expression using a reporter gene under the control of an endogenous ABC1 promoter.

This assay involves constructing a recombination vector that contains a promoterless
10 reporter gene and a selection marker gene. The vector is linearized and transfected into cells such that the reporter gene is integrated into the cellular genome downstream of the endogenous ABC1 promoter. Using this assay, expression of the reporter gene is driven by the endogenous ABC1 promoter in response to a test compound.

Construction of Reporter Plasmids: The recombination vector containing a promoterless
15 reporter gene can be made starting with a 7 kb EcoRI genomic fragment of ABC1 that contains exon 0 (which includes the ABC1 start sites) and part of intron1. Using site-directed mutagenesis, a SalI restriction site can be generated in the exon 0 sequence downstream of the two known start sites. The recombination vector is generated by inserting a DNA fragment containing a promoterless reporter gene, such as luciferase, and a promoterless selective marker,
20 such as puromycin resistance, into the SalI site. An internal ribosome entry signal should be inserted between the reporter gene and marker gene so that the genes will be transcribed in the correct orientation. The recombination vector contains two Eco47III sites, one of which must be eliminated, using, for example, site-directed mutagenesis. The remaining Eco47III site, located upstream of of exon 0, is used to linearize the vector.

25 Transfection of Reporter Constructs: The linearized recombination vector containing the reporter gene and marker gene is introduced into cultured cells, including human cells, by any of the various transfection methods known in the art. For example, the linearized vector can be transfected using the methods described in Example 15. The linearized recombination vector contains ABC1 sequences which allow the vector to integrate into the cellular genome at the site
30 of the endogenous ABC1 gene. The addition of an appropriate antibiotic to the culture medium allows the selection of only those cells in which the reporter gene and marker gene have integrated downstream of the endogenous ABC1 promoter in the proper orientation. For instance, if the vector contains a puromycin resistance gene inserted downstream of the reporter

gene, the transfected cells should be grown in the presence of puromycin. Only those cells that have a properly integrated puromycin resistance gene, and can thereby encode a functional protein, will survive in the presence of puromycin. Thus, the transfected cells should be grown under conditions that induce ABC1 promoter activity in the presence of the appropriate
5 antibiotic. Surviving cells can be clonally cultured and the DNA sequenced using PCR or southern blot analysis to test for proper integration of genomic sequences.

The resultant cells containing a reporter gene under the control of the endogenous ABC1 promoter can be used to determine whether a test compound modulates the expression of ABC1. The ABC1 modulating activity of a compound is determined by assaying the level of reporter
10 gene expression found in the cells exposed to the test compound. For example, cells having an integrated luciferase gene can be used to determine the ABC1 modulating activity of a test compound by measuring the amount of luciferase activity found in cells exposed to the compound.

15

EXAMPLE 17

This example demonstrates that ligands for nuclear receptors up-regulate the expression of a reporter gene under the control of the ABC1 promoter.

To determine whether ligands for the LXR α , LXR β and RXR α nuclear receptors could regulate ABC1 gene expression, the pAPR1 plasmid containing the luciferase reporter gene
20 under control of the ABC1 promoter was transfected into RAW 264.7 cells treated with at least one ligand for the nuclear receptors (Figure 12).

Construction and Transfection of Reporter Constructs: Reporter construct pAPR1 and control reporter construct pGL3-Basic were obtained as described in Example 15. RAW 264.7 cells were maintained in culture and transfected with either pGL3-Basic (1 μ g) or pAPR1 (1 μ g)
25 as described in Example 15. The transfected RAW 264.7 cells were treated with either ethanol (EtOH) (0.1% v/v), 20(S)-hydroxycholesterol (20(S) OH-cho1) (10 μ M), 9-*cis* retinoic acid (9-*cis* RA) (10 μ M) or both 20(S) OH-cho1 and 9-*cis* RA (20 μ M total) for 24 hours. The luciferase activity was measured and reported as relative light units as described in Example 15.

Results: The results of this study are shown in Figure 12. Control cells transfected with
30 pGL3-Basic showed no luciferase activity (data not shown). Cells transfected with pAPR1 produced a 19-fold increase in luciferase reporter activity in the presence of 20 OH-cho1, a 16-fold increase in luciferase activity in the presence of 9-*cis* RA, and a 280-fold increase in luciferase activity in the presence of both ligands compared with EtOH control. These results

indicate that both the sterol and retinoid elicit a strong transcription response from the ABC1 5' flanking sequence in pAPR1. Further, there is an apparent synergistic effect of the two classes of compounds, as can be seen by the dramatic increase in luciferase activity found in cells treated with both ligands. It is known that LXR α and RXR α receptors form active heterodimers. Thus, the ligand-induced activation of both nuclear receptors simultaneously could produce the observed synergistic increase in transcription.

These data demonstrate that hydroxy sterols, such as 20(S) hydroxycholesterol, and retinoids, such as 9-*cis* retinoic acid, activate the ABC1 promoter, indicating that these and related compounds can be useful in the development of therapeutic compounds to increase ABC1 expression in macrophage cells to rid peripheral sites of excess cholesterol. Additionally, the present ABC1 promoter/reporter gene screening assay can be used to screen other compounds that increase ABC1 expression to identify further therapeutic compounds.

EXAMPLE 18

This example demonstrates the further characterization of the ABC1 promoter region, including the identification of an LXR response element.

To determine which portion of the 5' flanking region of ABC1 retains transcriptional activity in response to nuclear ligands, various plasmids containing a different portion of the 5' flanking region and a luciferase reporter gene were transfected into RAW 264.7 cells treated with at least one ligand for the nuclear receptors. Using this system, an sterol response element corresponding to nucleotides 1480-1510 of SEQ ID NO: 3 was identified. The sterol response element contains a direct repeat-4 element TGACCGatagTAACCT. Confirmation of the sterol response element was obtained using site-directed mutagenesis and band-shift assay techniques.

Construction of Reporter Constructs: Reporter construct pAPR1 and control reporter construct pGL3-Basic were obtained as described in Example 15. Reporter constructs containing either nucleotides 1-1532, 1080-1643, 1181-1643, 1292-1643, or 1394-1643 of SEQ ID NO: 3 were also constructed. A reporter construct containing nucleotides 1080-1643 of SEQ ID NO: 3 (GL-6a) was constructed as described in Example 15. A reporter construct containing nucleotides 1-1532 of SEQ ID NO: 3 was constructed by digestion of pAPR1 with SpeI and NheI, and re-ligation of the gel-purified vector fragment. A reporter construct containing nucleotides 1181 to 1643 was constructed by firstly digesting GL-6a with StyI, blunting the cohesive ends with Klenow enzyme, digesting the resultant vector with SacI, and isolating the 462 base pair blunt-SacI cohesive end fragment. This was cloned into a vector obtained by digestion of GL-6a

with Acc65 I, blunting of the cohesive ends with Klenow enzyme, digestion with Sac I and gel isolation of the vector fragment. A reporter construct containing nucleotides 1292-1643 was constructed by consecutive digestion of GL-6a with Acc65 I, blunting the ends with Klenow enzyme, digestion with Sac II, blunting the ends with T4 polymerase, and re-ligation of the gel-isolated vector fragment. A reporter construct containing nucleotides 1394-1643 was constructed by digestion of GL-6a with Acc65 I, blunting the ends with Klenow enzyme, subsequent digestion with Apa I, end-blunting with T4 polymerase and re-ligation of the gel-isolated vector fragment.

Transfection of Reporter Constructs: The RAW 264.7 cells were maintained in culture and transfected with either pGL3-Basic (1µg), pAPR1 (1µg), or one of the other reporter constructs according to the method described in Example 15. The transfected RAW 264.7 cells were treated with either ethanol (EtOH) (0.1% v/v), 20(S)-hydroxycholesterol (20(S) OH-cho) (10µM), 22(R)-hydroxycholesterol (22(R) OH-cho) (10µM), 9-*cis* retinoic acid (9-*cis* RA) (10µM), or both 20(S) OH-cho and 9-*cis* RA (20µM total) for 24 hours. The luciferase activity was measured and reported as relative light units as described in Example 15.

Site-Directed Mutagenesis: The sterol response element corresponding to nucleotides 1480-1510 of SEQ ID NO: 3 was mutated in the 1080-1643 sequence described above using site-directed mutagenesis. Specifically, the response element containing a direct repeat-4 element TGACCGatagTAACCT was mutated to CTGCACatagTAACCT using the GeneEditor system (Promega, Madison, WI) according to the manufacturer's protocol.

Gel-Shift Assays: Nuclear extract was prepared from RAW 264.7 cells by the method of Ohlsson et al., *Cell*, 45:35-44 (1986). ³²P-labeled oligonucleotides (5 ng) corresponding to the LXR response element (TCGAGTGACCGATAGTAACCTCTCGA; SEQ ID NO: 56) and its mutated counterpart (TCGAGCTGCACATAGTAACCTCTCGA; SEQ ID NO: 57) were individually incubated with 5 µg of nuclear protein for 30 minutes at room temperature in 20mM HEPES, pH 7.9, 60mM KCL, 1mM MgCl₂, 1mM DTT, 66.6µg/ml poly(dIdC), and 10% glycerol in the presence or absence of 1µg antiserum to LXRα and LXRβ (Santa Cruz Biotechnology, Cat. No. SC-1591, Santa Cruz, CA) or antiserum to RXR (Santa Cruz Biotechnology, Cat. No. SC-774, Santa Cruz, CA). The protein-DNA complexes were applied to a 4% non-denaturing polyacrylamide gel for 1.5 hours at 150V in 0.5X TBE buffer. The protein-DNA complexes were detected by autoradiography of the dried gel.

Results: Transfection with the individual reporter constructs containing the 5' flanking region corresponding to nucleotides 1-1643 (i.e., pAPR1), 1-1532, 1080-1643, 1181-1643, 1292-

1643, or 1394-1643 of SEQ ID NO: 3 each produced the same results. All of the individual constructs produced a 3 to 4-fold increase in luciferase reporter activity in the presence of 20 (S) OH-chol or 22 (R) OH-chol compared with EtOH control. Also, all of the individual constructs produced an 8 to 10-fold increase luciferase reporter activity in the presence of 9-*cis* RA. In addition, transfection with any of the constructs produced a 25 to 50-fold increase in luciferase activity in the presence of oxysterol ligand (either (20 (S) OH-chol or 22(R) OH-chol)) and retinoid ligand (9-*cis* RA) together compared with EtOH control, indicating a synergistic interaction. Each of the described constructs demonstrated comparable levels of luciferase activity in response to the ligands tested, indicating that even the shorter 5' flanking sequences contained transcriptional regulatory sequences for sterols and retinoids. Specifically, these results demonstrated that the transcriptional regulatory sequences for sterols and retinoids are located in the 5' flanking region corresponding to nucleotides 1394-1532 of SEQ ID NO: 3.

These results were confirmed by luciferase assays using a reporter construct containing the wild-type sequence corresponding to nucleotides 1080-1643 of SEQ ID NO: 3 and a reporter construct containing a mutated sequence corresponding to nucleotides 1080-1643 of SEQ ID NO: 3, wherein the sterol response element found at nucleotides 1480-1500 was mutated as described above. Transfection with the wild-type sequence produced a transcriptional response, as measured by an increase in luciferase activity, in the presence of either 20 (S) OH-chol or 9-*cis* RA alone and produced a synergistic response in the presence of both ligands together. In contrast, transfection with the mutated sequence did not produce a transcriptional response in the presence of 20 (S) OH-chol or 22 (R) OH-chol. Transfection of the mutated sequence preserved a reduced response to 9-*cis* RA, producing a 4 to 5-fold increase in transcriptional activity, rather than the 8 to 10-fold increase observed with the wild-type sequence. Transfection of the mutated sequence also abolished the synergistic transcriptional response seen in the presence of 20 (S) OH-chol and 9-*cis* RA together. These results were further confirmed by gel-shift assays using the sterol consensus sequence (nucleotides 1480-1510) and its mutated counterpart. The gel-shift assays showed that while nuclear binding proteins isolated from RAW 264.7 cells bound to the sterol consensus sequence, nuclear proteins did not bind to the mutated sequence. Furthermore, incubation of nuclear proteins with the wild-type sterol consensus sequence in the presence of LXR antiserum resulted in the formation of supershifted complexes (i.e. antibody-protein-DNA complexes), identifying the sequence as a sterol response element that binds nuclear receptor LXR. In contrast, incubation of nuclear proteins with the wild-type sterol response element in the presence of RXR antiserum did not result in the formation of supershifted complexes,

indicating that RXR does not bind to this sequence. These results show that the mutation which destroys nuclear protein binding to the consensus sequence also abolishes the transcriptional response to LXR ligands and diminishes the response to RXR ligands. Furthermore, the nuclear binding studies performed in the presence of LXR or RXR antiserum confirmed that the
5 consensus sequence found at nucleotides 1480-1510 is an LXR response element. This element also mediates a partial response to 9-*cis* RA.

All of the references cited herein, including patents and publications, are hereby incorporated in their entireties by reference. While the invention has been described with an
10 emphasis upon preferred aspects of the invention, it will be readily apparent to those of ordinary skill in the art that variations of the preferred embodiments can be used and that it is intended that the invention can be practiced otherwise than is specifically described herein. Accordingly, the present invention includes all modifications encompassed within the spirit and scope of the invention as defined by the following claims.

What we claim is:

1. An isolated polypeptide comprising SEQ ID NO: 2.
2. An isolated polypeptide comprising an amino acid sequence that has at least 98% identity with the amino acid sequence of SEQ ID NO: 2.
- 5 3. An isolated polynucleotide selected from the group consisting of a polynucleotide encoding the polypeptide of claim 1, a polynucleotide comprising SEQ ID NO: 1, a polynucleotide comprising nucleotides 291-7074 of SEQ ID NO: 1, and a polynucleotide encoding the polypeptide of claim 2.
4. An isolated polynucleotide comprising a nucleotide sequence that is
10 complementary to the polynucleotide of claim 3.
5. An isolated polynucleotide comprising a nucleotide sequence that has at least 90% identity with the polynucleotide comprising SEQ ID NO: 1.
6. An isolated polynucleotide comprising a nucleotide sequence that has at least 95% identity with the polynucleotide comprising SEQ ID NO: 1.
- 15 7. An isolated polynucleotide comprising a nucleotide sequence that is complementary to the polynucleotide of claim 5.
8. A composition comprising the polynucleotide of claim 3 and a suitable carrier.
9. A composition comprising the polynucleotide of claim 4 and a suitable carrier.
10. A composition comprising the polynucleotide of claim 5 and a suitable carrier.
- 20 11. A recombinant vector comprising the polynucleotide of claim 3.
12. A recombinant vector comprising the polynucleotide of claim 4.
13. A recombinant vector comprising the polynucleotide of claim 5.
14. The recombinant vector of claim 11, further comprising a heterologous promoter polynucleotide.
- 25 15. The recombinant vector of claim 12, further comprising a heterologous promoter polynucleotide.
16. The recombinant vector of claim 13, further comprising a heterologous promoter polynucleotide.
- 30 17. The recombinant vector of claim 14, wherein said heterologous promoter is a cytomegalovirus promoter.
18. The recombinant vector of claim 17, wherein said vector is pCEPhABC1.
19. A composition comprising the recombinant vector of claim 11.

20. A composition comprising the recombinant vector of claim 13.
21. A composition comprising the recombinant vector of claim 14.
22. A host cell comprising the recombinant vector of claim 11.
23. A host cell comprising the recombinant vector of claim 13.
- 5 24. A host cell comprising the recombinant vector of claim 14.
25. An isolated polypeptide comprising SEQ ID NO: 8.
26. An isolated polynucleotide encoding the polypeptide of claim 25.
27. An isolated polynucleotide comprising SEQ ID NO: 7.
28. An isolated polypeptide comprising SEQ ID NO: 10.
- 10 29. An isolated polynucleotide encoding the polypeptide of claim 28.
30. An isolated polynucleotide comprising SEQ ID NO: 9.
31. A recombinant vector comprising the polynucleotide of claim 27.
32. A recombinant vector comprising the polynucleotide of claim 30.
33. A method for producing an ABC1 protein in a mammalian host cell comprising
- 15 the steps of:
- (a) transfecting the mammalian host cell with a recombinant expression vector comprising a polynucleotide encoding ABC1 in an amount sufficient to produce a detectable level of ABC1 protein; and
- (b) purifying the produced ABC1 protein.
- 20 34. The method of claim 33, wherein the recombinant expression vector comprises the polynucleotide of claim 3.
35. A method for expressing ABC1 in the cells of a mammalian subject comprising the step of administering to a mammalian subject a recombinant expression vector comprising a polynucleotide encoding ABC1 in an amount sufficient to express ABC1 in the cells of a
- 25 mammalian subject.
36. The method of claim 35, wherein the recombinant expression vector comprises the polynucleotide of claim 3.
37. A method suitable for increasing cholesterol efflux from cells of a mammalian subject comprising administering to the mammalian subject a recombinant expression vector
- 30 comprising a polynucleotide encoding ABC1 in an amount sufficient to increase cholesterol efflux from said cells.
38. The method of claim 37, wherein the recombinant expression vector comprises the polynucleotide of claim 3.

39. The method of claim 37, wherein the recombinant expression vector is a viral delivery vector.

40. The method of claim 39, wherein the viral delivery vector is an adenoviral vector.

41. The method of claim 39, wherein the viral vector is a lentiviral vector.

5 42. The method of claim 37, wherein the recombinant expression vector is a non-viral delivery vector.

43. A method suitable for increasing cholesterol efflux from cells of a mammalian subject comprising administering to the mammalian subject via a non-viral delivery system a polynucleotide encoding ABC1 in an amount sufficient to increase cholesterol efflux from said
10 cells.

44. The method of claim 43, wherein the polynucleotide encoding ABC1 comprises the polynucleotide of claim 3.

45. The method of claim 43, wherein the non-viral delivery system is selected from the group consisting of DNA-ligand complexes, adenovirus-ligand-DNA complexes, adeno-
15 associated virus-ligand-DNA complexes, direct injection of DNA, CaPO_4 precipitation, gene gun techniques, electroporation, liposomes, and lipofection.

46. A method suitable for increasing cholesterol efflux from cells of a mammalian subject comprising the step of administering to the mammalian subject a therapeutic amount of a compound that increases the expression of ABC1 in said cells.

20 47. The method of claim 46, wherein said compound is a cAMP analogue.

48. The method of claim 47, wherein said compound is selected from the group consisting of 8-bromo cAMP, N6-benzoyl cAMP, and 8-thiomethyl cAMP.

49. The method of claim 46, wherein said compound increases the synthesis of cAMP.

25 50. The method of claim 49, wherein said compound is forskolin.

51. The method of claim 46, wherein said compound is a phosphodiesterase inhibitor.

52. The method of claim 51, wherein said compound is selected from the group consisting of rolipram, theophylline, 3-isobutyl-1-methylxanthine, R020-1724, vinpocetine, zaprinast, dipyridamole, milrinone, amrinone, pimobendan, cilostamide, enoximone,
30 peroximone, and vesnarinone.

53. A method suitable for increasing the gene expression of ABC1 in the cells of a mammalian subject comprising the step of administering to the mammalian subject a cAMP analogue in an amount sufficient to increase the expression of ABC1 in said cells.

54. A method suitable for increasing cholesterol efflux of the cells of a mammalian subject by administering to the mammalian subject a compound that increases ABC1 activity in an amount sufficient to increase cholesterol efflux.

55. A method for screening a test compound to determine whether the test compound
5 promotes ABC1-mediated cholesterol efflux from cells in culture comprising the steps of:

assaying the level of cholesterol efflux in a sample of mammalian cells maintained in culture to determine a control level of cholesterol efflux;

contacting the cells with the test compound being screened;

assaying the level of cholesterol efflux in a sample of cells after contact with the test
10 compound; and

assaying the level of ABC1-mediated cholesterol efflux in a sample of cells after contact with the test compound, thereby determining whether the test compound promotes ABC1-mediated cholesterol efflux from cells in culture.

56. The method of claim 55, wherein the cultured cells are derived from a cell line.

15 57. The method of claim 56, wherein the cell line is selected from the group consisting of fibroblast, macrophage, hepatic, and intestinal cell lines.

58. The method of claim 57, wherein the cell line is RAW 264.7.

59. The method of claim 55, wherein the level of ABC1-dependent cholesterol efflux is assayed using an anti-ABC1 antibody that inhibits the activity of ABC1 upon binding.

20 60. The method of claim 55, wherein the level of ABC1-dependent cholesterol efflux is assayed using an anti-sense ABC1 polynucleotide.

61. The method of claim 60, wherein the polynucleotide comprises SEQ ID NO: 57.

62. A method for detecting the comparative level of ABC1 expression in the cells of a mammalian subject comprising the steps of:

25 obtaining a cell sample from the mammalian subject;

assaying the level of ABC1 mRNA expression in the cell sample; and

comparing the level of ABC1 mRNA expression in the cell sample with a pre-determined standard level of ABC1 mRNA expression, thereby detecting the comparative level of ABC1 gene expression in the cells of a mammalian subject.

30 63. The method of claim 62, wherein detection of the comparative level of ABC1 expression in cells of a mammalian subject is used to determine a susceptibility to coronary heart disease of the mammalian subject.

64. The method of claim 62, wherein the level of ABC1 mRNA expression is assayed by reverse transcription polymerase chain reaction, northern blot, or RNase protection assay.

65. A method for detecting the comparative level of ABC1 protein in the cells of a mammalian subject comprising the steps of :

- 5 obtaining a cell sample from the mammalian subject;
 assaying the amount of ABC1 protein in the cell sample; and
 comparing the amount of ABC1 protein in the cell sample with a pre-determined standard amount of ABC1 protein, thereby detecting the comparative level of ABC1 protein in the cells of the mammalian subject.

10 66. The method of claim 65, wherein detection of the comparative level of ABC1 protein in cells of a mammalian subject is used to determine a susceptibility to coronary heart disease of the mammalian subject.

67. The method of claim 65, wherein the assay to determine the amount of ABC1 protein comprises an immunoassay.

15 68. The method of claim 67, wherein the amount of ABC1 protein is determined by (a) contacting the cell sample with a population of anti-ABC1 antibodies and (b) detecting the specific binding ABC1 antibodies associated with the cell sample.

69. The method of claim 68, wherein the ABC1 antibodies are detected by western blotting, immunoprecipitation, or FACS.

20 70. An isolated antibody that binds specifically to the isolated polypeptide of claim 1 or claim 2.

71. The antibody of claim 70 wherein the antibody is a monoclonal antibody.

72. The antibody of claim 70 wherein the antibody is a polyclonal antibody.

25 73. The antibody of claim 70 wherein the antibody, upon binding to an ABC1 polypeptide, inhibits the cholesterol transport activity of the ABC1 polypeptide.

74. A kit suitable for screening a compound to determine whether the compound modulates ABC1-dependent cholesterol efflux comprising an inactivating anti-ABC1 antibody in an amount sufficient for at least one assay and instructions for use.

30 75. A kit suitable for screening a compound to determine whether the compound modulates ABC1-dependent cholesterol efflux comprising an antisense ABC1 oligonucleotide in an amount sufficient for at least one assay and instructions for use.

76. The kit of claim 75, wherein the antisense ABC1 oligonucleotide comprises SEQ ID NO: 53.

1/43




FIG. 1

2/13




FIG. 2

3/13




FIG. 3

4/13

CVT ABCA1 Gene Structure




FIG. 4

5143




FIG. 5

6/13




FIG. 6

7112




FIG. 7

8/13




FIG. 8

9/13




FIG. 9

10/17




FIG. 10

11/13




FIG. 11

12/13




FIG. 12

13/13


[illegible]

FIG. 13

SEQUENCE LISTING

<110> Lawn, Richard M.

Wade, David

Garvin, Michael

<120> Compositions and Methods for Increasing Cholesterol
Efflux and Raising HDL using ATP Binding Cassette
Transporter Protein ABC1

<130> 99,395-B

<140>

<141>

<150> US 60/140,264

<151> 1999-06-18

<150> US 60/153,872

<151> 1999-08-14

<150> US 60/166,573

<151> 1999-11-19

<160> 57

<170> PatentIn Ver. 2.0

<210> 1

<211> 10442

<212> DNA

<213> Homo sapiens

<400> 1

ggccgggacc cgcagagccg agccgaccct tctctcccgg gctgcggcag ggcagggcgg 60
ggagctccgc gcaccaacag agccggttct cagggcgctt tgctccttgt tttttccccc 120
gttctgtttt ctccccttct ccggaaggct tgtcaagggg taggagaaag agacgcaaac 180
acaaaagtgg aaaacagtta atgaccagcc acgggcgtcc ctgctgtgag ctctggccgc 240
tgcttccag ggctcccgag ccacacgctg ggcgtgctgg ctgagggaac atggcttggt 300
ggcctcagct gaggttgctg ctgtggaaga acctcacttt cagaagaaga caaacatgtc 360
agctgttact ggaagtggcc tggcctctat ttatcttctt gatcctgac tctgttcggc 420
tgagctaccc accctatgaa caacatgaat gccattttcc aaataaagcc atgccctctg 480
caggaacact tccttgggtt caggggatta tctgtaatgc caacaacccc tgtttccgtt 540
acccgactcc tggggaggct cccggagttg ttggaaactt taacaaatcc attgtggctc 600
gcctgttctc agatgctcgg aggttctctt tatacagcca gaaagacacc agcatgaagg 660
acatgcgcaa agttctgaga acattacagc agatcaagaa atccagctca aacttgaagc 720
ttcaagattt cctggtggac aatgaaacct tctctgggtt cctatatcac aacctctctc 780
tcccaaagtc tactgtggac aagatgctga gggctgatgt cattctccac aaggatattt 840
tgcaaggcta ccagttacat ttgacaagtc tgtgcaatgg atcaaaatca gaagagatga 900
ttcaacttgg tgaccaagaa gtttctgagc tttgtggcct accaaaggag aaactggctg 960
cagcagagcg agtacttcgt tccaacatgg acatcctgaa gccaatcctg agaacactaa 1020
actctacatc tcccttcccg agcaaggagc tggctgaagc cacaaaaaca ttgctgcata 1080
gtcttgggac tctggcccag gagctgttca gcatgagaag ctggagtgc atgcgacagg 1140
aggatgatgt tctgaccaat gtgaacagct ccagctctc caccctaatc taccaggctg 1200
tgtctcgtat tgtctgcggg catcccagg gaggggggct gaagatcaag tctctcaact 1260
ggatgagga caacaactac aaagccctct ttggaggcaa tggcactgag gaagatgctg 1320
aaaccttcta tgacaactct acaactcctt actgcaatga tttgatgaag aatttgaggt 1380

ctagtcctct tccccgatt atctggaaag ctctgaagcc gctgctcgtt gggaagatcc 1440
tgtatacacc tgacactcca gccacaaggc aggtcatggc tgaggtgaac aagaccttcc 1500
aggaaactggc tgtgttccat gatctggaag gcatgtggga ggaactcagc cccaagatct 1560
ggaccttcat ggagaacagc caagaaatgg acctgtccg gatgctgttg gacagcaggg 1620
acaatgacca cttttgggaa cagcagttgg atggcttaga ttggacagcc caagacatcg 1680
tggcgttttt ggccaagcac ccagaggatg tccagtccag taatggttct gtgtacacct 1740
ggagagaagc tttcaacgag actaaccagg caatccggac catatctcgc ttcattggagt 1800
gtgtcaacct gaacaagcta gaacccatag caacagaagt ctggctcatc aacaagtcca 1860
tggagctgct ggatgagagg aagttctggg ctggtattgt gttcactgga attactccag 1920
gcagcattga gctgccccat catgtcaagt acaagatccg aatggacatt gacaatgtgg 1980
agaggacaaa taaaatcaag gatgggtact gggaccctgg tcctcgagct gacccttttg 2040
aggacatgcg gtacgtctgg gggggcttcg cctacttgca ggatgtggtg gaggaggcaa 2100
tcacaggggt gctgacgggc accgagaaga aaactgggtg ctatatgcaa cagatgccct 2160
atccctgtta cgttgatgac atctttctgc ggggtgatgag ccggtcaatg cccctcttca 2220
tgacgctggc ctggatttac tcagtggctg tgatcatcaa gggcatcgtg tatgagaagg 2280
aggcacggct gaaagagacc atgcggatca tgggcctgga caacagcata ctctggttta 2340
gctggttcat tagtagctc attcctcttc ttgtgagcgc tggcctgcta gtggtcatcc 2400
tgaagttagg aaacctgctg ccctacagtg atcccagcgt ggtgtttgtc ttctgtccg 2460
tgtttgctgt ggtgacaatc ctgcagtgtc tcctgattag cacactcttc tccagagcca 2520
acctggcagc agcctgtggg ggcatcatct acttcacgct gtacctgcc tacgtcctgt 2580
gtgtggcatg gcaggactac gtgggcttca cactcaagat cttecgtagc ctgctgtctc 2640
ctgtggcttt tgggtttggc tgtgagtact ttgccctttt tgaggagcag ggcattggag 2700
tgcagtggga caacctgttt gagagtctg tggaggaaga tggcttcaat ctcaccactt 2760
cgatctccat gatgctgttt gacaccttcc tctatggggg gatgacctg tacattgagg 2820
ctgtctttcc aggccagtac ggaattccca ggccctggta ttttccttgc accaagtcct 2880
actggtttg cgaggaaagt gatgagaaga gccaccctgg tccaaccag aagagaatgt 2940
cagaaatctg catggaggag gaaccaccc acttgaagct gggcgtgtcc attcagaacc 3000
tggtaaaagt ctaccgagat gggatgaagg tggctgtcga tggcctggca ctgaattttt 3060

atgagggcca gatcacctcc ttcctgggcc acaatggagc ggggaagacg accaccatgt 3120
caatcctgac cgggttgttc cccccgacct cgggcaccgc ctacatcctg ggaaaagaca 3180
ttcgctctga gatgagcacc atccggcaga acctgggggt ctgtccccag cataacgtgc 3240
tgtttgacat gctgactgtc gaagaacaca tctggttcta tgcccgttg aaagggtctt 3300
ctgagaagca cgtgaaggcg gagatggagc agatggccct ggatgttggt ttgccatcaa 3360
gcaagctgaa aagcaaaaca agccagctgt cagggtggaat gcagagaaag ctatctgtgg 3420
ccttggcctt tgtcggggga tctaagggtg tcattctgga tgaaccaca gctggtgtgg 3480
acccttactc ccgcagggga atatgggagc tgctgctgaa ataccgacaa ggccgcacca 3540
ttattctctc tacacaccac atggatgaag cggacgtcct gggggacagg attgccatca 3600
tctcccatgg gaagctgtgc tgtgtgggt cctccctgtt tctgaagaac cagctgggaa 3660
caggctacta cctgacctg gtcaagaaag atgtggaatc ctccctcagt tcctgcagaa 3720
acagtagtag cactgtgtca tacctgaaaa aggaggacag tgtttctcag agcagttctg 3780
atgctggcct gggcagcgac catgagagtg acacgctgac catcgatgtc tctgctatct 3840
ccaacctcat caggaagcat gtgtctgaag cccggctggt ggaagacata gggcatgagc 3900
tgacctatgt gctgccatat gaagctgcta aggagggagc ctttgtggaa ctctttcatg 3960
agattgatga ccggtctca gacctgggca tttctagtta tggcatctca gagacgaccc 4020
tggaagaaat attcctcaag gtggccgaag agagtggggt ggatgctgag acctcagatg 4080
gtaccttgcc agcaagacga aacaggcggg ccttcgggga caagcagagc tgtcttcgcc 4140
cgttcactga agatgatgct gctgatccaa atgattctga catagacca gaatccagag 4200
agacagactt gctcagtggg atggatggca aagggccta ccaggtgaaa ggctggaaac 4260
ttacacagca acagtttgtg gcccttttgt ggaagagact gctaattgcc agacggagtc 4320
ggaaaggatt ttttgctcag attgtcttgc cagctgtgtt tgtctgcatt gcccttgtgt 4380
tcagcctgat cgtgccaccc tttggcaagt acccagcct ggaacttcag ccctggatgt 4440
acaacgaaca gtacacattt gtcagcaatg atgtcctga ggacacggga accctggaac 4500
tcttaaacgc cctcaccaa gaccctggct tcgggaccg ctgtatggaa ggaaaccaa 4560
tcccagacac gccctgccag gcaggggagg aagagtggac cactgcccc gttccccaga 4620
ccatcatgga cctcttcag aatgggaact ggacaatgca gaacccttca cctgcatgcc 4680
agtgtagcag cgacaaaatc aagaagatgc tgctgtgtg tccccaggg gcaggggggc 4740
tgctctctcc acaaagaaaa caaaacactg cagatctcct tcaggacctg acaggaagaa 4800

acatttcgga ttatctggtg aagacgtatg tgcagatcat agccaaaagc ttaaagaaca 4860
agatctgggt gaatgagttt aggtatggcg gcttttccct ggggtgtagt aatactcaag 4920
cacttcctcc gagtcaagaa gttaatgatg ccatcaaaca aatgaagaaa cacctaaagc 4980
tggccaagga cagttctgca gatcgatttc tcaacagctt gggaagattt atgacaggac 5040
tggacaccag aaataatgtc aagggtgtggt tcaataacaa gggctggcat gcaatcagct 5100
ctttcctgaa tgtcatcaac aatgccattc tccgggccaa cctgcaaaag ggagagaacc 5160
ctagccatta tggaattact gctttcaatc atccccgaa tctcaccaag cagcagctct 5220
cagaggtggc tctgatgacc acatcagtggt atgtccttgt gtccatctgt gtcattcttg 5280
caatgtcctt cgtcccagcc agctttgtcg tattcctgat ccaggagcgg gtcagcaaag 5340
caaaacacct gcagttcatc agtggagtga agcctgtcat ctactggctc tctaattttg 5400
tctgggatat gtgcaattac gttgtccctg ccacactggc cattatcatc ttcattctgt 5460
tccagcagaa gtcctatgtg tctccacca atctgcctgt gctagccctt ctacttttgc 5520
tgtatgggtg gtcaatcaca cctctcatgt acccagcctc ctttgtgttc aagatcccca 5580
gcacagccta tgtggtgctc accagcgtga acctcttcat tggcattaat ggcagcgtgg 5640
ccacctttgt gctggagctg ttcaccgaca ataagctgaa taatatcaat gatatcctga 5700
agtccgtgtt cttgatcttc ccacattttt gcctgggacg agggctcatc gacatggtga 5760
aaaaccaggc aatggctgat gccctggaaa ggtttgggga gaatcgcttt gtgtcaccat 5820
tatcttggga cttgggtggga cgaaacctct tcgccatggc cgtggaaggg gtggtgttct 5880
tctcattac tgttctgatc cagtacagat tcttcatcag gccagacct gtaaatacaa 5940
agctatctcc tctgaatgat gaagatgaag atgtgaggcg ggaaagacag agaattcttg 6000
atggtggagg ccagaatgac atcttagaaa tcaaggagt gacgaagata tatagaagga 6060
agcggaagcc tgctgttgac aggatttgcg tgggcattcc tcctggtgag tgctttgggc 6120
tcctgggagt taatggggct ggaaaatcat caactttcaa gatgttaaca ggagatacca 6180
ctgttaccag aggagatgct ttccttaaca aaaatagtat cttatcaaac atccatgaag 6240
tacatcagaa catgggctac tgccctcagt ttgatgccat cacagagctg ttgactggga 6300
gagaacacgt ggagttcttt gcccttttga gaggagtccc agagaaagaa gttggcaagg 6360
ttggtgagtg ggcgattcgg aaactgggcc tcgtgaagta tggagaaaaa tatgctggta 6420
actatagtgg aggcaacaaa cgcaagctct ctacagccat ggctttgatc ggcgggcctc 6480

ctgtggtggt tctggatgaa cccaccacag gcatggatcc caaagcccg cggttcttgt 6540
ggaattgtgc cctaagtgtt gtcaaggagg ggagatcagt agtgcttaca tctcatagta 6600
tggagaagtg tgaagctctt tgacttagga tggcaatcat ggtcaatgga aggttcaggt 6660
gccttggcag tgtccagcat ctaaaaata ggtttggaga tggttatata atagtgttac 6720
gaatagcagg gtccaacccg gacctgaagc ctgtccagga tttctttgga cttgcatttc 6780
ctggaagtgt tctaaaagag aaacaccgga acatgctaca ataccagctt ccatcttcat 6840
tatcttctct ggccaggata ttcagcatcc tctcccagag caaaaagcga ctccacatag 6900
aagactactc tgtttctcag acaacacttg accaagtatt tgtgaacttt gccaaaggacc 6960
aaagtgatga tgaccactta aaagacctct cattacacaa aaaccagaca gtagtggacg 7020
ttgcagttct cacatctttt ctacaggatg agaaagtga agaaagctat gtatgaagaa 7080
tcctgttcat acgggggtggc tgaaagtaaa gaggaactag actttccttt gcaccatgtg 7140
aagtgttgtg gagaaaagag ccagaagttg atgtgggaag aagtaaactg gatactgtac 7200
tgatactatt caatgcaatg caattcaatg caatgaaaac aaaattccat tacaggggca 7260
gtgcctttgt agcctatgtc ttgtatggct ctcaagtga agacttgaat ttagtttttt 7320
acctatacct atgtgaaact ctattatgga acccaatgga catatgggtt tgaactcaca 7380
cttttttttt ttttttgttc ctgtgtattc tcattggggg tgcaacaata attcatcaag 7440
taatcatggc cagcgattat tgatcaaaat caaaaggtaa tgcacatcct cattcactaa 7500
gccatgccat gccaggaga ctggtttccc ggtgacacat ccattgctgg caatgagtgt 7560
gccagagtta ttagtgccaa gtttttcaga aagtttgaag caccatgggtg tgtcatgctc 7620
acttttgtga aagctgctct gtcagagtc tatcaacatt gaatatcagt tgacagaatg 7680
gtgccatgcg tggctaacat cctgctttga ttccctctga taagctgttc tgggtggcagt 7740
aacatgcaac aaaaatgtgg gtgtctctag gcacgggaaa cttggttcca ttgttatatt 7800
gtcctatgct tcgagccatg ggtctacagg gtcattcctta tgagactctt aaatatactt 7860
agatcctggt aagaggcaaa gaatcaacag ccaaactgct ggggctgcaa gctgctgaag 7920
ccagggcatg ggattaaaga gattgtgctg tcaaacctag ggaagcctgt gccatttgt 7980
cctgactgtc tgctaacatg gtacactgca tctcaagatg tttatctgac acaagtgtat 8040
tatttctggc tttttgaatt aatctagaaa atgaaaagat ggagttgtat tttgacaaaa 8100
atgtttgtac tttttaatgt tatttggaat ttaagtctt atcagtgact tctgaatcct 8160
tagaatggcc tctttgtaga accctgtggt atagaggagt atggccactg cccactatt 8220

tttattttct tatgtaagtt tgcatatcag tcatgactag tgcctagaaa gcaatgtgat 8280
ggtcaggatc tcatgacatt atatttgagt ttctttcaga tcatttagga tactcttaat 8340
ctcacttcat caatcaaata ttttttgagt gtatgctgta gctgaaagag tatgtacgta 8400
cgtataagac tagagagata ttaagtctca gtacacttcc tgtgccatgt tattcagctc 8460
actggtttac aaatataggt tgtcttgagg ttgtaggagc ccactgtaac aatattgggc 8520
agcctttttt tttttttttt aattgcaaca atgcaaaagc caagaaagta taagggtcac 8580
aagtttaaac aatgaattct tcaacaggga aaacagctag cttgaaaact tgctgaaaaa 8640
cacaacttgt gtttatggca tttagtacct tcaaataatt ggctttgcag atattggata 8700
ccccattaaa tctgacagtc tcaaattttt catctcttca atcactagtc aagaaaaata 8760
taaaaacaac aaatacttcc atatggagca tttttcagag ttttctaacc cagtcttatt 8820
tttctagtca gtaaacattt gtaaaaatac tgtttacta atacttactg ttaactgtct 8880
tgagagaaaa gaaaaatatg agagaactat tgtttgggga agttcaagtg atctttcaat 8940
atcattacta acttcttcca ctttttccaa aatttgaata ttaacgctaa aggtgtaaga 9000
cttcagattt caaattaatc tttctatatt ttttaaattt acagaatatt atataacca 9060
ctgctgaaaa agaaaaaat gattgtttta gaagttaaag tcaatattga ttttaaatat 9120
aagtaatgaa ggcataattc caataactag tgatatggca tcgttgcaat ttacagtatc 9180
ttcaaaaata cagaatttat agaataattt ctcttcattt aatatttttc aaaatcaaag 9240
ttatggtttc ctcatcttac taaaatcgta ttctaattct tcattatagt aaatctatga 9300
gcaactcctt acttcggttc ctctgatttc aaggccatat tttaaaaaat caaaaggcac 9360
tgtgaactat tttgaagaaa acacgacatt ttaatacaga ttgaaaggac ctcttctgaa 9420
gctagaaaca atctatagtt atacatcttc attaatactg tgttaccttt taaaatagta 9480
attttttaca ttttcctgtg taaacctaata tgtggtagaa atttttacca actctatact 9540
caatcaagca aaatttctgt atattccctg tggaatgtac ctatgtgagt ttcagaaatt 9600
ctcaaaatac gtgttcaaaa atttctgctt ttgcatcttt gggacacctc agaaaactta 9660
ttaacaactg tgaatatgag aaatacagaa gaaaataata agccctctat acataaatgc 9720
ccagcacaat tcattgttaa aaaacaacca aacctcacac tactgtattt cattatctgt 9780
actgaaagca aatgctttgt gactattaaa tgttgacacat cattcattca ctgtatagta 9840
atcattgact aaagccattt gctgtgtttt cttcttgagg ntgnatatat caggtaaaat 9900

attttccaaa gagccatgtg tcatgtaata ctgaaccctt tgatattgag acattaattt 9960
 ggacccttgg tattatctac tagaataatg taatactgna gaaatattgc tctaattctt 10020
 tcaaaatggg gcatccccct taaaangttc tatttcata aggatttagc ttgcttatcc 10080
 cttcttatac cctaagatga agctgttttt gtgctctttg ttcattcattg gccctcattc 10140
 caagcacttt acgctgtctg taatgggacg tatttttgca ctggaatatac tgagaattgc 10200
 aaaactagac aaaagtttca caacagattt ctaagttaaa tcattttcat taaaaggaaa 10260
 aaagaaaaaa aattttgtat gtcaataact ttatatgaag tattaaaatg catatttcta 10320
 tgttgtaata taatgagtca caaaataaag ctgtgacagt tctgttaaaa aaaaaaaaaa 10380
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 10440
 aa 10442

<210> 2

<211> 2261

<212> PRT

<213> Homo sapiens

<400> 2

Met Ala Cys Trp Pro Gln Leu Arg Leu Leu Leu Trp Lys Asn Leu Thr

1 5 10 15

Phe Arg Arg Arg Gln Thr Cys Gln Leu Leu Leu Glu Val Ala Trp Pro

20 25 30

Leu Phe Ile Phe Leu Ile Leu Ile Ser Val Arg Leu Ser Tyr Pro Pro

35 40 45

Tyr Glu Gln His Glu Cys His Phe Pro Asn Lys Ala Met Pro Ser Ala

50 55 60

Gly Thr Leu Pro Trp Val Gln Gly Ile Ile Cys Asn Ala Asn Asn Pro
65 70 75 80

Cys Phe Arg Tyr Pro Thr Pro Gly Glu Ala Pro Gly Val Val Gly Asn
85 90 95

Phe Asn Lys Ser Ile Val Ala Arg Leu Phe Ser Asp Ala Arg Arg Leu
100 105 110

Leu Leu Tyr Ser Gln Lys Asp Thr Ser Met Lys Asp Met Arg Lys Val
115 120 125

Leu Arg Thr Leu Gln Gln Ile Lys Lys Ser Ser Ser Asn Leu Lys Leu
130 135 140

Gln Asp Phe Leu Val Asp Asn Glu Thr Phe Ser Gly Phe Leu Tyr His
145 150 155 160

Asn Leu Ser Leu Pro Lys Ser Thr Val Asp Lys Met Leu Arg Ala Asp
165 170 175

Val Ile Leu His Lys Val Phe Leu Gln Gly Tyr Gln Leu His Leu Thr
180 185 190

Ser Leu Cys Asn Gly Ser Lys Ser Glu Glu Met Ile Gln Leu Gly Asp
195 200 205

Gln Glu Val Ser Glu Leu Cys Gly Leu Pro Lys Glu Lys Leu Ala Ala

210 215 220
Ala Glu Arg Val Leu Arg Ser Asn Met Asp Ile Leu Lys Pro Ile Leu
225 230 235 240
Arg Thr Leu Asn Ser Thr Ser Pro Phe Pro Ser Lys Glu Leu Ala Glu
245 250 255
Ala Thr Lys Thr Leu Leu His Ser Leu Gly Thr Leu Ala Gln Glu Leu
260 265 270
Phe Ser Met Arg Ser Trp Ser Asp Met Arg Gln Glu Val Met Phe Leu
275 280 285
Thr Asn Val Asn Ser Ser Ser Ser Ser Thr Gln Ile Tyr Gln Ala Val
290 295 300
Ser Arg Ile Val Cys Gly His Pro Glu Gly Gly Gly Leu Lys Ile Lys
305 310 315 320
Ser Leu Asn Trp Tyr Glu Asp Asn Asn Tyr Lys Ala Leu Phe Gly Gly
325 330 335
Asn Gly Thr Glu Glu Asp Ala Glu Thr Phe Tyr Asp Asn Ser Thr Thr
340 345 350
Pro Tyr Cys Asn Asp Leu Met Lys Asn Leu Glu Ser Ser Pro Leu Ser
355 360 365

Arg Ile Ile Trp Lys Ala Leu Lys Pro Leu Leu Val Gly Lys Ile Leu

370

375

380

Tyr Thr Pro Asp Thr Pro Ala Thr Arg Gln Val Met Ala Glu Val Asn

385

390

395

400

Lys Thr Phe Gln Glu Leu Ala Val Phe His Asp Leu Glu Gly Met Trp

405

410

415

Glu Glu Leu Ser Pro Lys Ile Trp Thr Phe Met Glu Asn Ser Gln Glu

420

425

430

Met Asp Leu Val Arg Met Leu Leu Asp Ser Arg Asp Asn Asp His Phe

435

440

445

Trp Glu Gln Gln Leu Asp Gly Leu Asp Trp Thr Ala Gln Asp Ile Val

450

455

460

Ala Phe Leu Ala Lys His Pro Glu Asp Val Gln Ser Ser Asn Gly Ser

465

470

475

480

Val Tyr Thr Trp Arg Glu Ala Phe Asn Glu Thr Asn Gln Ala Ile Arg

485

490

495

Thr Ile Ser Arg Phe Met Glu Cys Val Asn Leu Asn Lys Leu Glu Pro

500

505

510

Ile Ala Thr Glu Val Trp Leu Ile Asn Lys Ser Met Glu Leu Leu Asp

515	520	525
Glu Arg Lys Phe Trp Ala Gly Ile Val Phe Thr Gly Ile Thr Pro Gly		
530	535	540
Ser Ile Glu Leu Pro His His Val Lys Tyr Lys Ile Arg Met Asp Ile		
545	550	555
		560
Asp Asn Val Glu Arg Thr Asn Lys Ile Lys Asp Gly Tyr Trp Asp Pro		
565	570	575
Gly Pro Arg Ala Asp Pro Phe Glu Asp Met Arg Tyr Val Trp Gly Gly		
580	585	590
Phe Ala Tyr Leu Gln Asp Val Val Glu Gln Ala Ile Ile Arg Val Leu		
595	600	605
Thr Gly Thr Glu Lys Lys Thr Gly Val Tyr Met Gln Gln Met Pro Tyr		
610	615	620
Pro Cys Tyr Val Asp Asp Ile Phe Leu Arg Val Met Ser Arg Ser Met		
625	630	635
		640
Pro Leu Phe Met Thr Leu Ala Trp Ile Tyr Ser Val Ala Val Ile Ile		
645	650	655
Lys Gly Ile Val Tyr Glu Lys Glu Ala Arg Leu Lys Glu Thr Met Arg		
660	665	670

Ile Met Gly Leu Asp Asn Ser Ile Leu Trp Phe Ser Trp Phe Ile Ser
675 680 685

Ser Leu Ile Pro Leu Leu Val Ser Ala Gly Leu Leu Val Val Ile Leu
690 695 700

Lys Leu Gly Asn Leu Leu Pro Tyr Ser Asp Pro Ser Val Val Phe Val
705 710 715 720

Phe Leu Ser Val Phe Ala Val Val Thr Ile Leu Gln Cys Phe Leu Ile
725 730 735

Ser Thr Leu Phe Ser Arg Ala Asn Leu Ala Ala Ala Cys Gly Gly Ile
740 745 750

Ile Tyr Phe Thr Leu Tyr Leu Pro Tyr Val Leu Cys Val Ala Trp Gln
755 760 765

Asp Tyr Val Gly Phe Thr Leu Lys Ile Phe Ala Ser Leu Leu Ser Pro
770 775 780

Val Ala Phe Gly Phe Gly Cys Glu Tyr Phe Ala Leu Phe Glu Glu Gln
785 790 795 800

Gly Ile Gly Val Gln Trp Asp Asn Leu Phe Glu Ser Pro Val Glu Glu
805 810 815

Asp Gly Phe Asn Leu Thr Thr Ser Ile Ser Met Met Leu Phe Asp Thr

820	825	830
Phe Leu Tyr Gly Val Met Thr Trp Tyr Ile Glu Ala Val Phe Pro Gly		
835	840	845
Gln Tyr Gly Ile Pro Arg Pro Trp Tyr Phe Pro Cys Thr Lys Ser Tyr		
850	855	860
Trp Phe Gly Glu Glu Ser Asp Glu Lys Ser His Pro Gly Ser Asn Gln		
865	870	875
		880
Lys Arg Met Ser Glu Ile Cys Met Glu Glu Glu Pro Thr His Leu Lys		
885	890	895
Leu Gly Val Ser Ile Gln Asn Leu Val Lys Val Tyr Arg Asp Gly Met		
900	905	910
Lys Val Ala Val Asp Gly Leu Ala Leu Asn Phe Tyr Glu Gly Gln Ile		
915	920	925
Thr Ser Phe Leu Gly His Asn Gly Ala Gly Lys Thr Thr Thr Met Ser		
930	935	940
Ile Leu Thr Gly Leu Phe Pro Pro Thr Ser Gly Thr Ala Tyr Ile Leu		
945	950	955
		960
Gly Lys Asp Ile Arg Ser Glu Met Ser Thr Ile Arg Gln Asn Leu Gly		
965	970	975

Val Cys Pro Gln His Asn Val Leu Phe Asp Met Leu Thr Val Glu Glu
980 985 990

His Ile Trp Phe Tyr Ala Arg Leu Lys Gly Leu Ser Glu Lys His Val
995 1000 1005

Lys Ala Glu Met Glu Gln Met Ala Leu Asp Val Gly Leu Pro Ser Ser
1010 1015 1020

Lys Leu Lys Ser Lys Thr Ser Gln Leu Ser Gly Gly Met Gln Arg Lys
1025 1030 1035 1040

Leu Ser Val Ala Leu Ala Phe Val Gly Gly Ser Lys Val Val Ile Leu
1045 1050 1055

Asp Glu Pro Thr Ala Gly Val Asp Pro Tyr Ser Arg Arg Gly Ile Trp
1060 1065 1070

Glu Leu Leu Leu Lys Tyr Arg Gln Gly Arg Thr Ile Ile Leu Ser Thr
1075 1080 1085

His His Met Asp Glu Ala Asp Val Leu Gly Asp Arg Ile Ala Ile Ile
1090 1095 1100

Ser His Gly Lys Leu Cys Cys Val Gly Ser Ser Leu Phe Leu Lys Asn
1105 1110 1115 1120

Gln Leu Gly Thr Gly Tyr Tyr Leu Thr Leu Val Lys Lys Asp Val Glu

1125	1130	1135	
Ser Ser Leu Ser Ser Cys Arg Asn Ser Ser Ser Thr Val Ser Tyr Leu			
1140	1145	1150	
Lys Lys Glu Asp Ser Val Ser Gln Ser Ser Ser Asp Ala Gly Leu Gly			
1155	1160	1165	
Ser Asp His Glu Ser Asp Thr Leu Thr Ile Asp Val Ser Ala Ile Ser			
1170	1175	1180	
Asn Leu Ile Arg Lys His Val Ser Glu Ala Arg Leu Val Glu Asp Ile			
1185	1190	1195	1200
Gly His Glu Leu Thr Tyr Val Leu Pro Tyr Glu Ala Ala Lys Glu Gly			
1205	1210	1215	
Ala Phe Val Glu Leu Phe His Glu Ile Asp Asp Arg Leu Ser Asp Leu			
1220	1225	1230	
Gly Ile Ser Ser Tyr Gly Ile Ser Glu Thr Thr Leu Glu Glu Ile Phe			
1235	1240	1245	
Leu Lys Val Ala Glu Glu Ser Gly Val Asp Ala Glu Thr Ser Asp Gly			
1250	1255	1260	
Thr Leu Pro Ala Arg Arg Asn Arg Arg Ala Phe Gly Asp Lys Gln Ser			
1265	1270	1275	1280

Cys Leu Arg Pro Phe Thr Glu Asp Asp Ala Ala Asp Pro Asn Asp Ser
1285 1290 1295

Asp Ile Asp Pro Glu Ser Arg Glu Thr Asp Leu Leu Ser Gly Met Asp
1300 1305 1310

Gly Lys Gly Ser Tyr Gln Val Lys Gly Trp Lys Leu Thr Gln Gln Gln
1315 1320 1325

Phe Val Ala Leu Leu Trp Lys Arg Leu Leu Ile Ala Arg Arg Ser Arg
1330 1335 1340

Lys Gly Phe Phe Ala Gln Ile Val Leu Pro Ala Val Phe Val Cys Ile
1345 1350 1355 1360

Ala Leu Val Phe Ser Leu Ile Val Pro Pro Phe Gly Lys Tyr Pro Ser
1365 1370 1375

Leu Glu Leu Gln Pro Trp Met Tyr Asn Glu Gln Tyr Thr Phe Val Ser
1380 1385 1390

Asn Asp Ala Pro Glu Asp Thr Gly Thr Leu Glu Leu Leu Asn Ala Leu
1395 1400 1405

Thr Lys Asp Pro Gly Phe Gly Thr Arg Cys Met Glu Gly Asn Pro Ile
1410 1415 1420

Pro Asp Thr Pro Cys Gln Ala Gly Glu Glu Glu Trp Thr Thr Ala Pro

1425	1430	1435	1440
Val Pro Gln Thr Ile Met Asp Leu Phe Gln Asn Gly Asn Trp Thr Met			
1445	1450	1455	
Gln Asn Pro Ser Pro Ala Cys Gln Cys Ser Ser Asp Lys Ile Lys Lys			
1460	1465	1470	
Met Leu Pro Val Cys Pro Pro Gly Ala Gly Gly Leu Pro Pro Pro Gln			
1475	1480	1485	
Arg Lys Gln Asn Thr Ala Asp Ile Leu Gln Asp Leu Thr Gly Arg Asn			
1490	1495	1500	
Ile Ser Asp Tyr Leu Val Lys Thr Tyr Val Gln Ile Ile Ala Lys Ser			
1505	1510	1515	1520
Leu Lys Asn Lys Ile Trp Val Asn Glu Phe Arg Tyr Gly Gly Phe Ser			
1525	1530	1535	
Leu Gly Val Ser Asn Thr Gln Ala Leu Pro Pro Ser Gln Glu Val Asn			
1540	1545	1550	
Asp Ala Ile Lys Gln Met Lys Lys His Leu Lys Leu Ala Lys Asp Ser			
1555	1560	1565	
Ser Ala Asp Arg Phe Leu Asn Ser Leu Gly Arg Phe Met Thr Gly Leu			
1570	1575	1580	

Asp Thr Arg Asn Asn Val Lys Val Trp Phe Asn Asn Lys Gly Trp His
1585 1590 1595 1600

Ala Ile Ser Ser Phe Leu Asn Val Ile Asn Asn Ala Ile Leu Arg Ala
1605 1610 1615

Asn Leu Gln Lys Gly Glu Asn Pro Ser His Tyr Gly Ile Thr Ala Phe
1620 1625 1630

Asn His Pro Leu Asn Leu Thr Lys Gln Gln Leu Ser Glu Val Ala Leu
1635 1640 1645

Met Thr Thr Ser Val Asp Val Leu Val Ser Ile Cys Val Ile Phe Ala
1650 1655 1660

Met Ser Phe Val Pro Ala Ser Phe Val Val Phe Leu Ile Gln Glu Arg
1665 1670 1675 1680

Val Ser Lys Ala Lys His Leu Gln Phe Ile Ser Gly Val Lys Pro Val
1685 1690 1695

Ile Tyr Trp Leu Ser Asn Phe Val Trp Asp Met Cys Asn Tyr Val Val
1700 1705 1710

Pro Ala Thr Leu Val Ile Ile Ile Phe Ile Cys Phe Gln Gln Lys Ser
1715 1720 1725

Tyr Val Ser Ser Thr Asn Leu Pro Val Leu Ala Leu Leu Leu Leu

1730	1735	1740
Tyr Gly Trp Ser Ile Thr Pro Leu Met Tyr Pro Ala Ser Phe Val Phe		
1745	1750	1755
1760		
Lys Ile Pro Ser Thr Ala Tyr Val Val Leu Thr Ser Val Asn Leu Phe		
1765	1770	1775
Ile Gly Ile Asn Gly Ser Val Ala Thr Phe Val Leu Glu Leu Phe Thr		
1780	1785	1790
Asp Asn Lys Leu Asn Asn Ile Asn Asp Ile Leu Lys Ser Val Phe Leu		
1795	1800	1805
Ile Phe Pro His Phe Cys Leu Gly Arg Gly Leu Ile Asp Met Val Lys		
1810	1815	1820
Asn Gln Ala Met Ala Asp Ala Leu Glu Arg Phe Gly Glu Asn Arg Phe		
1825	1830	1835
1840		
Val Ser Pro Leu Ser Trp Asp Leu Val Gly Arg Asn Leu Phe Ala Met		
1845	1850	1855
Ala Val Glu Gly Val Val Phe Phe Leu Ile Thr Val Leu Ile Gln Tyr		
1860	1865	1870
Arg Phe Phe Ile Arg Pro Arg Pro Val Asn Ala Lys Leu Ser Pro Leu		
1875	1880	1885

Asn Asp Glu Asp Glu Asp Val Arg Arg Glu Arg Gln Arg Ile Leu Asp
1890 1895 1900

Gly Gly Gly Gln Asn Asp Ile Leu Glu Ile Lys Glu Leu Thr Lys Ile
1905 1910 1915 1920

Tyr Arg Arg Lys Arg Lys Pro Ala Val Asp Arg Ile Cys Val Gly Ile
1925 1930 1935

Pro Pro Gly Glu Cys Phe Gly Leu Leu Gly Val Asn Gly Ala Gly Lys
1940 1945 1950

Ser Ser Thr Phe Lys Met Leu Thr Gly Asp Thr Thr Val Thr Arg Gly
1955 1960 1965

Asp Ala Phe Leu Asn Lys Asn Ser Ile Leu Ser Asn Ile His Glu Val
1970 1975 1980

His Gln Asn Met Gly Tyr Cys Pro Gln Phe Asp Ala Ile Thr Glu Leu
1985 1990 1995 2000

Leu Thr Gly Arg Glu His Val Glu Phe Phe Ala Leu Leu Arg Gly Val
2005 2010 2015

Pro Glu Lys Glu Val Gly Lys Val Gly Glu Trp Ala Ile Arg Lys Leu
2020 2025 2030

Gly Leu Val Lys Tyr Gly Glu Lys Tyr Ala Gly Asn Tyr Ser Gly Gly

2035	2040	2045
Asn Lys Arg Lys Leu Ser Thr Ala Met Ala Leu Ile Gly Gly Pro Pro		
2050	2055	2060
Val Val Phe Leu Asp Glu Pro Thr Thr Gly Met Asp Pro Lys Ala Arg		
2065	2070	2075
Arg Phe Leu Trp Asn Cys Ala Leu Ser Val Val Lys Glu Gly Arg Ser		
2085	2090	2095
Val Val Leu Thr Ser His Ser Met Glu Glu Cys Glu Ala Leu Cys Thr		
2100	2105	2110
Arg Met Ala Ile Met Val Asn Gly Arg Phe Arg Cys Leu Gly Ser Val		
2115	2120	2125
Gln His Leu Lys Asn Arg Phe Gly Asp Gly Tyr Thr Ile Val Val Arg		
2130	2135	2140
Ile Ala Gly Ser Asn Pro Asp Leu Lys Pro Val Gln Asp Phe Phe Gly		
2145	2150	2155
Leu Ala Phe Pro Gly Ser Val Leu Lys Glu Lys His Arg Asn Met Leu		
2165	2170	2175
Gln Tyr Gln Leu Pro Ser Ser Leu Ser Ser Leu Ala Arg Ile Phe Ser		
2180	2185	2190

Ile Leu Ser Gln Ser Lys Lys Arg Leu His Ile Glu Asp Tyr Ser Val
 2195 2200 2205

Ser Gln Thr Thr Leu Asp Gln Val Phe Val Asn Phe Ala Lys Asp Gln
 2210 2215 2220

Ser Asp Asp Asp His Leu Lys Asp Leu Ser Leu His Lys Asn Gln Thr
 2225 2230 2235 2240

Val Val Asp Val Ala Val Leu Thr Ser Phe Leu Gln Asp Glu Lys Val
 2245 2250 2255

Lys Glu Ser Tyr Val
 2260

<210> 3

<211> 1643

<212> DNA

<213> Homo sapiens

<400> 3

gaattccttg ctggtggctc cacatgcact tccagggcct gcttggetct tctatgggtc 60
 tgtcctgagt gttgatagaa ccactgatgt gaggacctgg gcttgagcgt ggcctggaga 120
 tcctgttgac tgtagcatgg agggggcctg tcagctgaat gtctgtatgc aggtggtggg 180
 agttctggaa tatgatggag ctggaggtgg gaagagaagt aggcttgggg cagctctctc 240
 atgccacctc attctggcca aaactcaggt caaactgtga agagtctaaa tgtgaatctg 300
 cccttcaagg tggctacaaa ggtatctttg tcaaggtagg agaccttggt gcctccacgt 360

gcacttccag ggcctgcttg gcctcttcta cgggtctgtc ctgagtcttc tatgaatctc 420
 ccttcagggc agattcatat ttagactctt cacagtttga cctgagtttt ggccagaata 480
 aggtgacatt tagtttggtg gcttgatgaa tgacttaaatt atttagacat atgggtgtgta 540
 ggcctgcatt cctactcttg cctttttttt tgccctcca gtgttttggg tagttttgct 600
 cccctacag ccaaaggcaa acagataagt tggaggtctg gagtggctac ataattttac 660
 acgactgcaa ttctctggct gcacttcaca aatgtataca aactaaatac aagtcctgtg 720
 tttttatcac agggaggctg atcaatataa tgaaattaaa aggggggctg tcccatattg 780
 ttctgtgttt ttgtttgttt gtttcttttt ttgtttttgt ggctccttc ctctcaattt 840
 atgaagagaa gcagtaagat gttcctctcg ggtcctctga gggacctggg gagctcaggc 900
 tgggaatctc caaggcagta ggtcgcttat caaaaatcaa agtccagggt tgtgggggga 960
 aaacaaaagc agcccattac ccagaggact gtccgccttc cctcaccctc agcctaggcc 1020
 tttgaaagga aacaaaagac aagacaaaat gattggcgctc ctgagggaga ttcagcctag 1080
 agctctctct cccccaatcc ctccctccgg ctgaggaaac taacaaagga aaaaaaatt 1140
 gcggaagca ggatttagag gaagcaaatt ccactgggtgc ccttggctgc cgggaacgtg 1200
 gactagagag tctgcggcgc agccccgagc ccagcgcttc ccgcgcgtct taggccggcg 1260
 ggcccgggcg ggggaagggg acgcagaccg cggaccctaa gacacctgct gtaccctcca 1320
 cccccacccc acccacctcc ccccaactcc ctagatgtgt cgtgggcggc tgaacgtcgc 1380
 ccgtttaagg ggcgggcccc ggctccacgt gctttctgct gagtgaactga actacataaa 1440
 cagaggccgg gaacggggcg gggaggaggg agagcacagg ctttgaccga tagtaacctc 1500
 tgcgctcggg gcagccgaat ctataaaagg aactagtccc ggcaaaaacc ccgtaattgc 1560
 gagcgagagt gagtggggcc gggaccgca gagccgagcc gacccttctc tcccgggctg 1620
 cggcagggca gggcggggag ctc 1643

<210> 4

<211> 748

<212> DNA

<213> Homo sapiens

<400> 4

attccaanaa cattttccng catctgtggt tgccaactca caatgtcttt cattttctga 60
 acttaccenc caaatgaagn tcggcacgca attatgtagt agcnactctt agggccccgg 120
 cttacactta tgcttccggc tngttgtgtg ggaattggac ggataccatt tcacncagga 180
 aacagatatg nccatgatta cgccaagtta tttagggtgcn cgatagaata ctcaagcttg 240
 gaattcgcgg ccgcagtcga cggacccccg ggaaagattc ctctcattac acaaaaacca 300
 gacagtagtg gacgttgcag ttctcacatc ttctctacag gatgagaaag tgaaagaaag 360
 ctatgtatga agaatcctgt tcatacgggg tggctgaaag taaagaggaa ctagactttc 420
 ctttgcacca tgtgaagtgt tgtggagaaa agagccagaa gttgatgtgg gaagaagtaa 480
 actggatact gtactgatac tattcaatgc aatgcaattc aatgcaatga aaacaaaatt 540
 ccattacagg ggcagtgcct ttgtagccta tgtcttgtat ggctctcaag tgaaagactt 600
 gaatttagtt ttttacctat acctatgtga aactctatta tggaacccaa tggacatatg 660
 ggtttgaact cacacttttt ttttttgtt cctgtgtatt ctcatgggg ttgcaacaat 720
 aattcatcaa gtataaaaaa aaaaaaaa 748

<210> 5

<211> 2011

<212> DNA

<213> Homo sapiens

<400> 5

agaatcctgt tcatactggg gtggcttgaa agtaaattgga ggaactagac ttctcttgc 60
 accatgtgaa gtgttgtgga gaaaagagcc agaagttgat gtgggaagaa gtaaactgga 120
 tactgtactg atactattca atgcaatgca attcaatgca atgaaaacaa aattccatta 180
 caggggcaag tgcctttgta gcccatgtct tgtatggctc tcaagtgaaa gacttgaatt 240
 tagtttttta cctataccta tgtgaaactc tattatggaa cccaatggac atatgggttt 300
 gaactcacac tttttttttt tttgttcct gtgtattctc attgggggtg caacaataat 360
 tcatcaagta atcatggcca gcgattattg atcaaaatca aaaggtaatg cacatcctca 420
 ttcactaagc catgccatgc ccaggagact ggtttcccg tgacacatcc attgctggca 480

atgagtgatgc cagagttatt agtgccaagt ttttcagaaa gtttgaagca ccatgggtgtg 540
 tcatgctcac ttttgtgaaa gctgctctgc tcagagtcta tcaacattga atatcagttg 600
 acagaatggt gccatgctg gctaacaatcc tgctttgatt cctcttgata agctgttctg 660
 gtggcagtaa catgcaacaa aaatgtgggt gtctctaggg acgggaaact tggttccatt 720
 gttatattgt cctatgcttc gagccatggg tctacagggg catccttatg agactcttaa 780
 atatacttag atcctggtaa gaggcaaaga atcaacagcc aaactgctgg ggctgcaagc 840
 tgctgaagcc agggcatggg attaaagaga ttgtgcgttc aaacctaggg aagcctgtgc 900
 ccatttgtcc tgactgtctg ctaacatggg aactgcatc tcaagatgtt tatctgacac 960
 aagtgtatta tttctggctt tttgaattaa tctagaaaat gaaaagatgg agttgtatTT 1020
 tgacaaaaat gtttgtactt tttaatgtta tttggaattt taagttctat cagtgacttc 1080
 tgaatcctta gaatggcctc tttgtagaac cctgtggtat agaggagtat ggccactgcc 1140
 ccactatTTT tattttctta tgtaagtttg catatcagtc atgactagtg cctagaagc 1200
 aatgtgatgg tcaggatctc atgacattat atttgagttt ctttcagatc atttaggata 1260
 ctcttaatct cacttcatca atcaaatatt ttttgagtgt atgctgtagc tgaaagagta 1320
 tgtacgtacg tataagacta gagagatatt aagtctcagt acacttcctg tgccatgtta 1380
 ttcagctcac tggtttacia atataggttg tcttgtgggt gtaggagccc actgtaacaa 1440
 tattgggcag cttttttttt tttttttttt aattgcaaca atgcaaaagc caagaaagta 1500
 taagggtcac aagtctaaac aatgaattct tcaacagggg aaacagctag ctagaaaact 1560
 tgctgaaaac acaacttggt tttatggcat ttagtacctt caaataattg gctttgcaga 1620
 tattgggatac cccattaaat ctgacagtct caaatTTTt atctcttcaa tcactagtca 1680
 agaaaaatat aaaaacaaca aatacttcca tatggagcat ttttcagagt tttctaacc 1740
 agtcttattt ttctagtcag taaacatttg taaaaatact gtttcaactaa tacttactgt 1800
 taactgtctt gagagaaaag aaaaatatga gagaactatt gtttggggaa gttcaagtga 1860
 tctttcaata tcattactaa cttcttcac tttttccaaa atttgaatat taacgctaaa 1920
 ggtgtaagga cttcagattt caaattaatc tttctatatt ttttaaattt acagaatatt 1980
 atataacca ctgctgaaaa aaaaaaaaaa a 2011

<210> 6

<211> 3366

<212> DNA

<213> Homo sapiens

<400> 6

agaatcctgt tcatacgggg tggctgaaag taaagaggaa ctagactttc ctttgcacca 60
tgtgaagtgt tgtggagaaa agagccagaa gttgatgtgg gaagaagtaa actggatact 120
gtactgatac tattcaatgc aatgcaattc aatgcaatga aaacaaaatt ccattacagg 180
ggcagtgcct ttgtagccta tgtcttgtat ggctctcaag tgaaagactt gaatttagtt 240
ttttacctat acctatgtga aactctatta tggaacccaa tggacatatg ggtttgaact 300
cacacttttt tttttttttt gttcctgtgt attctcattg gggttgcaac aataattcat 360
caagtaatca tggccagcga ttattgatca aaatcaaaag gtaatgcaca tcttcattca 420
ctaagccatg ccatgcccag gagactgggt tcccgggtgac acatccattg ctggcaatga 480
gtgtgccaga gttattagt ccaagttttt cagaaagttt gaagcaccat ggtgtgtcat 540
gtcacttttt gtgaaagctg ctctgctcag agtctatcaa cattgaatat cagttgacag 600
aatggtgcc a tgcgtggcta acatcctgct ttgattccct ctgataagct gttctgggtg 660
cagtaacatg caacaaaaat gtgggtgtct ctaggcacgg gaaacttgggt tccattgtta 720
tattgtccta tgcttcgagc catgggtcta cagggtcac cttatgagac tcttaaata 780
acttagatcc tggtaagagg caaagaatca acagccaaac tgctggggct gcaagctgct 840
gaagccaggg catgggatta aagagattgt gcgttcaaac ctagggaagc ctgtgcccac 900
ttgtcctgac tgtctgctaa catggtacac tgcattctca gatgtttatc tgacacaagt 960
gtattatttc tggctttttg aattaatcta gaaaatgaaa agatggagtt gtattttgac 1020
aaaaatgttt gtacttttta atgttatttg gaattttaag ttctatcagt gacttctgaa 1080
tccttagaat ggctcttttg tagaaccctg tggatatagag gagtatggcc actgccccac 1140
tatttttatt ttcttatgta agtttgcata tcagtcatga ctagtgccta gaaagcaatg 1200
tgatggtcag gatctcatga cattatattt gagtttcttt cagatcattt aggatactct 1260
taatctcact tcatcaatca aatatttttt gagtgtatgc tgtagctgaa agagtatgta 1320
cgtacgtata agactagaga gatattaagt ctacgtacac ttctgtgccc atgttattca 1380
gtcactgggt ttacaaatat aggttgtctt gtgggtgtag gagccactg taacaatatt 1440

gggcagcctt tttttttttt ttttaattgc aacaatgcaa aagccaagaa agtataaggg 1500
tcacaagttt aaacaatgaa ttcttcaaca gggaaaacag ctagcttgaa aacttgctga 1560
aaaacacaac ttgtgtttat ggcatttagt accttcaa attggttt gcagatattg 1620
gatacccat taaatctgac agtctcaa ttttcatctc ttcaatcact agtcaagaaa 1680
aatataaaaa caacaatac ttccatattg agcatttttc agagttttct aacctagtct 1740
tatttttcta gtcagtaaac atttgtaaaa atactgtttc actaatactt actgttaact 1800
gtcttgagag aaaagaaaaa tatgagagaa ctattgtttg gggagttca agtgatcttt 1860
caatatcatt actaacttct tccacttttt ccaaaatttg aatattaacg ctaaagggtg 1920
aagacttcag atttcaaatt aatcttttcta tattttttta atttacagaa tattatataa 1980
cccactgctg aaaaagaaaa aaatgattgt tttagaagtt aaagtcaata ttgattttta 2040
atataagtaa tgaaggcata tttccaataa ctagtgatat ggcatcgttg caatttacag 2100
tatcttcaaa aatacagaat ttatagaata atttctctc atttaata tttcaaaatc 2160
aaagttatgg tttctcatt ttactaaaat cgtattctaa ttcttcatta tagtaaatct 2220
atgagcaact cttactctg gttctctga tttcaaggcc atatttttaa aaatcaaaag 2280
gcactgtgaa ctattttgaa gaaaacacga cattttaata cagattgaaa ggacctctc 2340
tgaagctaga aacaatctat agttatacat cttcattaat actgtgttac cttttaaaat 2400
agtaattttt tacattttcc tgtgtaaacc taattgtggt agaaattttt accaactcta 2460
tactcaatca agcaaaattt ctgtatatc cctgtggaat gtacctatgt gagtttcaga 2520
aattctcaaa atacgtgttc aaaaatttct gcttttgc atttgggaca cctcagaaaa 2580
cttattaaca actgtgaata tgagaaatac agaagaaaat aataagccct ctatacataa 2640
atgccagca caattcattg ttaaaaaaca accaaacctc acactactgt atttcattat 2700
ctgtactgaa agcaaatgct ttgtgactat taaatgttgc acatcattca ttcactgtat 2760
agtaatcatt gactaaagcc atttgctgtg ttttcttctt gtggnatgnat atatcaggta 2820
aaatattttc caaagagcca tgtgtcatgt aatactgaac ctttgatat tgagacatta 2880
atttggaccc ttggtattat ctactagaat aatgtaatac tgnagaaata ttgctcta 2940
tctttcaaaa tgggtcatcc cccttaaaan gttctatttc cataaggatt tagcttgctt 3000
atcccttctt ataccctaag atgaagctgt tttgtgtc tttgttcac attggccctc 3060
attccaagca ctttacgtg tctgtaatgg gatctatttt tgactggaa tatctgagaa 3120
ttgcaaaact agacaaaagt ttcacaacag atttctaagt taaatcattt tcattaaaa 3180

gaaaaaagaa aaaaaatttt gtatgtcaat aactttatat gaagtattaa aatgcatatt 3240
tctatgttgt aatataatga gtcacaaaat aaagctgtga cagttctgtt aaaaaaaaaa 3300
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3360
aaaaaa 3366

<210> 7

<211> 10474

<212> DNA

<213> Homo sapiens

<400> 7

tctagaactg ggtaccagct gctagcaagc ttcggcacga gccgcagagc cgagccgacc 60
cttctctccc gggctgcggc agggcagggc ggggagctcc gcgcaccaac agagccgggt 120
ctcagggcgc tttgctcctt gttttttccc cggttctgtt ttctcccctt ctccggaagg 180
cttgtcaagg ggtaggagaa agagacgcaa acacaaaagt ggaaaacagt taatgaccag 240
ccacgggcgt ccctgctgtg agctctggcc gctgccttcc agggctcccg agccacacgc 300
tgggcgtgct ggctgagggg acatggcttg ttggcctcag ctgagggtgc tgctgtggaa 360
gaacctcact ttcagaagaa gacaaacatg tcagctgtta ctggaagtgg cctggcctct 420
atttatcttc ctgatcctga tctctgttcg gctgagctac ccaccctatg aacaacatga 480
atgccatttt ccaataaaag ccatgccctc tgcaggaaca cttccttggg ttcaggggat 540
tatctgtaat gccacaacc cctgtttccg ttacccgact cctggggagg ctcccggagt 600
tgttgaaaac ttaacaaat ccattgtggc tcgcctgttc tcagatgctc ggaggcttct 660
tttatacagc cagaaagaca ccagcatgaa ggacatgcgc aaagttctga gaacattaca 720
gcagatcaag aaatccagct caaacttgaa gcttcaagat ttctgtgtgg acaatgaaac 780
cttctctggg ttcttatatc acaacctctc tctcccaaag tctactgtgg acaagatgct 840
gagggctgat gtcattctcc acaaggtatt tttgcaaggc taccagttac atttgacaag 900
tctgtgcaat ggatcaaaat cagaagagat gattcaactt ggtgaccaag aagtttctga 960
gctttgtggc ctaccaagg agaaactggc tgcagcagag cgagtacttc gttccaacat 1020

ggacatcctg aagccaatcc tgagaacact aaactctaca tctcccttcc cgagcaagga 1080
gctggctgaa gccacaaaaa cattgctgca tagtcttggg actctggccc aggagctgtt 1140
cagcatgaga agctggagtg acatgcgaca ggaggtgatg tttctgacca atgtgaacag 1200
ctccagctcc tccacccaaa tctaccaggc tgtgtctcgt attgtctgcg ggcattccga 1260
gggagggggg ctgaagatca agtctctcaa ctggtatgag gacaacaact acaaagccct 1320
ctttggaggc aatggcactg aggaagatgc tgaaaccttc tatgacaact ctacaactcc 1380
ttactgcaat gatttgatga agaatttgga gtctagtcct ctttcccgca ttatctggaa 1440
agctctgaag ccgctgctcg ttgggaagat cctgtatata cctgacactc cagccacaag 1500
gcaggtcatg gctgaggtga acaagacctt ccaggaactg gctgtgttcc atgatctgga 1560
aggcatgtgg gaggaactca gcccgaagat ctggaccttc atggagaaca gccaaagaaat 1620
ggaccttgtc cggatgctgt tggacagcag ggacaatgac cacttttggg aacagcagtt 1680
ggatggctta gattggacag cccaagacat cgtggcgttt ttggccaagc acccagagga 1740
tgtccagtcc agtaatggtt ctgtgtacac ctggagagaa gctttcaacg agactaacca 1800
ggcaatccgg accatatctc gcttcatgga gtgtgtcaac ctgaacaagc tagaaccat 1860
agcaacagaa gtctggctca tcaacaagtc catggagctg ctggatgaga ggaagtctctg 1920
ggctgggtatt gtgttcaactg gaattactcc aggagcatt gagctgcccc atcatgtcaa 1980
gtacaagatc cgaatggaca ttgacaatgt ggagaggaca aataaaatca aggatgggta 2040
ctgggaccct ggtcctcgag ctgacccctt tgaggacatg cggtagctct gggggggctt 2100
cgctacttg cgggatgtgg tggagcaggc aatcatcagg gtgctgacgg gcaccgagaa 2160
gaaaactggg gtctatatgc aacagatgcc ctatccctgt tacgttgatg acatctttct 2220
gcgggtgatg agccgggtcaa tgccctctt catgacgctg gcctggattt actcagtggc 2280
tgtgatcatc aagggcacg tgatatgaga ggaggcacgg ctgaaagaga ccatgcggat 2340
catgggcctg gacaacagca tactctgggt tagctgggtc attagtagcc tcattcctct 2400
tcttgtagc gctggcctgc tagtggtcat cctgaagtta ggaaacctgc tgccctacag 2460
tgatcccagc gtggtgtttg tcttctgtc cgtgtttgct gtggtgacaa tcctgcagt 2520
cttctgatt agcacactct tctccagagc caacctggca gcagcctgtg ggggcatcat 2580
ctacttcacg ctgtacctgc cctacgtcct gtgtgtggca tggcaggact acgtgggctt 2640
cacactcaag atcttcgcta gcctgctgtc tcctgtggct tttgggtttg gctgtgagta 2700
ctttgccctt tttgaggagc agggcattgg agtgcagtgg gacaacctgt ttgagagtcc 2760

tgtggaggaa gatggcttca atctcaccac ttcgatctcc atgatgctgt ttgacacctt 2820
cctctatggg gtgatgacct ggtacattga ggctgtcttt ccaggccagt acggaattcc 2880
caggccctgg tattttcctt gcaccaagtc ctactggttt ggcgaggaaa gtgatgagaa 2940
gagccaccct ggttccaacc agaagagaat gtcagaaatc tgcattggagg aggaaccac 3000
ccacttgaag ctgggcgtgt ccattcagaa cctggtaaaa gtctaccgag atgggatgaa 3060
ggtagctgtc gatggcctgg cactgaattt ttatgagggc cagatcacct ccttctggg 3120
ccacaatgga gcggggaaga cgaccaccat gtcaatcctg accgggttgt tcccccgac 3180
ctcgggcacc gcctacatcc tgggaaaaga cattcgtctc gagatgagca ccattccgga 3240
gaacctgggg gtctgtcccc agcataacgt gctgtttgac atgctgactg tcgaagaaca 3300
catctgggtc tatgcccgt tgaagggtct ctctgagaag cacgtgaagg cggagatgga 3360
gcagatggcc ctggatgttg gtttccatc aagcaagctg aaaagcaaaa caagccagct 3420
gtcaggtgga atgcagagaa agctatctgt ggccttggcc tttgtcgggg gatctaagg 3480
tgtcattctg gatgaacca cagctggtgt ggacccttac tcccgcaggg gaatatggga 3540
gctgctgctg aaataccgac aaggccgcac cattattctc tctacacacc acatggatga 3600
agcggacgtc ctgggggaca ggattgccat catctcccat gggaaagctgt gctgtgtggg 3660
ctcctccctg tttctgaaga accagctggg aacaggctac tacctgacct tggtaagaa 3720
agatgtggaa tcctccctca gttcctgcag aaacagtagt agcactgtgt catacctgaa 3780
aaaggaggac agtggttctc agagcagttc tgatgctggc ctgggcagcg accatgagag 3840
tgacacgctg accatcgatg tctctgctat ctccaacctc atcaggaagc atgtgtctga 3900
agccccgctg gtggaagaca tagggcatga gctgacctat gtgctgccat atgaagctgc 3960
taaggaggga gcctttgtgg aactctttca tgagattgat gaccggctct cagacctggg 4020
catttctagt tatggcatct cagagacgac cctggaagaa atattcctca aggtggccga 4080
agagagtggg gtggatgctg agacctcaga tggtagcttg ccagcaagac gaaacaggcg 4140
ggccttcggg gacaagcaga gctgtcttcg cccgttcact gaagatgatg ctgctgatcc 4200
aatgattct gacatagacc cagaatccag agagacagac ttgctcagtg ggatggatgg 4260
caaagggtcc taccaggtga aaggctggaa acttacacag caacagtttg tggccctttt 4320
gtggaagaga ctgctaattg ccagacggag tcggaaagga ttttttgctc agattgtctt 4380
gccagctgtg tttgtctgca ttgcccttgt gttcagcctg atcgtgccac cctttggcaa 4440

gtaccccagc ctggaacttc agccctggat gtacaacgaa cagtacacat ttgtcagcaa 4500
tgatgctcct gaggacacgg gaaccctgga actcttaaac gccctcacca aagaccctgg 4560
cttcgggacc cgctgtatgg aaggaaaccc aatcccagac acgccctgcc aggcagggga 4620
ggaagagtgg accactgccc cagttcccca gaccatcatg gacctcttcc agaatgggaa 4680
ctggacaatg cagaaccctt cacctgcatg ccagtgtagc agcgacaaaa tcaagaagat 4740
gctgcctgtg tgtccccag gggcaggggg gctgcctcct ccacaaagaa aacaaaacac 4800
tgcagatcct cttcaggacc tgacaggaag aaacatttcg gattatctgg tgaagacgta 4860
tgtgcagatc atagccaaaa gcttaaagaa caagatctgg gtgaatgagt ttaggtatgg 4920
cggcttttcc ctgggtgtca gtaatactca agcacttcct ccgagtcaag aagttaatga 4980
tgccatcaaa caaatgaaga aacacctaaa gctggccaag gacagttctg cagatcgatt 5040
tctcaacagc ttgggaagat ttatgacagg actggacacc agaaataatg tcaagggtgtg 5100
gttcaataac aagggctggc atgcaatcag ctctttcctg aatgtcatca acaatgccat 5160
tctccgggcc aacctgcaa aggagagaa ccctagccat tatggaatta ctgctttcaa 5220
tcatccccctg aatctcacca agcagcagct ctgagagggtg gctctgatga ccacatcagt 5280
ggatgtcctt gtgtccatct gtgtcatctt tgcaatgtcc ttcgtcccag ccagctttgt 5340
cgtattcctg atccaggagc gggtcagcaa agcaaaacac ctgcagttca tcagtggagt 5400
gaagcctgtc atctactggc tctctaattt tgtctgggat atgtgcaatt acgttggtccc 5460
tgccacactg gtcattatca tcttcatctg cttccagcag aagtcctatg tgtcctccac 5520
caatctgcct gtgctagccc ttctactttt gctgtatggg tggatcaatca cacctctcat 5580
gtacccagcc tcctttgtgt tcaagatccc cagcacagcc tatgtgggtg tcaccagcgt 5640
gaacctcttc attggcatta atggcagcgt ggccaccttt gtgctggagc tgttcaccga 5700
caataagctg aataatatca atgatatcct gaagtccgtg ttcttgatct tcccacattt 5760
ttgcctggga cgagggtcga tcgacatggt gaaaaaccag gcaatggctg atgccctgga 5820
aaggtttggg gagaatcgct ttgtgtcacc attatcttgg gacttgggtg gacgaaacct 5880
cttcgccatg gccgtggaag ggggtggtgtt cttcctcatt actgttctga tccagtacag 5940
attcttcatc aggccagac ctgtaaagtc aaagctatct cctctgaatg atgaagatga 6000
agatgtgagg cgggaaagac agagaattct tgatgggtgga ggccagaatg acatcttaga 6060
aatcaaggag ttgacgaaga tatatagaag gaagcggaag cctgctgttg acaggatttg 6120
cgtgggcatt cctcctgggtg agtgcttttg gctcctggga gttaatgggg ctggaaaatc 6180

atcaactttc aagatgttaa caggagatac cactgttacc agaggagatg ctttccttaa 6240
caaaaatagt atcttatcaa acatccatga agtacatcag aacatgggct actgccctca 6300
gtttgatgcc atcacagagc tgttgactgg gagagaacac gtggagtctt ttgccctttt 6360
gagaggagtc ccagagaaaag aagttggcaa ggttggtgag tgggcgattc ggaaactggg 6420
cctogtgaag tatggagaaa aatatgctgg taactatagt ggaggcaaca aacgcaagct 6480
ctctacagcc atggctttga tcggcggggc tcctgtggtg tttctggatg aaccaccac 6540
aggcatggat cccaaagccc ggcggttctt gtggaattgt gccctaagt tttgcaagga 6600
ggggagatca gtagtgctta catctcatag tatggaagaa tgtgaagctc tttgcactag 6660
gatggcaatc atgggtcaatg gaaggttcag gtgccttggc agtgtccagc atctaaaaaa 6720
taggtttgga gatggttata caatagttgt acgaatagca gggccaacc cggacctgaa 6780
gcctgtccag gatttctttg gacttgcatt tcctggaagt gttctaaaag agaaacaccg 6840
gaacatgcta caataccagc ttccatcttc attatcttct ctggccagga tattcagcat 6900
cctctcccag agcaaaaagc gactccacat agaagactac tctgtttctc agacaacact 6960
tgaccaagta tttgtgaact ttgccaagga ccaagtgat gatgaccact taaaagacct 7020
ctcattacac aaaaaccaga cagtagtgga cgttgcagtt ctcacatctt ttctacagga 7080
tgagaaagtg aaagaaagct atgtatgaag aatcctgttc atacgggggtg gctgaaagta 7140
aagaggaact agactttcct ttgcaccatg tgaagtgttg tggagaaaag agccagaagt 7200
tgatgtggga agaagtaaac tggatactgt actgatacta ttcaatgcaa tgcaattcaa 7260
tgcaatgaaa acaaaattcc attacagggg cagtgccttt gtagcctatg tcttgtatgg 7320
ctctcaagtg aaagacttga atttagtttt ttacctatac ctatgtgaaa ctctattatg 7380
gaaccaatg gacatatggg tttgaactca cacttttttt ttttttttgt tcctgtgtat 7440
tctcattggg gttgcaacaa taattcatca agtaatcatg gccagcgatt attgatcaaa 7500
atcaaaaggt aatgcacatc ctcattcact aagccatgcc atgcccagga gactggtttc 7560
ccggtgacac atccattgct ggcaatgagt gtgccagagt tattagtgcc aagtttttca 7620
gaaagtttga agcaccatgg tgtgtcatgc tcacttttgt gaaagctgct ctgctcagag 7680
tctatcaaca ttgaatatca gttgacagaa tggtgccatg cgtggctaac atcctgcttt 7740
gattccctct gataagctgt tctggtggca gtaacatgca acaaaaatgt ggggtgtctct 7800
aggcacggga aacttggttc cattgttata ttgtcctatg cttcgagcca tgggtctaca 7860

gggtcatcct tatgagactc ttaaataatac ttagatcctg gtaagaggca aagaatcaac 7920
agccaaactg ctggggctgc aagctgctga agccagggca tgggattaaa gagattgtgc 7980
gttcaaacct aggggaagcct gtgcccattt gtccctgactg tctgctaaca tggtagactg 8040
catctcaaga tgtttatctg acacaagtgt attatttctg gctttttgaa ttaatctaga 8100
aaatgaaaag atggagttgt attttgacaa aaatgtttgt actttttaat gttatttgga 8160
attttaagtt ctatcagtga cttctgaatc cttagaatgg cctctttgta gaacctgtg 8220
gtatagagga gtatggccac tgccccacta tttttatttt cttatgtaag tttgcatatc 8280
agtcattgact agtgcctaga aagcaatgtg atggctcagga tctcatgaca ttatatattga 8340
gtttctttca gatcatttag gataactctta atctcacttc atcaatcaaa tattttttga 8400
gtgtatgctg tagctgaaag agtatgtacg tacgtataag actagagaga tattaagtct 8460
cagtacactt cctgtgccat gttattcagc tcaactggtt acaaatatag gttgtcttgt 8520
ggttgtagga gccactgta acaatattgg gcagcctttt tttttttttt ttaattgcaa 8580
caatgcaaaa gccagaaaag tataagggtc acaagttaa acaatgaatt cttcaacagg 8640
gaaaacagct agcttgaaaa cttgctgaaa aacacaactt gtgtttatgg catttagtac 8700
cttcaaataa ttggctttgc agatattgga taccaccatta aatctgacag tctcaaattt 8760
ttcatctctt caatcactag tcaagaaaaa tataaaaaaca acaataactt ccatatggag 8820
catttttctag agttttctaa cccagtctta tttttctagt cagtaaacad ttgtaaaaat 8880
actgtttcac taatacttac tgtaactgt cttgagagaa aagaaaaata tgagagaact 8940
attgtttggg gaagttcaag tgatctttca atatcattac taacttcttc cactttttcc 9000
aaaatttgaa tattaacgct aaagggtgaa gacttcagat ttcaaattaa tctttctata 9060
ttttttaaat ttacagaata ttatataacc cactgctgaa aaagaaaaaa atgattgttt 9120
tagaagttaa agtcaatatt gatttttaaat ataagtaatg aaggcatatt tccaataact 9180
agtgatatgg catcgttgca atttacagta tcttcaaaaa tacagaattt atagaataat 9240
ttctcctcat ttaatatatt tcaaaatcaa agttatggtt tcctcatttt actaaaatcg 9300
tattctaatt cttcattata gtaaacttat gagcaactcc ttacttcggt tcctctgatt 9360
tcaaggccat attttaaaaa atcaaaaggc actgtgaact attttgaaga aaacacgaca 9420
ttttaataca gattgaaagg acctcttctg aagctagaaa caatctatag ttatacatct 9480
tcattaatac tgtgttacct tttaaaatag taatttttta cattttctctg tgtaaaccta 9540
attgtggtag aaatttttac caactctata ctcaatcaag caaaatttct gtatatcccc 9600

tgtggaatgt acctatgtga gtttcagaaa ttctcaaaat acgtgttcaa aaatttctgc 9660
 ttttgcattct ttgggacacc tcagaaaact tattaacaac tgtgaatatg agaaatacag 9720
 aagaaaataa taagccctct atacataaat gcccgacaca attcattggt aaaaaacaac 9780
 caaacctcac actactgtat ttcattatct gtactgaaag caaatgcttt gtgactatta 9840
 aatgttgacac atcattcatt cactgtatag taatcattga ctaaagccat ttgctgtggt 9900
 ttcttcttgt ggntgnatat atcaggtaaa atattttcca aagagccatg tgtcatgtaa 9960
 tactgaaccc ttgatattg agacattaat ttggaccctt ggtattatct actagaataa 10020
 tgtaatactg nagaaatatt gctctaattc tttcaaaatg gtgcatcccc cttaaaangt 10080
 tctattttcca taaggattta gcttgcttat cccttcttat accctaagat gaagctgttt 10140
 ttgtgctctt tggtcatcat tggccctcat tccaagcact ttacgctgtc tgtaatggga 10200
 tctatttttg cactggaata tctgagaatt gcaaaactag acaaaagttt cacaacagat 10260
 ttctaagtta aatcattttc attaaaagga aaaaagaaaa aaaattttgt atgtcaataa 10320
 ctttatatga agtattaaaa tgcataattc tatgttgtaa tataatgagt cacaaaaataa 10380
 agctgtgaca gttctgttaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 10440
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 10474

<210> 8

<211> 2261

<212> PRT

<213> Homo sapiens

<400> 8

Met Ala Cys Trp Pro Gln Leu Arg Leu Leu Leu Trp Lys Asn Leu Thr

1 , 5 10 15

Phe Arg Arg Arg Gln Thr Cys Gln Leu Leu Leu Glu Val Ala Trp Pro

20 25 30

Leu Phe Ile Phe Leu Ile Leu Ile Ser Val Arg Leu Ser Tyr Pro Pro
35 40 45

Tyr Glu Gln His Glu Cys His Phe Pro Asn Lys Ala Met Pro Ser Ala
50 55 60

Gly Thr Leu Pro Trp Val Gln Gly Ile Ile Cys Asn Ala Asn Asn Pro
65 70 75 80

Cys Phe Arg Tyr Pro Thr Pro Gly Glu Ala Pro Gly Val Val Gly Asn
85 90 95

Phe Asn Lys Ser Ile Val Ala Arg Leu Phe Ser Asp Ala Arg Arg Leu
100 105 110

Leu Leu Tyr Ser Gln Lys Asp Thr Ser Met Lys Asp Met Arg Lys Val
115 120 125

Leu Arg Thr Leu Gln Gln Ile Lys Lys Ser Ser Ser Asn Leu Lys Leu
130 135 140

Gln Asp Phe Leu Val Asp Asn Glu Thr Phe Ser Gly Phe Leu Tyr His
145 150 155 160

Asn Leu Ser Leu Pro Lys Ser Thr Val Asp Lys Met Leu Arg Ala Asp
165 170 175

Val Ile Leu His Lys Val Phe Leu Gln Gly Tyr Gln Leu His Leu Thr
180 185 190

Ser Leu Cys Asn Gly Ser Lys Ser Glu Glu Met Ile Gln Leu Gly Asp

195

200

205

Gln Glu Val Ser Glu Leu Cys Gly Leu Pro Lys Glu Lys Leu Ala Ala

210

215

220

Ala Glu Arg Val Leu Arg Ser Asn Met Asp Ile Leu Lys Pro Ile Leu

225

230

235

240

Arg Thr Leu Asn Ser Thr Ser Pro Phe Pro Ser Lys Glu Leu Ala Glu

245

250

255

Ala Thr Lys Thr Leu Leu His Ser Leu Gly Thr Leu Ala Gln Glu Leu

260

265

270

Phe Ser Met Arg Ser Trp Ser Asp Met Arg Gln Glu Val Met Phe Leu

275

280

285

Thr Asn Val Asn Ser Ser Ser Ser Thr Gln Ile Tyr Gln Ala Val

290

295

300

Ser Arg Ile Val Cys Gly His Pro Glu Gly Gly Gly Leu Lys Ile Lys

305

310

315

320

Ser Leu Asn Trp Tyr Glu Asp Asn Asn Tyr Lys Ala Leu Phe Gly Gly

325

330

335

Asn Gly Thr Glu Glu Asp Ala Glu Thr Phe Tyr Asp Asn Ser Thr Thr
340 345 350

Pro Tyr Cys Asn Asp Leu Met Lys Asn Leu Glu Ser Ser Pro Leu Ser
355 360 365

Arg Ile Ile Trp Lys Ala Leu Lys Pro Leu Leu Val Gly Lys Ile Leu
370 375 380

Tyr Thr Pro Asp Thr Pro Ala Thr Arg Gln Val Met Ala Glu Val Asn
385 390 395 400

Lys Thr Phe Gln Glu Leu Ala Val Phe His Asp Leu Glu Gly Met Trp
405 410 415

Glu Glu Leu Ser Pro Lys Ile Trp Thr Phe Met Glu Asn Ser Gln Glu
420 425 430

Met Asp Leu Val Arg Met Leu Leu Asp Ser Arg Asp Asn Asp His Phe
435 440 445

Trp Glu Gln Gln Leu Asp Gly Leu Asp Trp Thr Ala Gln Asp Ile Val
450 455 460

Ala Phe Leu Ala Lys His Pro Glu Asp Val Gln Ser Ser Asn Gly Ser
465 470 475 480

Val Tyr Thr Trp Arg Glu Ala Phe Asn Glu Thr Asn Gln Ala Ile Arg
485 490 495

Thr Ile Ser Arg Phe Met Glu Cys Val Asn Leu Asn Lys Leu Glu Pro

500

505

510

Ile Ala Thr Glu Val Trp Leu Ile Asn Lys Ser Met Glu Leu Leu Asp

515

520

525

Glu Arg Lys Phe Trp Ala Gly Ile Val Phe Thr Gly Ile Thr Pro Gly

530

535

540

Ser Ile Glu Leu Pro His His Val Lys Tyr Lys Ile Arg Met Asp Ile

545

550

555

560

Asp Asn Val Glu Arg Thr Asn Lys Ile Lys Asp Gly Tyr Trp Asp Pro

565

570

575

Gly Pro Arg Ala Asp Pro Phe Glu Asp Met Arg Tyr Val Trp Gly Gly

580

585

590

Phe Ala Tyr Leu Arg Asp Val Val Glu Gln Ala Ile Ile Arg Val Leu

595

600

605

Thr Gly Thr Glu Lys Lys Thr Gly Val Tyr Met Gln Gln Met Pro Tyr

610

615

620

Pro Cys Tyr Val Asp Asp Ile Phe Leu Arg Val Met Ser Arg Ser Met

625

630

635

640

Pro Leu Phe Met Thr Leu Ala Trp Ile Tyr Ser Val Ala Val Ile Ile

645

650

655

Lys Gly Ile Val Tyr Glu Lys Glu Ala Arg Leu Lys Glu Thr Met Arg

660

665

670

Ile Met Gly Leu Asp Asn Ser Ile Leu Trp Phe Ser Trp Phe Ile Ser

675

680

685

Ser Leu Ile Pro Leu Leu Val Ser Ala Gly Leu Leu Val Val Ile Leu

690

695

700

Lys Leu Gly Asn Leu Leu Pro Tyr Ser Asp Pro Ser Val Val Phe Val

705

710

715

720

Phe Leu Ser Val Phe Ala Val Val Thr Ile Leu Gln Cys Phe Leu Ile

725

730

735

Ser Thr Leu Phe Ser Arg Ala Asn Leu Ala Ala Ala Cys Gly Gly Ile

740

745

750

Ile Tyr Phe Thr Leu Tyr Leu Pro Tyr Val Leu Cys Val Ala Trp Gln

755

760

765

Asp Tyr Val Gly Phe Thr Leu Lys Ile Phe Ala Ser Leu Leu Ser Pro

770

775

780

Val Ala Phe Gly Phe Gly Cys Glu Tyr Phe Ala Leu Phe Glu Glu Gln

785

790

795

800

Gly Ile Gly Val Gln Trp Asp Asn Leu Phe Glu Ser Pro Val Glu Glu

805

810

815

Asp Gly Phe Asn Leu Thr Thr Ser Ile Ser Met Met Leu Phe Asp Thr

820

825

830

Phe Leu Tyr Gly Val Met Thr Trp Tyr Ile Glu Ala Val Phe Pro Gly

835

840

845

Gln Tyr Gly Ile Pro Arg Pro Trp Tyr Phe Pro Cys Thr Lys Ser Tyr

850

855

860

Trp Phe Gly Glu Glu Ser Asp Glu Lys Ser His Pro Gly Ser Asn Gln

865

870

875

880

Lys Arg Met Ser Glu Ile Cys Met Glu Glu Glu Pro Thr His Leu Lys

885

890

895

Leu Gly Val Ser Ile Gln Asn Leu Val Lys Val Tyr Arg Asp Gly Met

900

905

910

Lys Val Ala Val Asp Gly Leu Ala Leu Asn Phe Tyr Glu Gly Gln Ile

915

920

925

Thr Ser Phe Leu Gly His Asn Gly Ala Gly Lys Thr Thr Thr Met Ser

930

935

940

Ile Leu Thr Gly Leu Phe Pro Pro Thr Ser Gly Thr Ala Tyr Ile Leu
945 950 955 960

Gly Lys Asp Ile Arg Ser Glu Met Ser Thr Ile Arg Gln Asn Leu Gly
965 970 975

Val Cys Pro Gln His Asn Val Leu Phe Asp Met Leu Thr Val Glu Glu
980 985 990

His Ile Trp Phe Tyr Ala Arg Leu Lys Gly Leu Ser Glu Lys His Val
995 1000 1005

Lys Ala Glu Met Glu Gln Met Ala Leu Asp Val Gly Leu Pro Ser Ser
1010 1015 1020

Lys Leu Lys Ser Lys Thr Ser Gln Leu Ser Gly Gly Met Gln Arg Lys
1025 1030 1035 1040

Leu Ser Val Ala Leu Ala Phe Val Gly Gly Ser Lys Val Val Ile Leu
1045 1050 1055

Asp Glu Pro Thr Ala Gly Val Asp Pro Tyr Ser Arg Arg Gly Ile Trp
1060 1065 1070

Glu Leu Leu Leu Lys Tyr Arg Gln Gly Arg Thr Ile Ile Leu Ser Thr
1075 1080 1085

His His Met Asp Glu Ala Asp Val Leu Gly Asp Arg Ile Ala Ile Ile
1090 1095 1100

Ser His Gly Lys Leu Cys Cys Val Gly Ser Ser Leu Phe Leu Lys Asn

1105 1110 1115 1120

Gln Leu Gly Thr Gly Tyr Tyr Leu Thr Leu Val Lys Lys Asp Val Glu

 1125 1130 1135

Ser Ser Leu Ser Ser Cys Arg Asn Ser Ser Ser Thr Val Ser Tyr Leu

 1140 1145 1150

Lys Lys Glu Asp Ser Val Ser Gln Ser Ser Ser Asp Ala Gly Leu Gly

 1155 1160 1165

Ser Asp His Glu Ser Asp Thr Leu Thr Ile Asp Val Ser Ala Ile Ser

 1170 1175 1180

Asn Leu Ile Arg Lys His Val Ser Glu Ala Arg Leu Val Glu Asp Ile

1185 1190 1195 1200

Gly His Glu Leu Thr Tyr Val Leu Pro Tyr Glu Ala Ala Lys Glu Gly

 1205 1210 1215

Ala Phe Val Glu Leu Phe His Glu Ile Asp Asp Arg Leu Ser Asp Leu

 1220 1225 1230

Gly Ile Ser Ser Tyr Gly Ile Ser Glu Thr Thr Leu Glu Glu Ile Phe

 1235 1240 1245

Leu Lys Val Ala Glu Glu Ser Gly Val Asp Ala Glu Thr Ser Asp Gly

1250

1255

1260

Thr Leu Pro Ala Arg Arg Asn Arg Arg Ala Phe Gly Asp Lys Gln Ser

1265

1270

1275

1280

Cys Leu Arg Pro Phe Thr Glu Asp Asp Ala Ala Asp Pro Asn Asp Ser

1285

1290

1295

Asp Ile Asp Pro Glu Ser Arg Glu Thr Asp Leu Leu Ser Gly Met Asp

1300

1305

1310

Gly Lys Gly Ser Tyr Gln Val Lys Gly Trp Lys Leu Thr Gln Gln Gln

1315

1320

1325

Phe Val Ala Leu Leu Trp Lys Arg Leu Leu Ile Ala Arg Arg Ser Arg

1330

1335

1340

Lys Gly Phe Phe Ala Gln Ile Val Leu Pro Ala Val Phe Val Cys Ile

1345

1350

1355

1360

Ala Leu Val Phe Ser Leu Ile Val Pro Pro Phe Gly Lys Tyr Pro Ser

1365

1370

1375

Leu Glu Leu Gln Pro Trp Met Tyr Asn Glu Gln Tyr Thr Phe Val Ser

1380

1385

1390

Asn Asp Ala Pro Glu Asp Thr Gly Thr Leu Glu Leu Leu Asn Ala Leu

1395

1400

1405

Thr Lys Asp Pro Gly Phe Gly Thr Arg Cys Met Glu Gly Asn Pro Ile

1410

1415

1420

Pro Asp Thr Pro Cys Gln Ala Gly Glu Glu Glu Trp Thr Thr Ala Pro

1425

1430

1435

1440

Val Pro Gln Thr Ile Met Asp Leu Phe Gln Asn Gly Asn Trp Thr Met

1445

1450

1455

Gln Asn Pro Ser Pro Ala Cys Gln Cys Ser Ser Asp Lys Ile Lys Lys

1460

1465

1470

Met Leu Pro Val Cys Pro Pro Gly Ala Gly Gly Leu Pro Pro Pro Gln

1475

1480

1485

Arg Lys Gln Asn Thr Ala Asp Ile Leu Gln Asp Leu Thr Gly Arg Asn

1490

1495

1500

Ile Ser Asp Tyr Leu Val Lys Thr Tyr Val Gln Ile Ile Ala Lys Ser

1505

1510

1515

1520

Leu Lys Asn Lys Ile Trp Val Asn Glu Phe Arg Tyr Gly Gly Phe Ser

1525

1530

1535

Leu Gly Val Ser Asn Thr Gln Ala Leu Pro Pro Ser Gln Glu Val Asn

1540

1545

1550

Asp Ala Ile Lys Gln Met Lys Lys His Leu Lys Leu Ala Lys Asp Ser
1555 1560 1565

Ser Ala Asp Arg Phe Leu Asn Ser Leu Gly Arg Phe Met Thr Gly Leu
1570 1575 1580

Asp Thr Arg Asn Asn Val Lys Val Trp Phe Asn Asn Lys Gly Trp His
1585 1590 1595 1600

Ala Ile Ser Ser Phe Leu Asn Val Ile Asn Asn Ala Ile Leu Arg Ala
1605 1610 1615

Asn Leu Gln Lys Gly Glu Asn Pro Ser His Tyr Gly Ile Thr Ala Phe
1620 1625 1630

Asn His Pro Leu Asn Leu Thr Lys Gln Gln Leu Ser Glu Val Ala Leu
1635 1640 1645

Met Thr Thr Ser Val Asp Val Leu Val Ser Ile Cys Val Ile Phe Ala
1650 1655 1660

Met Ser Phe Val Pro Ala Ser Phe Val Val Phe Leu Ile Gln Glu Arg
1665 1670 1675 1680

Val Ser Lys Ala Lys His Leu Gln Phe Ile Ser Gly Val Lys Pro Val
1685 1690 1695

Ile Tyr Trp Leu Ser Asn Phe Val Trp Asp Met Cys Asn Tyr Val Val
1700 1705 1710

Pro Ala Thr Leu Val Ile Ile Ile Phe Ile Cys Phe Gln Gln Lys Ser

1715

1720

1725

Tyr Val Ser Ser Thr Asn Leu Pro Val Leu Ala Leu Leu Leu Leu

1730

1735

1740

Tyr Gly Trp Ser Ile Thr Pro Leu Met Tyr Pro Ala Ser Phe Val Phe

1745

1750

1755

1760

Lys Ile Pro Ser Thr Ala Tyr Val Val Leu Thr Ser Val Asn Leu Phe

1765

1770

1775

Ile Gly Ile Asn Gly Ser Val Ala Thr Phe Val Leu Glu Leu Phe Thr

1780

1785

1790

Asp Asn Lys Leu Asn Asn Ile Asn Asp Ile Leu Lys Ser Val Phe Leu

1795

1800

1805

Ile Phe Pro His Phe Cys Leu Gly Arg Gly Leu Ile Asp Met Val Lys

1810

1815

1820

Asn Gln Ala Met Ala Asp Ala Leu Glu Arg Phe Gly Glu Asn Arg Phe

1825

1830

1835

1840

Val Ser Pro Leu Ser Trp Asp Leu Val Gly Arg Asn Leu Phe Ala Met

1845

1850

1855

Ala Val Glu Gly Val Val Phe Phe Leu Ile Thr Val Leu Ile Gln Tyr

1860

1865

1870

Arg Phe Phe Ile Arg Pro Arg Pro Val Asn Ala Lys Leu Ser Pro Leu

1875

1880

1885

Asn Asp Glu Asp Glu Asp Val Arg Arg Glu Arg Gln Arg Ile Leu Asp

1890

1895

1900

Gly Gly Gly Gln Asn Asp Ile Leu Glu Ile Lys Glu Leu Thr Lys Ile

1905

1910

1915

1920

Tyr Arg Arg Lys Arg Lys Pro Ala Val Asp Arg Ile Cys Val Gly Ile

1925

1930

1935

Pro Pro Gly Glu Cys Phe Gly Leu Leu Gly Val Asn Gly Ala Gly Lys

1940

1945

1950

Ser Ser Thr Phe Lys Met Leu Thr Gly Asp Thr Thr Val Thr Arg Gly

1955

1960

1965

Asp Ala Phe Leu Asn Lys Asn Ser Ile Leu Ser Asn Ile His Glu Val

1970

1975

1980

His Gln Asn Met Gly Tyr Cys Pro Gln Phe Asp Ala Ile Thr Glu Leu

1985

1990

1995

2000

Leu Thr Gly Arg Glu His Val Glu Phe Phe Ala Leu Leu Arg Gly Val

2005

2010

2015

Pro Glu Lys Glu Val Gly Lys Val Gly Glu Trp Ala Ile Arg Lys Leu

2020

2025

2030

Gly Leu Val Lys Tyr Gly Glu Lys Tyr Ala Gly Asn Tyr Ser Gly Gly

2035

2040

2045

Asn Lys Arg Lys Leu Ser Thr Ala Met Ala Leu Ile Gly Gly Pro Pro

2050

2055

2060

Val Val Phe Leu Asp Glu Pro Thr Thr Gly Met Asp Pro Lys Ala Arg

2065

2070

2075

2080

Arg Phe Leu Trp Asn Cys Ala Leu Ser Val Val Lys Glu Gly Arg Ser

2085

2090

2095

Val Val Leu Thr Ser His Ser Met Glu Glu Cys Glu Ala Leu Cys Thr

2100

2105

2110

Arg Met Ala Ile Met Val Asn Gly Arg Phe Arg Cys Leu Gly Ser Val

2115

2120

2125

Gln His Leu Lys Asn Arg Phe Gly Asp Gly Tyr Thr Ile Val Val Arg

2130

2135

2140

Ile Ala Gly Ser Asn Pro Asp Leu Lys Pro Val Gln Asp Phe Phe Gly

2145

2150

2155

2160

Leu Ala Phe Pro Gly Ser Val Leu Lys Glu Lys His Arg Asn Met Leu
2165 2170 2175

Gln Tyr Gln Leu Pro Ser Ser Leu Ser Ser Leu Ala Arg Ile Phe Ser
2180 2185 2190

Ile Leu Ser Gln Ser Lys Lys Arg Leu His Ile Glu Asp Tyr Ser Val
2195 2200 2205

Ser Gln Thr Thr Leu Asp Gln Val Phe Val Asn Phe Ala Lys Asp Gln
2210 2215 2220

Ser Asp Asp Asp His Leu Lys Asp Leu Ser Leu His Lys Asn Gln Thr
2225 2230 2235 2240

Val Val Asp Val Ala Val Leu Thr Ser Phe Leu Gln Asp Glu Lys Val
2245 2250 2255

Lys Glu Ser Tyr Val
2260

<210> 9

<211> 10474

<212> DNA

<213> Homo sapiens

<400> 9

tctagaactg ggtaccagct gctagcaagc ttcggcacga gccgcagagc cgagccgacc 60

cttctctccc gggctgcggc agggcagggc ggggagctcc gcgcaccaac agagccgggtt 120
ctcagggcgc tttgctcctt gttttttccc cggttctgtt ttctccctt ctccggaagg 180
cttgtcaagg ggtaggagaa agagacgcaa acacaaaagt ggaaaacagt taatgaccag 240
ccacgggcgt ccttgcgtgtg agctctggcc gctgccttcc agggctcccc agccacacgc 300
tgggcgtgct ggctgagggg acatggcttg ttggcctcag ctgaggttgc tgctgtggaa 360
gaacctcact ttcagaagaa gacaaacatg tcagctgtta ctggaagtgg cctggcctct 420
atttatcttc ctgatcctga tctctgttcg gctgagctac ccaccctatg aacaacatga 480
atgccatfff ccaaataaag ccatgccctc tgcaggaaca cttccttggg ttcaggggat 540
tatctgtaat gccaacaacc cctgtttccg ttaccctgact cctggggagg ctcccggagt 600
tggttgaaac ttaacaaat ccattgtggc tcgcctgttc tcagatgctc ggaggcttct 660
tttatacagc cagaaagaca ccagcatgaa ggacatgcgc aaagtcttga gaacattaca 720
gcagatcaag aaatccagct caaacttgaa gcttcaagat ttcctgggtg acaatgaaac 780
cttctctggg ttcctatata acaacctctc tctcccaaag tctactgtgg acaagatgct 840
gagggctgat gtcattctcc acaaggtatt tttgcaaggc taccagttac atttgacaag 900
tctgtgcaat ggatcaaaat cagaagagat gattcaactt ggtgaccaag aagtttctga 960
gctttgtggc ctaccaaagg agaaactggc tgcagcagag cgagtacttc gttccaacat 1020
ggacatcctg aagccaatcc tgagaacact aaactctaca tctcccttcc cgagcaaggga 1080
gctggctgaa gccacaaaaa cattgctgca tagtcttggg actctggccc aggagctgtt 1140
cagcatgaga agctggagtg acatgcgaca ggaggtgatg tttctgacca atgtgaacag 1200
ctccagctcc tccacccaaa tctaccaggc tgtgtctcgt attgtctgcg ggcacccga 1260
gggagggggg ctgaagatca agtctctcaa ctggtatgag gacaacaact acaaagccct 1320
ctttggaggc aatggcactg aggaagatgc tgaaaccttc tatgacaact ctacaactcc 1380
ttactgcaat gatttgatga agaatttgga gtctagtcct ctttcccgca ttatctggaa 1440
agctctgaag ccgctgctcg ttgggaagat cctgtatata cctgacactc cagccacaag 1500
gcaggtcatg gctgaggtga acaagacctt ccaggaactg gctgtgttcc atgatctgga 1560
aggcatgtgg gaggaactca gcccgaagat ctggaccttc atggagaaca gccaaagaaat 1620
ggaccttgct cggtatgctgt tggacagcag ggacaatgac cacttttggg aacagcagtt 1680
ggatggctta gattggacag cccaagacat cgtggcgttt ttggccaagc acccagagga 1740

tgtccagtcc agtaatgggt ctgtgtacac ctggagagaa gctttcaacg agactaacca 1800
ggcaatccgg accatatctc gcttcatgga gtgtgtcaac ctgaacaagc tagaaccat 1860
agcaacagaa gtctggctca tcaacaagtc catggagctg ctggatgaga ggaagttctg 1920
ggctgggtatt gtgttctactg gaattactcc aggcagcatt gagctgcccc atcatgtcaa 1980
gtacaagatc cgaatggaca ttgacaatgt ggagaggaca aataaaatca aggatgggta 2040
ctgggaccct ggctctcgag ctgaccctt tgaggacatg tggtagctct gggggggctt 2100
cgctacttg caggatgtgg tggagcaggc aatcatcagg gtgctgacgg gcaccgagaa 2160
gaaaactggg gtctatatgc aacagatgcc ctatccctgt tacgttgatg acatctttct 2220
gcgggtgatg agccgggtcaa tgccctctt catgacgctg gcctggattt actcagtggc 2280
tgtgatcatc aagggcatcg tgtatgagaa ggaggcacgg ctgaaagaga ccatgcggat 2340
catgggcctg gacaacagca tactctgggt tagctgggtc attagtagcc tcattctctt 2400
tcttgtagc gctggcctgc tagtgggtcat cctgaagta ggaaacctgc tgccctacag 2460
tgatcccagc gtgggtgttg tcttctgtc cgtgtttgct gtggtgacaa tcctgcagt 2520
cttctgatt agcacactct tctccagagc caacctggca gcagcctgtg ggggcatcat 2580
ctacttcacg ctgtacctgc cctacgtcct gtgtgtggca tggcaggact acgtgggctt 2640
cacactcaag atcttcgcta gcctgctgtc tcctgtggct tttgggtttg gctgtgagta 2700
ctttgccctt tttgaggagc agggcattgg agtgcagtgg gacaacctgt ttgagagtcc 2760
tgtggaggaa gatggcttca atctcaccac ttcgatctcc atgatgctgt ttgacacctt 2820
cctctatggg gtgatgacct ggtacattga ggctgtctt ccaggccagt acggaattcc 2880
caggccctgg tattttctt gcaccaagtc ctactggtt ggcgaggaaa gtgatgagaa 2940
gagccacctt ggttccaacc agaagagaat gtcagaaatc tgcattggagg aggaaccac 3000
ccacttgaag ctgggcgtgt ccattcagaa cctggtaaaa gtctaccgag atgggatgaa 3060
gggtggctgt gatggcctgg cactgaattt ttatgagggc cagatcacct cttctctggg 3120
ccacaatgga gcggggaaga cgaccacat gtcaatctg accgggttgt tcccccgac 3180
ctcgggcacc gcctacatcc tgggaaaaga cattcgctct gagatgagca ccatccggca 3240
gaacctgggg gtctgtcccc agcataacgt gctgtttgac atgctgactg tcgaagaaca 3300
catctgggtc tatgcccgt tgaaagggt ctctgagaag cacgtgaagg cggagatgga 3360
gcagatggcc ctggatgttg gtttccatc aagcaagctg aaaagcaaaa caagccagct 3420
gtcaggtgga atgcagagaa agctatctgt ggccttggcc tttgtcggg gatctaaggt 3480

tgtcattctg gatgaaccca cagctggtgt ggacccttac tcccgaggg gaatatggga 3540
gctgctgctg aaataccgac aaggccgcac cattattctc tctacacacc acatggatga 3600
agcggacgctc ctgggggaca ggattgccat catctcccat gggaagctgt gctgtgtggg 3660
ctcctccctg tttctgaaga accagctggg aacaggctac tacctgacct tggtaagaa 3720
agatgtggaa tcctccctca gttctgcag aaacagtagt agcactgtgt catacctgaa 3780
aaaggaggac agtggtttctc agagcagttc tgatgctggc ctgggcagcg accatgagag 3840
tgacacgctg accatcgatg tctctgctat ctccaacctc atcaggaagc atgtgtctga 3900
agcccggctg gtggaagaca tagggcatga gctgacctat gtgctgccat atgaagctgc 3960
taaggaggga gcctttgtgg aactctttca tgagattgat gaccggctct cagacctggg 4020
catttctagt tatggcatct cagagacgac cctggaagaa atattcctca aggtggccga 4080
agagagtggg gtggatgctg agacctcaga tggtagcttg ccagcaagac gaaacaggcg 4140
ggccttcggg gacaagcaga gctgtcttcg cccgttcact gaagatgatg ctgctgatcc 4200
aaatgattct gacatagacc cagaatccag agagacagac ttgctcagtg ggatggatgg 4260
caaagggtcc taccaggtga aaggctggaa acttacacag caacagtttg tggccctttt 4320
gtggaagaga ctgctaattg ccagacggag tcggaaagga ttttttctc agattgtctt 4380
gccagctgtg tttgtctgca ttgcccttgt gttcagcctg atcgtgccac cctttggcaa 4440
gtacccacgc ctggaacttc agccctggat gtacaacgaa cagtacacat ttgtcagcaa 4500
tgatgtcctt gaggacacgg gaacctgga actcttaaac gccctacca aagacctggg 4560
cttcgggacc cgctgtatgg aaggaaacct aatcccagac acgccctgcc aggcagggga 4620
ggaagagtgg accactgccc cagttcccca gaccatcatg gacctcttcc agaatgggaa 4680
ctggacaatg cagaacctt cacctgcatg ccagtgtagc agcgacaaa tcaagaagat 4740
gctgcctgtg tgtccccag gggcagggg gctgcctcct ccacaaagaa aacaaaacac 4800
tgacagatgc cttcaggacc tgacaggaag aaacatttcg gattatctgg tgaagacgta 4860
tgtgcagatc atagccaaaa gcttaaagaa caagatctgg gtgaatgagt ttaggtatgg 4920
cggcttttcc ctgggtgtca gtaatactca agcacttcct ccgagtcaag aagttaatga 4980
tgccatcaaa caaatgaaga aacacctaaa gctggccaag gacagttctg cagatcgatt 5040
tctcaacagc ttgggaagat ttatgacagg actggacacc agaaataatg tcaaggtgtg 5100
gttcaataac aagggtggc atgcaatcag ctctttcctg aatgtcatca acaatgccat 5160

tctccgggcc aacctgcaaa agggagagaa ccctagccat tatggaatta ctgctttcaa 5220
tcaccccctg aatctcacca agcagcagct ctcagagggtg gctctgatga ccacatcagt 5280
ggatgtcctt gtgtccatct gtgtcatctt tgcaatgtcc ttcgtcccag ccagctttgt 5340
cgtattcctg atccaggagc gggtcagcaa agcaaaacac ctgcagttca tcagtggagt 5400
gaagcctgtc atctactggc tctctaattt tgtctgggat atgtgcaatt acgttgctcc 5460
tgccacactg gtcattatca tttcatctg cttccagcag aagtcctatg tgcctccac 5520
caatctgcct gtgctagccc ttctactttt gctgtatggg tggcfaatca cacctctcat 5580
gtaccagacc tccttttgtt tcaagatccc cagcacagcc tatgtgggtg tcaccagcgt 5640
gaacctcttc attggcatta atggcagcgt ggccaccttt gtgctggagc tgttcaccga 5700
caataagctg aataatatca atgatatcct gaagtccgtg ttcttgatct tcccacattt 5760
ttgcctggga cgagggctca tcgacatggt gaaaaaccag gcaatggctg atgccctgga 5820
aaggtttggg gagaatcgct ttgtgtcacc attatcttgg gacttggtgg gacgaaacct 5880
cttcgccatg gccgtggaag ggggtggtgtt ctctctcatt actgttctga tccagtacag 5940
attcttcatc aggccagac ctgtaaatgc aaagctatct cctctgaatg atgaagatga 6000
agatgtgagg cgggaaagac agagaattct tgatggtgga ggccagaatg acatcttaga 6060
aatcaaggag ttgacgaaga tatatagaag gaagcggaag cctgctgttg acaggatttg 6120
cgtgggcatt cctcctggtg agtgctttgg gctcctggga gttaatgggg ctggaaaatc 6180
atcaactttc aagatgttaa caggagatac cactgttacc agaggagatg ctttcttaa 6240
caaaaatagt atcttatcaa acatccatga agtacatcag aacatgggct actgcctca 6300
gtttgatgcc atcacagagc tgttgactgg gagagaacac gtggagtctt ttgcctttt 6360
gagaggagtc ccagagaaaag aagttggcaa ggttggtgag tgggcgattc ggaaactggg 6420
cctcgtgaag tatggagaaa aatatgctgg taactatagt ggaggcaaca aacgcaagct 6480
ctctacagcc atggctttga tcggcgggcc tcctgtggtg tttctggatg aaccaccac 6540
aggcatggat ccaaagccc ggcggttctt gtggaattgt gccctaagtg ttgtcaagga 6600
ggggagatca gtagtgctta catctcatag tatggaagaa tgtgaagctc tttgactag 6660
gatggcaatc atggtcaatg gaaggttcag gtgccttggc agtgtccagc atctaaaaa 6720
taggtttgga gatgggtata caatagttgt acgaatagca gggccaacc cggaacctgaa 6780
gcctgtccag gatttctttg gacttgcatt tcctggaagt gttctaaaag agaaacaccg 6840
gaacatgcta caataccagc ttcatcttc attatcttct ctggccagga tattcagcat 6900

cctctcccag agcaaaaagc gactccacat agaagactac tctgtttctc agacaacact 6960
tgaccaagta tttgtgaact ttgccaagga ccaaagtgat gatgaccact taaaagacct 7020
ctcattacac aaaaaccaga cagtagtgga cgttgacagtt ctcacatctt ttctacagga 7080
tgagaaagtg aaagaaagct atgtatgaag aatcctgttc atacgggggtg gctgaaagta 7140
aagaggaact agactttcct ttgcaccatg tgaagtgttg tggagaaaag agccagaagt 7200
tgatgtggga agaagtaaac tggatactgt actgatacta ttcaatgcaa tgcaattcaa 7260
tgcaatgaaa acaaaattcc attacagggg cagtgccttt gtagcctatg tcttgtatgg 7320
ctctcaagtg aaagacttga atttagtttt ttacctatac ctatgtgaaa ctctattatg 7380
gaacccaatg gacatatggg tttgaactca cacttttttt ttttttttgt tcttgtgtat 7440
tctcattggg gttgcaacaa taattcatca agtaatcatg gccagcgatt attgatcaaa 7500
atcaaaaggt aatgcacatc ctcattcact aagccatgcc atgcccagga gactggtttc 7560
ccggtgacac atccattgct ggcaatgagt gtgccagagt tattagtgcc aagtttttca 7620
gaaagtttga agcaccatgg tgtgtcatgc tcacttttgt gaaagctgct ctgctcagag 7680
tctatcaaca ttgaatatca gttgacagaa tgggtgccatg cgtgggctaac atcctgcttt 7740
gattccctct gataagctgt tctgggtggca gtaacatgca acaaaaatgt ggggtgtctct 7800
aggcacggga aacttggttc cattgttata ttgtcctatg cttcgagcca tgggtctaca 7860
gggtcatcct tatgagactc ttaaataac ttagatcctg gtaagaggca aagaatcaac 7920
agccaaactg ctggggctgc aagctgctga agccaggga tgggattaaa gagattgtgc 7980
gttcaaacct agggaagcct gtgcccattt gtcttgactg tctgctaaca tggtagactg 8040
catctcaaga tgtttatctg acacaagtgt attatttctg gctttttgaa ttaatctaga 8100
aaatgaaaag atggagtgtg attttgacaa aaatgtttgt actttttaat gttatttggg 8160
attttaagtt ctatcagtga cttctgaatc cttagaatgg cctctttgta gaaccctgtg 8220
gtatagagga gtatggccac tgcccacta tttttatttt cttatgtaag ttgcatatc 8280
agtcatgact agtgccatga aagcaatgtg atggtcagga tctcatgaca ttatatttga 8340
gtttctttca gatcatttag gatactctta atctcacttc atcaatcaaa tattttttga 8400
gtgtatgctg tagctgaaag agtatgtacg tacgtataag actagagaga tattaagtct 8460
cagtacactt cctgtgccat gttattcagc tcaactggtt acaaatatag gttgtcttgt 8520
ggttgtagga gcccaactga acaatattgg gcagcctttt tttttttttt ttaattgcaa 8580

caatgcaaaa gccaaagaaag tataaggggtc acaagtttaa acaatgaatt cttcaacagg 8640
gaaaacagct agcttgaaaa cttgctgaaa aacacaactt gtgtttatgg catttagtac 8700
cttcaaataa ttggctttgc agatattgga taccacatta aatctgacag tctcaaattt 8760
ttcatctctt caatcactag tcaagaaaaa tataaaaaaca acaataactt ccatatggag 8820
cattttttcag agtttttctaa cccagtctta tttttctagt cagtaaacad ttgtaaaaat 8880
actgtttcac taatacttac tgttaactgt cttgagagaa aagaaaaata tgagagaact 8940
attgtttggg gaagttcaag tgatctttca atatcattac taacttcttc cactttttcc 9000
aaaatttgaa tattaacgct aaaggtgtaa gacttcagat ttcaaattaa tctttctata 9060
ttttttaaat ttacagaata ttatataacc cactgctgaa aaagaaaaaa atgattgttt 9120
tagaagttaa agtcaatatt gatttttaaat ataagtaatg aaggcatatt tccaataact 9180
agtgatatgg catcgttgca atttacagta tcttcaaaaa tacagaattt atagaataat 9240
ttctcctcat ttaatatatt tcaaaatcaa agttatgggt tcttcatttt actaaaatcg 9300
tattctaatt cttcattata gtaaattctat gagcaactcc ttacttcggt tctcttgatt 9360
tcaaggccat attttaaaaa atcaaaaggc actgtgaact attttgaaga aaacacgaca 9420
ttttaataca gattgaaagg acctcttctg aagctagaaa caatctatag ttatacatct 9480
tcattaatac tgtgttacct tttaaaatag taatttttta catttttctg tgtaaaccta 9540
attgtggtag aaatttttac caactctata ctcaatcaag caaaatttct gtatatccc 9600
tgtggaatgt acctatgtga gtttcagaaa ttctcaaaat acgtgttcaa aaatttctgc 9660
ttttgcatct ttgggacacc tcagaaaact tattaacaac tgtgaatatg agaaatacag 9720
aagaaaataa taagccctct atacataaat gccagcaca attcattgtt aaaaaacaac 9780
caaacctcac actactgtat ttcattatct gtactgaaag caaatgcttt gtgactatta 9840
aatgttgac atcattcatt cactgtatag taatcattga cttaaagccat ttgctgtgtt 9900
ttcttcttgt ggntgnatat atcaggtaaa atattttcca aagagccatg tgtcatgtaa 9960
tactgaaccc ttgatattg agacattaat ttggaccctt ggtattatct actagaataa 10020
tgtaatactg nagaaatatt gctctaattc tttcaaaatg gtgcatcccc cttaaaangt 10080
tctattttcca taaggattta gcttgcttat ccttcttat accctaagat gaagctgttt 10140
ttgtgctctt tgttcattcat tggccctcat tccaagcact ttacgctgtc tgtaatggga 10200
tctatttttg cactggaata tctgagaatt gcaaaactag acaaaagttt cacaacagat 10260
ttctaagtta aatcattttc attaaaagga aaaaagaaaa aaaattttgt atgtcaataa 10320

ctttatatga agtattaaaa tgcataatttc tatgttgtaa tataatgagt cacaaaaataa 10380
agctgtgaca gttctgttaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 10440
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 10474

<210> 10

<211> 2261

<212> PRT

<213> Homo sapiens

<400> 10

Met Ala Cys Trp Pro Gln Leu Arg Leu Leu Leu Trp Lys Asn Leu Thr
1 5 10 15

Phe Arg Arg Arg Gln Thr Cys Gln Leu Leu Leu Glu Val Ala Trp Pro
20 25 30

Leu Phe Ile Phe Leu Ile Leu Ile Ser Val Arg Leu Ser Tyr Pro Pro
35 40 45

Tyr Glu Gln His Glu Cys His Phe Pro Asn Lys Ala Met Pro Ser Ala
50 55 60

Gly Thr Leu Pro Trp Val Gln Gly Ile Ile Cys Asn Ala Asn Asn Pro
65 70 75 80

Cys Phe Arg Tyr Pro Thr Pro Gly Glu Ala Pro Gly Val Val Gly Asn
85 90 95

Phe Asn Lys Ser Ile Val Ala Arg Leu Phe Ser Asp Ala Arg Arg Leu
100 105 110

Leu Leu Tyr Ser Gln Lys Asp Thr Ser Met Lys Asp Met Arg Lys Val
115 120 125

Leu Arg Thr Leu Gln Gln Ile Lys Lys Ser Ser Ser Asn Leu Lys Leu
130 135 140

Gln Asp Phe Leu Val Asp Asn Glu Thr Phe Ser Gly Phe Leu Tyr His
145 150 155 160

Asn Leu Ser Leu Pro Lys Ser Thr Val Asp Lys Met Leu Arg Ala Asp
165 170 175

Val Ile Leu His Lys Val Phe Leu Gln Gly Tyr Gln Leu His Leu Thr
180 185 190

Ser Leu Cys Asn Gly Ser Lys Ser Glu Glu Met Ile Gln Leu Gly Asp
195 200 205

Gln Glu Val Ser Glu Leu Cys Gly Leu Pro Lys Glu Lys Leu Ala Ala
210 215 220

Ala Glu Arg Val Leu Arg Ser Asn Met Asp Ile Leu Lys Pro Ile Leu
225 230 235 240

Arg Thr Leu Asn Ser Thr Ser Pro Phe Pro Ser Lys Glu Leu Ala Glu
245 250 255

Ala Thr Lys Thr Leu Leu His Ser Leu Gly Thr Leu Ala Gln Glu Leu

260

265

270

Phe Ser Met Arg Ser Trp Ser Asp Met Arg Gln Glu Val Met Phe Leu

275

280

285

Thr Asn Val Asn Ser Ser Ser Ser Ser Thr Gln Ile Tyr Gln Ala Val

290

295

300

Ser Arg Ile Val Cys Gly His Pro Glu Gly Gly Gly Leu Lys Ile Lys

305

310

315

320

Ser Leu Asn Trp Tyr Glu Asp Asn Asn Tyr Lys Ala Leu Phe Gly Gly

325

330

335

Asn Gly Thr Glu Glu Asp Ala Glu Thr Phe Tyr Asp Asn Ser Thr Thr

340

345

350

Pro Tyr Cys Asn Asp Leu Met Lys Asn Leu Glu Ser Ser Pro Leu Ser

355

360

365

Arg Ile Ile Trp Lys Ala Leu Lys Pro Leu Leu Val Gly Lys Ile Leu

370

375

380

Tyr Thr Pro Asp Thr Pro Ala Thr Arg Gln Val Met Ala Glu Val Asn

385

390

395

400

Lys Thr Phe Gln Glu Leu Ala Val Phe His Asp Leu Glu Gly Met Trp

405

410

415

Glu Glu Leu Ser Pro Lys Ile Trp Thr Phe Met Glu Asn Ser Gln Glu

420

425

430

Met Asp Leu Val Arg Met Leu Leu Asp Ser Arg Asp Asn Asp His Phe

435

440

445

Trp Glu Gln Gln Leu Asp Gly Leu Asp Trp Thr Ala Gln Asp Ile Val

450

455

460

Ala Phe Leu Ala Lys His Pro Glu Asp Val Gln Ser Ser Asn Gly Ser

465

470

475

480

Val Tyr Thr Trp Arg Glu Ala Phe Asn Glu Thr Asn Gln Ala Ile Arg

485

490

495

Thr Ile Ser Arg Phe Met Glu Cys Val Asn Leu Asn Lys Leu Glu Pro

500

505

510

Ile Ala Thr Glu Val Trp Leu Ile Asn Lys Ser Met Glu Leu Leu Asp

515

520

525

Glu Arg Lys Phe Trp Ala Gly Ile Val Phe Thr Gly Ile Thr Pro Gly

530

535

540

Ser Ile Glu Leu Pro His His Val Lys Tyr Lys Ile Arg Met Asp Ile

545

550

555

560

Asp Asn Val Glu Arg Thr Asn Lys Ile Lys Asp Gly Tyr Trp Asp Pro

565

570

575

Gly Pro Arg Ala Asp Pro Phe Glu Asp Met Trp Tyr Val Trp Gly Gly

580

585

590

Phe Ala Tyr Leu Gln Asp Val Val Glu Gln Ala Ile Ile Arg Val Leu

595

600

605

Thr Gly Thr Glu Lys Lys Thr Gly Val Tyr Met Gln Gln Met Pro Tyr

610

615

620

Pro Cys Tyr Val Asp Asp Ile Phe Leu Arg Val Met Ser Arg Ser Met

625

630

635

640

Pro Leu Phe Met Thr Leu Ala Trp Ile Tyr Ser Val Ala Val Ile Ile

645

650

655

Lys Gly Ile Val Tyr Glu Lys Glu Ala Arg Leu Lys Glu Thr Met Arg

660

665

670

Ile Met Gly Leu Asp Asn Ser Ile Leu Trp Phe Ser Trp Phe Ile Ser

675

680

685

Ser Leu Ile Pro Leu Leu Val Ser Ala Gly Leu Leu Val Val Ile Leu

690

695

700

Lys Leu Gly Asn Leu Leu Pro Tyr Ser Asp Pro Ser Val Val Phe Val
705 710 715 720

Phe Leu Ser Val Phe Ala Val Val Thr Ile Leu Gln Cys Phe Leu Ile
725 730 735

Ser Thr Leu Phe Ser Arg Ala Asn Leu Ala Ala Ala Cys Gly Gly Ile
740 745 750

Ile Tyr Phe Thr Leu Tyr Leu Pro Tyr Val Leu Cys Val Ala Trp Gln
755 760 765

Asp Tyr Val Gly Phe Thr Leu Lys Ile Phe Ala Ser Leu Leu Ser Pro
770 775 780

Val Ala Phe Gly Phe Gly Cys Glu Tyr Phe Ala Leu Phe Glu Glu Gln
785 790 795 800

Gly Ile Gly Val Gln Trp Asp Asn Leu Phe Glu Ser Pro Val Glu Glu
805 810 815

Asp Gly Phe Asn Leu Thr Thr Ser Ile Ser Met Met Leu Phe Asp Thr
820 825 830

Phe Leu Tyr Gly Val Met Thr Trp Tyr Ile Glu Ala Val Phe Pro Gly
835 840 845

Gln Tyr Gly Ile Pro Arg Pro Trp Tyr Phe Pro Cys Thr Lys Ser Tyr
850 855 860

Trp Phe Gly Glu Glu Ser Asp Glu Lys Ser His Pro Gly Ser Asn Gln
865 870 875 880

Lys Arg Met Ser Glu Ile Cys Met Glu Glu Glu Pro Thr His Leu Lys
885 890 895

Leu Gly Val Ser Ile Gln Asn Leu Val Lys Val Tyr Arg Asp Gly Met
900 905 910

Lys Val Ala Val Asp Gly Leu Ala Leu Asn Phe Tyr Glu Gly Gln Ile
915 920 925

Thr Ser Phe Leu Gly His Asn Gly Ala Gly Lys Thr Thr Thr Met Ser
930 935 940

Ile Leu Thr Gly Leu Phe Pro Pro Thr Ser Gly Thr Ala Tyr Ile Leu
945 950 955 960

Gly Lys Asp Ile Arg Ser Glu Met Ser Thr Ile Arg Gln Asn Leu Gly
965 970 975

Val Cys Pro Gln His Asn Val Leu Phe Asp Met Leu Thr Val Glu Glu
980 985 990

His Ile Trp Phe Tyr Ala Arg Leu Lys Gly Leu Ser Glu Lys His Val
995 1000 1005

Lys Ala Glu Met Glu Gln Met Ala Leu Asp Val Gly Leu Pro Ser Ser

1010

1015

1020

Lys Leu Lys Ser Lys Thr Ser Gln Leu Ser Gly Gly Met Gln Arg Lys

1025

1030

1035

1040

Leu Ser Val Ala Leu Ala Phe Val Gly Gly Ser Lys Val Val Ile Leu

1045

1050

1055

Asp Glu Pro Thr Ala Gly Val Asp Pro Tyr Ser Arg Arg Gly Ile Trp

1060

1065

1070

Glu Leu Leu Leu Lys Tyr Arg Gln Gly Arg Thr Ile Ile Leu Ser Thr

1075

1080

1085

His His Met Asp Glu Ala Asp Val Leu Gly Asp Arg Ile Ala Ile Ile

1090

1095

1100

Ser His Gly Lys Leu Cys Cys Val Gly Ser Ser Leu Phe Leu Lys Asn

1105

1110

1115

1120

Gln Leu Gly Thr Gly Tyr Tyr Leu Thr Leu Val Lys Lys Asp Val Glu

1125

1130

1135

Ser Ser Leu Ser Ser Cys Arg Asn Ser Ser Ser Thr Val Ser Tyr Leu

1140

1145

1150

Lys Lys Glu Asp Ser Val Ser Gln Ser Ser Ser Asp Ala Gly Leu Gly

1155

1160

1165

Ser Asp His Glu Ser Asp Thr Leu Thr Ile Asp Val Ser Ala Ile Ser

1170

1175

1180

Asn Leu Ile Arg Lys His Val Ser Glu Ala Arg Leu Val Glu Asp Ile

1185

1190

1195

1200

Gly His Glu Leu Thr Tyr Val Leu Pro Tyr Glu Ala Ala Lys Glu Gly

1205

1210

1215

Ala Phe Val Glu Leu Phe His Glu Ile Asp Asp Arg Leu Ser Asp Leu

1220

1225

1230

Gly Ile Ser Ser Tyr Gly Ile Ser Glu Thr Thr Leu Glu Glu Ile Phe

1235

1240

1245

Leu Lys Val Ala Glu Glu Ser Gly Val Asp Ala Glu Thr Ser Asp Gly

1250

1255

1260

Thr Leu Pro Ala Arg Arg Asn Arg Arg Ala Phe Gly Asp Lys Gln Ser

1265

1270

1275

1280

Cys Leu Arg Pro Phe Thr Glu Asp Asp Ala Ala Asp Pro Asn Asp Ser

1285

1290

1295

Asp Ile Asp Pro Glu Ser Arg Glu Thr Asp Leu Leu Ser Gly Met Asp

1300

1305

1310

Gly Lys Gly Ser Tyr Gln Val Lys Gly Trp Lys Leu Thr Gln Gln Gln
1315 1320 1325

Phe Val Ala Leu Leu Trp Lys Arg Leu Leu Ile Ala Arg Arg Ser Arg
1330 1335 1340

Lys Gly Phe Phe Ala Gln Ile Val Leu Pro Ala Val Phe Val Cys Ile
1345 1350 1355 1360

Ala Leu Val Phe Ser Leu Ile Val Pro Pro Phe Gly Lys Tyr Pro Ser
1365 1370 1375

Leu Glu Leu Gln Pro Trp Met Tyr Asn Glu Gln Tyr Thr Phe Val Ser
1380 1385 1390

Asn Asp Ala Pro Glu Asp Thr Gly Thr Leu Glu Leu Leu Asn Ala Leu
1395 1400 1405

Thr Lys Asp Pro Gly Phe Gly Thr Arg Cys Met Glu Gly Asn Pro Ile
1410 1415 1420

Pro Asp Thr Pro Cys Gln Ala Gly Glu Glu Glu Trp Thr Thr Ala Pro
1425 1430 1435 1440

Val Pro Gln Thr Ile Met Asp Leu Phe Gln Asn Gly Asn Trp Thr Met
1445 1450 1455

Gln Asn Pro Ser Pro Ala Cys Gln Cys Ser Ser Asp Lys Ile Lys Lys
1460 1465 1470

Met Leu Pro Val Cys Pro Pro Gly Ala Gly Gly Leu Pro Pro Pro Gln

1475

1480

1485

Arg Lys Gln Asn Thr Ala Asp Ile Leu Gln Asp Leu Thr Gly Arg Asn

1490

1495

1500

Ile Ser Asp Tyr Leu Val Lys Thr Tyr Val Gln Ile Ile Ala Lys Ser

1505

1510

1515

1520

Leu Lys Asn Lys Ile Trp Val Asn Glu Phe Arg Tyr Gly Gly Phe Ser

1525

1530

1535

Leu Gly Val Ser Asn Thr Gln Ala Leu Pro Pro Ser Gln Glu Val Asn

1540

1545

1550

Asp Ala Ile Lys Gln Met Lys Lys His Leu Lys Leu Ala Lys Asp Ser

1555

1560

1565

Ser Ala Asp Arg Phe Leu Asn Ser Leu Gly Arg Phe Met Thr Gly Leu

1570

1575

1580

Asp Thr Arg Asn Asn Val Lys Val Trp Phe Asn Asn Lys Gly Trp His

1585

1590

1595

1600

Ala Ile Ser Ser Phe Leu Asn Val Ile Asn Asn Ala Ile Leu Arg Ala

1605

1610

1615

Asn Leu Gln Lys Gly Glu Asn Pro Ser His Tyr Gly Ile Thr Ala Phe
1620 1625 1630

Asn His Pro Leu Asn Leu Thr Lys Gln Gln Leu Ser Glu Val Ala Leu
1635 1640 1645

Met Thr Thr Ser Val Asp Val Leu Val Ser Ile Cys Val Ile Phe Ala
1650 1655 1660

Met Ser Phe Val Pro Ala Ser Phe Val Val Phe Leu Ile Gln Glu Arg
1665 1670 1675 1680

Val Ser Lys Ala Lys His Leu Gln Phe Ile Ser Gly Val Lys Pro Val
1685 1690 1695

Ile Tyr Trp Leu Ser Asn Phe Val Trp Asp Met Cys Asn Tyr Val Val
1700 1705 1710

Pro Ala Thr Leu Val Ile Ile Ile Phe Ile Cys Phe Gln Gln Lys Ser
1715 1720 1725

Tyr Val Ser Ser Thr Asn Leu Pro Val Leu Ala Leu Leu Leu Leu
1730 1735 1740

Tyr Gly Trp Ser Ile Thr Pro Leu Met Tyr Pro Ala Ser Phe Val Phe
1745 1750 1755 1760

Lys Ile Pro Ser Thr Ala Tyr Val Val Leu Thr Ser Val Asn Leu Phe
1765 1770 1775

Ile Gly Ile Asn Gly Ser Val Ala Thr Phe Val Leu Glu Leu Phe Thr

1780

1785

1790

Asp Asn Lys Leu Asn Asn Ile Asn Asp Ile Leu Lys Ser Val Phe Leu

1795

1800

1805

Ile Phe Pro His Phe Cys Leu Gly Arg Gly Leu Ile Asp Met Val Lys

1810

1815

1820

Asn Gln Ala Met Ala Asp Ala Leu Glu Arg Phe Gly Glu Asn Arg Phe

1825

1830

1835

1840

Val Ser Pro Leu Ser Trp Asp Leu Val Gly Arg Asn Leu Phe Ala Met

1845

1850

1855

Ala Val Glu Gly Val Val Phe Phe Leu Ile Thr Val Leu Ile Gln Tyr

1860

1865

1870

Arg Phe Phe Ile Arg Pro Arg Pro Val Asn Ala Lys Leu Ser Pro Leu

1875

1880

1885

Asn Asp Glu Asp Glu Asp Val Arg Arg Glu Arg Gln Arg Ile Leu Asp

1890

1895

1900

Gly Gly Gly Gln Asn Asp Ile Leu Glu Ile Lys Glu Leu Thr Lys Ile

1905

1910

1915

1920

Tyr Arg Arg Lys Arg Lys Pro Ala Val Asp Arg Ile Cys Val Gly Ile

1925

1930

1935

Pro Pro Gly Glu Cys Phe Gly Leu Leu Gly Val Asn Gly Ala Gly Lys

1940

1945

1950

Ser Ser Thr Phe Lys Met Leu Thr Gly Asp Thr Thr Val Thr Arg Gly

1955

1960

1965

Asp Ala Phe Leu Asn Lys Asn Ser Ile Leu Ser Asn Ile His Glu Val

1970

1975

1980

His Gln Asn Met Gly Tyr Cys Pro Gln Phe Asp Ala Ile Thr Glu Leu

1985

1990

1995

2000

Leu Thr Gly Arg Glu His Val Glu Phe Phe Ala Leu Leu Arg Gly Val

2005

2010

2015

Pro Glu Lys Glu Val Gly Lys Val Gly Glu Trp Ala Ile Arg Lys Leu

2020

2025

2030

Gly Leu Val Lys Tyr Gly Glu Lys Tyr Ala Gly Asn Tyr Ser Gly Gly

2035

2040

2045

Asn Lys Arg Lys Leu Ser Thr Ala Met Ala Leu Ile Gly Gly Pro Pro

2050

2055

2060

Val Val Phe Leu Asp Glu Pro Thr Thr Gly Met Asp Pro Lys Ala Arg

2065

2070

2075

2080

Arg Phe Leu Trp Asn Cys Ala Leu Ser Val Val Lys Glu Gly Arg Ser

2085

2090

2095

Val Val Leu Thr Ser His Ser Met Glu Glu Cys Glu Ala Leu Cys Thr

2100

2105

2110

Arg Met Ala Ile Met Val Asn Gly Arg Phe Arg Cys Leu Gly Ser Val

2115

2120

2125

Gln His Leu Lys Asn Arg Phe Gly Asp Gly Tyr Thr Ile Val Val Arg

2130

2135

2140

Ile Ala Gly Ser Asn Pro Asp Leu Lys Pro Val Gln Asp Phe Phe Gly

2145

2150

2155

2160

Leu Ala Phe Pro Gly Ser Val Leu Lys Glu Lys His Arg Asn Met Leu

2165

2170

2175

Gln Tyr Gln Leu Pro Ser Ser Leu Ser Ser Leu Ala Arg Ile Phe Ser

2180

2185

2190

Ile Leu Ser Gln Ser Lys Lys Arg Leu His Ile Glu Asp Tyr Ser Val

2195

2200

2205

Ser Gln Thr Thr Leu Asp Gln Val Phe Val Asn Phe Ala Lys Asp Gln

2210

2215

2220

Ser Asp Asp Asp His Leu Lys Asp Leu Ser Leu His Lys Asn Gln Thr

2225

2230

2235

2240

Val Val Asp Val Ala Val Leu Thr Ser Phe Leu Gln Asp Glu Lys Val

2245

2250

2255

Lys Glu Ser Tyr Val

2260

<210> 11

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<220>

<220>

<220>

<223> Description of Artificial Sequence: ABC1

amplification primer

<400> 11

cctctcatta cacaaaaacc agac

24

<210> 12

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<220>

<220>

<220>

<223> Description of Artificial Sequence: ABC1
amplification primer

<400> 12

gctttcttttc acttctcatc ctg

23

<210> 13

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1 RT-PCR
primer

<400> 13

tccttggggtt caggggatta tc

22

<210> 14

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1 RT-PCR
primer

<400> 14

caatgttttt gtggcttcgg c

21

<210> 15

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1 RT-PCR
primer

<400> 15

74

agtcgagctc caaacatgctc agctgttact ggaagtggcc

40

<210> 16

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1 RT-PCR
primer

<400> 16

tctctggatt ctgggtctat gtcag

25

<210> 17

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1 RT-PCR
primer

<400> 17

gggagccttt gtggaactct ttc

23

<210> 18

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1 RT-PCR
primer

<400> 18

actggtcgac cattgaattg cattgcattg aatagtatca g

41

<210> 19

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 19

tttcctggtg gacaatgaa

19

<210> 20

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 20

agtgacatgc gacaggag

18

<210> 21

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 21

gatctggaag gcatgtgg

18

77

<210> 22

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 22

ccaggcagca ttgagctg

18

<210> 23

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 23

ggcctggaca acagcata

18

<210> 24

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 24

ggacaacctg tttgagagt

19

<210> 25

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 25

aagacgacca ccatgtca

18

<210> 26

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 26

atatgggagc tgctgctg

18

<210> 27

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 27

gggcatgagc tgacctatgt gctg

24

<210> 28

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 28

aagagactgc taattgcc

18

<210> 29

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 29

agcgacaaaa tcaagaag

18

<210> 30

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 30

tggcattgcaa tcagctct

18

<210> 31

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 31

tcctccacca atctgcct

18

<210> 32

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 32

ttcttctctca ttactggt

18

<210> 33

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 33

gatgccatca cagagctg

18

<210> 34

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 34

agtgtccagc atctaaa

17

<210> 35

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 35

caaagttcac aaatactt

18

<210> 36

<211> 19

<212> DNA

84

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 36

cttagggcac aattccaca

19

<210> 37

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 37

tgaaagttga tgattttc

18

<210> 38

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 38

tttttcacca tgtcgatga

19

<210> 39

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 39

ctccactgat gaactgc

17

<210> 40

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 40

gtttcttcat ttgtttga

18

<210> 41

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 41

agggcgtgtc tgggattg

18

<210> 42

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 42

cagaatcatt tggatcag

18

<210> 43

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 43

catcagaact gctctgag

18

<210> 44

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 44

agctggcttg ttttgcctt

19

<210> 45

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 45

tggacacgcc cagcttca

18

<210> 46

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 46

cctgccatgc cacacaca

18

<210> 47

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 47

ctcatcaccc gcagaaag

18

<210> 48

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 48

cacactccat gaagcgag

18

<210> 49

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 49

tccagataat gcgggaaa

18

<210> 50

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 50

tcaggattgg cttcagga

18

<210> 51

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1
sequencing primer

<400> 51

aagtttgagc tggatttctt g

21

<210> 52

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: beta-globin
antisense oligonucleotide

<400> 52

cctcttacct cagttacaat ttata

25

<210> 53

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: ABC1 antisense
oligonucleotide

<400> 53

catgttggtc ataggggtggg tagctc

26

<210> 54

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: beta-actin
amplification primer

<400> 54

tcacccacac tgtgccatct acga

24

<210> 55

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: beta-actin
amplification primer

<400> 55

cagcgggaacc gctcattgcc aatgg

25

<210> 56

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: sterol
response element oligonucleotide

<400> 56

tcgagtgacc gatagtaacc tctcga

26

<210> 57

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<220>

<223> Description of Artificial Sequence: mutated sterol
response element oligonucleotide

<400> 57

tcgagctgca catagtaacc tctcga

26

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number
WO 00/78971 A3

(51) International Patent Classification⁷: **C12N 15/52**,
C07K 14/705, C12Q 1/68, C07K 16/18, A61K 38/00

(21) International Application Number: PCT/US00/16591

(22) International Filing Date: 16 June 2000 (16.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/140,264 18 June 1999 (18.06.1999) US
60/153,872 14 September 1999 (14.09.1999) US
60/166,573 19 November 1999 (19.11.1999) US

(71) Applicants (for all designated States except US): CV
THERAPEUTICS, INC. [US/US]; 3172 Porter Drive,
Palo Alto, CA 94304 (US). UNIVERSITY OF WASH-
INGTON [US/US]; Office of Technology Transfer, Suite
200, 1107 N.E. Street, Seattle, WA 98195 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LAWN, Richard,

M. [US/US]; 1927 8th Avenue, San Francisco, CA 94116
(US). WADE, David [GB/US]; Apartment L, 383 Curt-
ner Avenue, Palo Alto, CA 94306 (US). ORAM, John, F.
[US/US]; 1738 N.E. Naomi Place, Seattle, WA 98195 (US).
GARVIN, Michael [US/US]; 17 White Street, San Fran-
cisco, CA 94019 (US).


(74) Agent: HUGHES, A., Blair; McDonnell Boehnen Hul-
bert & Berghoff, 32nd Floor, 300 South Wacker Drive,
Chicago, IL 60606 (US).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: ATP BINDING CASSETTE TRANSPORTER PROTEIN ABC1 POLYPEPTIDES



(57) Abstract: The present invention relates to novel ABC1 polypeptides and nucleic acid molecules encoding the same. The invention also relates to recombinant vectors, host cells, and compositions comprising ABC1 polynucleotides, as well as to methods for producing ABC1 polypeptides. The invention also relates to antibodies that bind specifically to ABC1 polypeptides. In addition, the invention relates to methods for increasing cholesterol efflux as well as to methods for increasing ABC1 expression and activity. The present invention further relates to methods for identifying compounds that modulate the expression of ABC1 and methods for detecting the comparative level of ABC1 polypeptides and polynucleotides in a mammalian subject. The present invention also provides kits and compositions suitable for screening compounds to determine the ABC1 expression modulating activity of the compound, as well as kits and compositions suitable to determine whether a compound modulates ABC1-dependent cholesterol efflux.



WO 00/78971 A3



Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(88) Date of publication of the international search report:
17 January 2002

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/16591

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/52 C07K14/705 C12Q1/68 C07K16/18 A61K38/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, EP0-Internal, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LANGMANN T, KLUCKEN J, REIL M, LIEBISCH G, LUCIANI MF, CHIMINI G, KAMINSKI WE, SCHMITZ G: "Molecular cloning of the human ATP-binding cassette transporter 1 (hABC1): evidence for sterol-dependent regulation in macrophages" BIOCHEM BIOPHYS RES COMMUN, vol. 257, no. 1, 2 April 1999 (1999-04-02), pages 29-33, XP000877240	1-3
X	cited in the application the whole document --- -/--	33,35

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 March 2001

Date of mailing of the international search report

12 07. 2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

CHAMBONNET, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/16591

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>SCHWARTZ K ET AL: "ABC1 GENE EXPRESSION AND APOA-I-MEDIATED CHOLESTEROL EFFLUX ARE REGULATED BY LXR" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 274, no. 3, 11 August 2000 (2000-08-11), pages 794-802, XP000960872 ISSN: 0006-291X the whole document -----</p>	1

INTERNATIONAL SEARCH REPORT

international application No.
PCT/US 00/16591

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 35 to 45 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see further information sheet invention 1.

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-24 33-36 38 44 70-73 and partially 37 39-43 45
74

An isolated polypeptide comprising SEQ ID NO 2 or an amino acid sequence that has at least 98% identity with it; an isolated polynucleotide selected from the group of consisting of a polynucleotide encoding said polypeptide or comprising at least nucleotides 291-7074 of SEQ ID NO 1, or that is at least 90% identical to SEQ ID NO 1, or complementary thereof; a composition comprising such a polynucleotide and a suitable carrier; a recombinant vector comprising such a polynucleotide; a composition or a host cell comprising said vector; a method for producing an ABC1 protein using said vector; methods for increasing cholesterol influx or ABC1 expression from cells of a mammalian subject using thereof; an isolated antibody that binds specifically to said isolated polypeptide and a kit for screening containing said antibody;

2. Claims: 25-27 31

An isolated polypeptide comprising SEQ ID NO 8; an isolated polynucleotide encoding said polypeptide or comprising SEQ ID NO 7; a recombinant vector comprising said polynucleotide.

3. Claims: 28-30 32

An isolated polypeptide comprising SEQ ID NO 10; an isolated polynucleotide encoding said polypeptide or comprising SEQ ID NO 9; a recombinant vector comprising said polynucleotide.

4. Claims: partially 37 39-43 45 74

A method suitable for increasing cholesterol efflux from the cells of a mammalian subject comprising the step of administering to the mammalian subject a polynucleotide encoding ABC1 in an amount sufficient to increase cholesterol efflux from said cells; as far as not covered by the previous subject.

5. Claims: 46-54

A method suitable for increasing cholesterol efflux of the cells of a mammalian subject comprising the step of

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

administering to the mammalian subject a therapeutic amount of a compound that increases the expression of ABC1 in said cells; a method suitable for increasing the expression of ABC1 in the cells of a mammalian subject comprising the step of administering to the mammalian subject a cAMP analogue in an amount sufficient to increase the expression of ABC1 in said cells; a method suitable for increasing cholesterol efflux of the cells of a mammalian subject comprising the step of administering to the mammalian subject a compound that increases ABC1 activity in said cells; as far as not covered by the previous subject;

6. Claims: 55-61 and partially 74 75 76

A method for screening a test compound to determine whether the test compound promotes ABC1-mediated cholesterol efflux from cells in culture comprising the steps of : assaying the level of cholesterol efflux in a sample of mammalian cells maintained in culture to determine a control level of cholesterol efflux; contacting the cells with the test compound being screened; assaying the level of cholesterol efflux in a sample of cells after contact with the test compound; and assaying the level of ABC1-mediated cholesterol efflux in a sample of cells after contact with the test compound, thereby determining whether the test compound promotes ABC1-mediated cholesterol efflux from cells in culture; a kit suitable for said screening method comprising an inactivating ABC1 antibody or an antisense ABC1 oligonucleotide; as far as not covered by a previous subject;

7. Claims: 62-69

A method for detecting the comparative level of ABC1 expression in the cells of a mammalian subject comprising the steps of : obtaining a cell sample from the mammalian subject; assaying the level of ABC1 mRNA expression in the cell sample with a pre-determined standard level of ABC1 mRNA expression, thereby detecting the comparative level of ABC1 expression in the cells of a mammalian subject ;

8. Claim : Partially 74

A kit suitable for screening a compound to determine whether the compound modulates ABC-1 dependent cholesterol efflux comprising an inactivating ABC1 antibody in an amount sufficient for at least one assay and instructions for use; as far as not covered by a previous subject.

9. Claim : Partially 75 and 76

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

A kit suitable for screening a compound to determine whether the compound modulates ABC1-dependent cholesterol efflux comprising an antisense ABC1 oligonucleotide in an amount sufficient for at least one assay and instructions for use; as far as not covered by a previous subject.